

Estimating Social Network Models with Missing Links*

Arthur Lewbel, Xi Qu, and Xun Tang

December 23, 2022

Abstract

We propose an adjusted 2SLS estimator for social network models when some existing network links are missing from the sample (due, e.g., to recall errors by survey respondents, or lapses in data input). In the feasible structural form, missing links make all covariates endogenous and add a new source of correlation between the structural errors and endogenous peer outcomes (in addition to simultaneity), thus invalidating conventional estimators used in the literature. We resolve these issues by rescaling peer outcomes with estimates of missing rates and constructing instruments that exploit properties of the noisy network measures. We apply our method to study peer effects in household decisions to participate in a microfinance program in Indian villages. We find that ignoring missing links and applying conventional instruments would result in a sizeable upward bias in peer effect estimates.

*We are grateful to seminar and conference participants at CalTech, Chinese University of Hong Kong, Texas Camp Econometrics, U Penn, and UT Austin for useful comments and suggestions. Lewbel and Tang are grateful for the financial support from National Science Foundation (Grant SES-1919489). Any or all remaining errors are our own.

1 Introduction

In many social and economic environments, an individual’s behavior or outcome (such as a consumption choice or a test score) depends not only on his or her own characteristics, but also on the behavior and characteristics of other individuals. Call such dependence between two individuals a *link*. A *social network* consists of a group of individuals, some of whom are linked to others. The econometrics literature on social networks has largely focused on disentangling various channels of social effects based on observed outcomes and characteristics of network members. These include identifying the effects on each individual’s outcome of (i) the individual’s own characteristics (*individual effects*), (ii) the characteristics of people linked to the individual (*contextual effects*) and (iii) the outcomes of people linked to the individual (*peer effects*). See Blume et al. (2011) and Graham (2020) for extensive surveys about identifying such effects in social network models.

A popular approach for estimating social network models is to use two-stage least squares (2SLS). This requires researchers to construct instruments for the endogenous peer outcomes, using *perfect knowledge* of the network structure, as given by the *adjacency* matrix (i.e., the matrix that lists all links in the network). See, for example, Bramoullé et al. (2009), Kelejian and Prucha (1998), Lee (2007), and Lin (2010). In practice, samples of network links are often collected from survey responses. Such samples may suffer from missing links, due, e.g., to recall errors or misunderstandings by survey respondents, or lapses in data input.

Missing links in the sample pose major methodological challenges for estimators like 2SLS. To see this, consider a data-generating process (DGP) from which a large number of independent networks (i.e., groups) are drawn. Each group has n individual members. The issues we raise and the solutions we propose also apply to other contexts, such as a large number of independent networks with different sizes or a single growing network, but are easiest to illustrate in the context of many independent, identically sized groups.

Suppose that in each group, a vector of individual outcomes $y \in \mathbb{R}^n$ is determined by a structural model:

$$y = \lambda Gy + X\beta + \varepsilon, \text{ where } E(\varepsilon|X, G) = 0.$$

In this model, the adjacency matrix, G , is an n -by- n matrix of dummy variables that describes the group’s network: the element in row j and column k of G equals one if individual j is linked to member k , and zero otherwise. Here X is an n -by- K matrix of exogenous covariates, and ε is an n -vector of structural errors. The random arrays y , G , X , and ε all vary across the groups in the sample, while the coefficients λ and β are the same across groups. We drop group subscripts for clarity.

For simplicity we have for now omitted contextual effects, i.e., a term given by $GX\gamma$ in equation (1). We also omit group-level fixed effects for now. Extensions of our results that include either or both are provided later.

The regressors in the model are GY and X . While X is exogenous, the regressors GY are correlated with ε . The issue of simultaneity arises here, because any one individual’s outcome depends on, and is determined simultaneously with, the outcomes of other group peers. A simple estimator of the peer effect λ and individual effects β that deals with this simultaneity problem is 2SLS, using GX or G^2X as instruments for GY , as in Bramoullé et al. (2009).¹

But now suppose that, in each group, a researcher does not observe G perfectly, but instead observes a noisy measure H , which differs from G by randomly missing some actual existing links while correctly reporting others. The goal now is to estimate λ and β from the “feasible” structural form:

$$y = \lambda Hy + X\beta + u, \tag{1}$$

where $u \equiv [\varepsilon + \lambda(G - H)y]$ is a vector of *composite* errors.

The missing links in H aggravate endogeneity issues in (1) in two important ways. First, they lead to correlation between X and u through $\lambda(G - H)y$, a component in u that is due to the measurement error in the adjacency matrix. As a result, unlike using GX or G^2X as instruments, 2SLS estimates based on the feasible instruments HX or H^2X would be inconsistent, due to a failure of instrument exogeneity.

Second, these missing links cause an additional source of endogeneity in Hy . Like Gy ,

¹If the model included contextual effects $GX\gamma$ in its structural form, then G^2X could be used as instruments for Gy , otherwise use of GX as instruments suffices.

the feasible Hy is correlated with the model error ε due to simultaneity. But in addition, Hy is also correlated with u through the measurement errors in $\lambda(G - H)y$. For all these reasons, standard 2SLS estimators of this model become inconsistent in the presence of missing links.²

In this paper, we introduce an *adjusted-2SLS* estimator, which resolves these challenges and consistently estimates (λ, β) using alternative valid instruments constructed from H despite the missing links. We first introduce the main idea for a benchmark case, where actual links in G are missing randomly from H in the sample at an unknown rate $p \in (0, 1)$. Later, we extend our method to allow the missing rates $p(X)$ to depend on covariates.

Our method is based on a series of new insights that have not been explored in the literature. *First*, we observe that by rescaling the noisy measure of peer outcomes Hy with the inverse probability of reporting correctly $1/(1 - p)$, we restore the exogeneity of X in a *rescaled* structural form. Formally, this means if we reparametrize (1) as

$$y = \lambda^*Hy + X\beta + v \text{ with } \lambda^* \equiv \lambda/(1 - p), \tag{2}$$

then the reparametrized errors $v \equiv \varepsilon + (\lambda G - \lambda^*H)y$ satisfy $E(v|X, G) = 0$. This holds regardless of how the actual network G is formed, as long as $E(\varepsilon|X, G) = 0$.

Second, despite the restored exogeneity of X in (2), conventional instruments such as HX or H^2X remain invalid, because the reparametrized errors v depend on H . To address this issue, we provide alternative functions of H and X that are valid instruments. For example, we show that if the true network G is symmetric, and the observed H is asymmetric with links missing independently, then $H'X$ is uncorrelated with v (where H' denotes the transpose of H). Thus, we can use $H'X$ as valid instruments in an adjusted-2SLS where peer outcomes are rescaled by $1/(1 - p)$. To the best of our knowledge, no other paper in the literature has proposed this use of $H'X$ as instruments.

For cases where the above conditions do not hold (e.g., if H is symmetric and/or the symmetry of G is unknown), we provide an alternative way to construct valid instruments,

²While we focus on the 2SLS estimator in this paper, the same arguments apply to show that conventional maximum likelihood, and the generalized least squares estimators based on (1) are also inconsistent.

based on observing two different H matrices.³ For example, in our empirical application, for an undirected link between two households A and B, we observe two proxy measures of the same link: whether A visited B, and whether B visited A. This yields two different observed H matrices corresponding to the same true undirected link in the G matrix. Observing these two H matrices allows us to construct valid instruments.

Third, under either of the two scenarios above that permit construction of valid instruments (that is, either asymmetric H with symmetric G , or observing two different H matrices regardless of symmetry), we provide simple methods to identify and estimate the unknown missing rate p .⁴

Building on these insights, we construct an adjusted 2SLS estimator for (λ, β) , and provide its limiting distribution as the number of groups grows to infinity. This estimator essentially applies 2SLS to the rescaled peer outcomes in (2), using our proposed new instruments, and a sample analog estimator for the missing rate p . The estimator is easy to implement, and we demonstrate good finite-sample performance in monte carlo simulation.

We then generalize the model and our estimator in several directions. We show how to include contextual effects (a $GX\gamma$ term) as well as group-level fixed effects into the structural form in (1). We also allow the missing rates p to be heterogeneous and depend on individual covariates in X .

Furthermore, we extend our method to the case of a single large network. In this case, the asymptotic experiment is to increase the number of individuals on a single network, rather than increasing the number of small, fixed-sized groups. For this extension we propose two possible settings where some form of weak dependence exists between the outcomes of individuals who are “sufficiently far” from each other, either in the sense of not being in the same group (Section 6.1) or in terms of a latent distance metric (Section 6.2). In either case, we show that under such weak dependence our adjusted 2SLS estimator, when pooled over individuals in the sample, still converges to the intended estimand.

³We also show yet another way to construct valid instruments is to use nonlinear functions of X .

⁴The approach we take in this step differs from, and is simpler than, other papers that use multiple measures to deal with misclassification in discrete explanatory variables (e.g. Mahajan (2006), Lewbel (2007), and Hu (2008)). This is because, for implementing our adjusted-2SLS, it is only necessary to estimate the missing rate p , rather than the distribution of outcomes conditional on the actual G .

Finally, we apply our method to estimate peer effects in household decisions to participate in a microfinance program in Indian villages, using data from Banerjee et al. (2013). We match the individual survey to the household survey there, yielding a sample of 4134 households in 43 villages in South India. The parameter of interest is the peer (endorsement) effect, which reflects how a household’s decision is influenced by the microfinance program participation of other households to which it is linked. Survey information about directed visits between the households provides two noisy measures of network links (i.e., two H matrices). We estimate missing rates in each of these two measures using our methodology, and then we apply these rates in our adjusted-2SLS procedure to estimate the endorsement peer effects.

We find that participation by another linked household increases a household’s own participation rate by around 4.6%. This effect is economically significant, compared to the average participation rate of 18.2% in the sample. We also find that ignoring the missing links in the noisy measures and applying conventional 2SLS estimation results in a sizeable upward bias in the estimates of these peer effects.

Roadmap. Section 2 reviews the related literature, and explains our contribution in its context. Section 3 specifies the model, and illustrates the main ideas in a benchmark model with independent and identical missing rates. Section 4 defines an analog estimator for missing rates, and provides our adjusted-2SLS estimator for social effects. Section 5 extends the method to more general settings with contextual effects, heterogeneous missing rates, and group fixed effects. Section 6 shows how our estimator works when the sample consists of a single, large network. Section 7 presents monte carlo simulation results. Section 8 applies our method to analyze peer effects in microfinance participation in India. Proofs are collected in the Appendix.

2 Related Literature

Missing or misclassified links is an important topic in the social networks literature. Shalizi and Rinaldo (2013) note the challenge of dealing with missing network links in Random

Graph Models. Advani and Malde (2018) show that even a relatively low misreporting rate can lead to large bias in causal effect estimates.

The econometrics literature on the estimation of peer effects with network measurement issues is fast growing. Butts (2003) proposes a hierarchical Bayesian model to infer social structure in the presence of measurement errors. Chandrasekhar and Lewis (2011) show how egocentrically sampled network data can be used to predict the full network in a graphical reconstruction process. Liu (2013) shows that when the adjacency matrix is not row-normalized, instrumental variable estimators based on an out-degree distribution can be valid.

Goldsmith-Pinkham and Imbens (2013) examine network endogeneity and investigate simultaneously alternative definitions of links and the possibility of peer effects arising through multiple networks. They explicitly model network formation, with estimation based on maximum likelihood, using a Bayesian approach for computational convenience and feasibility. Hardy et al. (2019) estimate treatment effects on a social network when the reported links are a noisy representation of true spillover pathways. They use a mixture model that accounts for missing links as unobserved network heterogeneity, and estimate it using an Expectation-Maximization algorithm. This approach requires a parametric model of how links are determined and treatment is assigned, and requires enumerating the likelihood conditional on all possible treatment exposures (which in turn depends on the latent unobserved network).

In contrast with these papers above, we focus on social effect parameters in a linear social network model, and exploit implications of randomly missing links for identification. Our method does not require modeling link formation. Our estimator is essentially a rescaled 2SLS, which has closed form and is easy to compute.

Boucher and Houndetoungan (2020) estimate peer effects when the social networks in the sample are subject to measurement issues, such as missing or misclassified links. Their method can be applied when the researcher only has access to aggregated relational data, but assumes the researcher knows, or has a consistent estimator of, the distribution of the actual network. They construct instruments by drawing from this distribution, and use

2SLS to estimate the peer effects. In comparison, the method we propose does not require such prior knowledge or estimates of network distribution.

Griffith (2021) studies the case where links are censored in the sample (e.g., when each individual is restricted to naming 5 or fewer links with other people, even if the actual number of people the individual is linked with is larger). Griffith (2021) analytically characterizes the bias in a reduced-form regression (i.e., when the outcome vector y is regressed on the exogenous variables X and GX). In addition, for a model with *no* endogenous peer effect, Griffith (2021) shows that the bias can be consistently estimated under an order invariance condition (i.e., the covariance of characteristics of those one is linked with is invariant to the order in which links are reported or censored). In comparison, we consider different settings where links are missing at random in a model with a non-zero endogenous peer effect $\lambda \neq 0$. (This is later generalized to the case with heterogeneous missing rates.) We show that the 2SLS estimand in this case contains a simple augmentation bias in peer effects (in the sense of converging to $\lambda/(1-p)$, with p being the missing rate), and no bias in other individual effects. Bias correction in our case is immediate once the missing rate is estimated using a simple approach that we provide.

3 Model and Identification

Consider a DGP from which a large number of small, independent networks (groups) are drawn. Each group s consists of n_s individual members, with $n_s \geq 3$ being finite integers. In Section 3-5, we identify and estimate a linear social network model with missing links in the data as the number of groups in the sample approaches infinity. Later we consider the extension to a single growing network.

To simplify exposition, let the group sizes $n_s = n$ be fixed across groups $s = 1, \dots, S$. This allows us to drop the group subscript s while presenting our identification argument. We will later add back these group subscripts and allow for variation in group sizes when we define our estimator in Section 4.

The structural form for the n -vector of individual outcomes y in each group is:

$$y = \lambda Gy + X\beta + \varepsilon, \quad (3)$$

where the peer effect λ and the K -vector of direct effects β are constant parameters of interest, X is an n -by- K matrix of individual- or group-level explanatory variables, and $G \in \{0, 1\}^{n \times n}$ is a network (adjacency) matrix with its (i, j) -th entry $G_{ij} = 1$ if and only if the individual members i and j are linked.

Note that, like y , X , and ε , the adjacency matrix G varies by group, and so it too has an s subscript that has been dropped for now. Only the coefficients λ and β are constants that do not vary across groups. Assume that $(I - \lambda G)$ is invertible almost surely. (A sufficient condition for this is that $\|\lambda G\| < 1$ for *any* matrix norm $\|\cdot\|$.) Solving equation (3) for y gives the reduced form for outcomes:

$$y = M(X\beta + \varepsilon), \text{ where } M \equiv (I - \lambda G)^{-1}. \quad (4)$$

For each group, the sample only reports a noisy measure of the adjacency matrix G , with randomly missing links. Denote this noisy measure by $H \in \{0, 1\}^{n \times n}$. Let $G_{ii} = 0$ and $H_{ii} = 0$ by convention. Assume:

$$(A1) \ E(H_{ij}|G, X) = E(H_{ij}|G_{ij}, X) \text{ for all } i \text{ and } j;$$

$$(A2) \ E(H_{ij}|G_{ij} = 1, X) = 1 - p \text{ and } E(H_{ij}|G_{ij} = 0, X) = 0 \text{ for all } i \neq j;$$

$$(A3) \ E(\varepsilon|X, G, H) = 0.$$

Condition (A1) states that the incidence of missing a link between two individual members i and j is conditionally independent from the state of links involving other individuals $l \notin \{i, j\}$. Condition (A2) specifies that misclassification of links is one-sided in that existent links are missing from the sample at a rate of $p \in (0, 1)$ while non-existent links are never mistakenly coded as existent. Condition (A3) rules out endogeneity in link

formation, by assuming that (X, G, H) are exogenous to the structural error ε .

Under (A1) and (A2), we can write:

$$E(H|G, X) = (1 - p)G. \tag{5}$$

In the next subsection we show how this property, along with condition (A3), leads to a simple expression for the 2SLS estimand despite missing links.

3.1 Augmentation bias in two-stage least squares

In place of equation (1), we write a feasible structural form using H instead of G as:

$$y = \lambda^*Hy + X\beta + \underbrace{\varepsilon + (\lambda G - \lambda^*H)y}_{\equiv v}, \text{ where } \lambda^* \equiv \lambda/(1 - p). \tag{6}$$

Note the peer effect λ is replaced with a rescaled version λ^* in (6). Lemma 1 shows how this replacement restores the exogeneity of X with the new composite error v in (6).

Lemma 1. *Under (A1), (A2), and (A3), $E(v|X, G) = 0$.*

Lemma 1 may seem rather surprising *ex ante*, because one would expect (X, G) to be generically correlated with the composite error v which depends on y . The intuition for this result is as follows. Once we condition on the actual network G and explanatory variables in X , the randomness in individual outcomes y is solely due to the actual structural errors ε , which are uncorrelated with both X and (H, G) under (A3). As a result, any potential correlation between v and (X, G) could only be due to the reparametrized measurement error $\lambda G - \lambda^*H$. But equation (5) implies that $\lambda G - \lambda^*H$, and consequently the reparametrized error v , are mean independent from (X, G) .

As discussed earlier, even with Lemma 1 establishing exogeneity of X in (6) by replacing λ with λ^* , there is still endogeneity in the term Hy because $E[(Hy)'v] \neq 0$ in general.⁵ We therefore next investigate the estimand from 2SLS given appropriate instruments for Hy .

⁵To see this, note $E(H'H|G, X) \neq (1 - p)^2G'G = (1 - p)E(H'G|G, X)$ under (A1), (A2) and even with the addition of a stronger condition (A4) in Section 3.2. It then follows from (A3) and an application of the law of iterated expectation that $E(y'H'Hy) \neq (1 - p)E(y'H'Gy)$ in general.

Based on Lemma 1 nonlinear functions of X can serve as instruments, if the usual rank (instrument relevance) condition is satisfied. However, nonlinear functions of X might not be relevant, or might be weak as instruments, since the structural model is linear in X . To deal with this possibility, we later show that it is also possible to use the noisy network measure H to construct instruments just from linear functions of X . One example we give later in Section 3.2 is that $H'X$ can be a valid instrument, meaning $E[(H'X)'v] = 0$, when H consists of directed links.

More generally, let ζ be a generic n -by- L matrix of instruments for Hy . This may either be nonlinear functions of explanatory variables $\zeta(X)$ if these are related to the link formation in G , or functions of the noisy network measure H such as $\zeta(X, H) = H'X$ for asymmetric H as discussed later in Section 3.2. Denote $R \equiv (Hy, X)$, $Z \equiv (\zeta, X)$ so that $E(R'v) \neq 0$ while $E(Z'v) = 0$. Assume instruments satisfy the following rank condition:

$$(IV-R) \ E(Z'R) \text{ and } E(Z'Z) \text{ have full rank.}$$

Let $\Pi \equiv [E(Z'Z)]^{-1} E(Z'R)$. By (6) and Lemma 1,

$$\begin{aligned} \Pi' E(Z'y) &= \Pi' E(Z'R)(\lambda^*, \beta')' + \Pi' E(Z'v) \\ \Rightarrow (\lambda^*, \beta')' &= [\Pi' E(Z'R)]^{-1} [\Pi' E(Z'y)]. \end{aligned} \tag{7}$$

We formalize this result in the next proposition.

Proposition 1. *Suppose (A1), (A2), and (A3) hold, and that (IV-R) holds for instruments Z . The two-stage least-squares estimand using Z for (6) is then $(\lambda^*, \beta')'$.*

Proposition 1 shows that when links are missing at random in the sample, 2SLS estimation using valid instruments leads to *augmentation bias* in the peer effect, because 2SLS estimates λ^* instead of λ . Intuitively, when links are missing in the sample, their contribution to peer effects are erroneously attributed to the remaining observed links, thereby exaggerating the magnitude of peer effects attributed to the observed links. In contrast to peer effects, the individual effects β are consistently estimated by 2SLS (with valid instruments) despite missing links.

Based on Proposition 1, we have two main requirements for estimating the model. First, we need to construct valid instruments for 2SLS, and second, we need an estimator of p , to convert the 2SLS estimate of λ^* into an estimate of λ .

3.2 Constructing instruments from a noisy network measure

We return to the question about how to construct instruments using a noisy network measure H . Assume:

(A4) Conditional on (G, X) , H_{ij} and H_{kl} are independent whenever $(i, j) \neq (k, l)$.

This condition states the incidence of missing two links that do not involve the same individual are independent conditional on actual link status. This rules out the case where H and G are both symmetric so $H_{ik} = H_{ki}$ and $G_{ik} = G_{ki}$ for all i, k .⁶ We later give a method for constructing instruments in the symmetric matrix case of undirected links. With (A4), we can construct instruments using H and X as follows.

Proposition 2. *Suppose (A1), (A2), (A3), and (A4) hold. Then $E(Z'v) = 0$, where $Z \equiv (H'X, X)$.*

There is a simple interpretation of the instruments $H'X$: the i -th component (row) of $H'X$ is the sum of characteristics of all individuals who are observed to report links with i .

Recall that GX are valid instruments when G is perfectly observed. Therefore, one may wonder why we use $H'X$ instead of HX as instruments here. To understand this, note the composite error v in (6) contains the reparametrized measurement error $(\lambda G - \lambda^* H)$, and so in particular contains H . Hence, even under (A1)-(A4), HX is correlated with this reparametrized measurement error in v through H . In contrast, using a *transpose* of H in $H'X$ removes such correlation, because under (A4) the events of missing links between different pairs of (i, j) are conditionally independent.⁷ Therefore, $H'X$ are valid instruments

⁶Suppose $(H_{ik}, G_{ik}) = (H_{ki}, G_{ki})$. Under (A1)-(A2), $E(H_{ik}|G, X)E(H_{ki}|G, X) = (1-p)^2 G_{ik} \neq (1-p)G_{ik} = E(H_{ik}H_{ki}|G, X)$. Hence (A4) does not hold.

⁷Formally, $(HX)'v$ contains $H'H$ (and consequently H_{ik}^2 terms), while $(H'X)'v$ contains H^2 (and consequently $H_{ik}H_{ki}$ terms) instead. Under (A4), $E(H^2|X, G) = (1-p)^2 G^2$. This equality, along with

while HX are not.

To apply 2SLS, the instruments need to satisfy the rank conditions in (IV-R). The next proposition specifies sufficient conditions for these rank conditions in terms of moments of functions of (X, G) .

Proposition 3. *Suppose (A1), (A2), (A3), (A4) hold, and $E(X'X)$ is non-singular. Then (IV-R) holds for $Z \equiv (H'X, X)$ if*

$$\begin{pmatrix} E(X'X) & E(X'M^{-1}X) \\ E(X'MX) & E(X'X) \end{pmatrix} \text{ and } \begin{pmatrix} E(X'G^2X) & E(X'GX) \\ E(X'GX) & E(X'X) \end{pmatrix} \text{ are non-singular.} \quad (8)$$

The rank conditions in (8) hold generically for random link formation models. Our simulations show that these conditions hold even for very restrictive cases where links are i.i.d. Bernoulli and independent from X . Violations of these conditions in (8) do exist in special cases. One such example is the linear-in-means social interactions model where G^k is proportional to a square matrix of ones for all positive integers k . It is worth noting that such an example of linear-in-means model also violates the rank condition for identifying social effects in Bramoullé et al. (2009), which requires I , G , and G^2 be linearly independent.

3.3 Instruments based on multiple symmetric measures

The method in Section 3.2 to construct instruments assumes we observe an asymmetric network measure matrix H . In this section we show that we can alternatively construct instruments if the sample provides two (or more) symmetric measures of the network. Call these two measures $H^{(1)}$ and $H^{(2)}$.

For example, Banerjee et al. (2013) provide multiple measures of undirected links between households in rural villages across the State of Karnataka, India. For each pair of households, the survey asks which households you visited, and which ones visited you. Banerjee et al. (2013) symmetrize each of these two measures, yielding symmetric matrices

the fact that $E(HG|X, G) = (1 - p)G^2$ under (A1)-(A2), implies $E[(H'X)'v|G, X] = 0$. In contrast, $E[H'H|X, G] \neq (1 - p)^2G^2$, and as a result, $E[(HX)'v|G, X] \neq 0$.

we can call $H^{(1)}$ and $H^{(2)}$. These two matrices are both measures of the same underlying symmetric network G . However, as we show later, these two matrices empirically differ substantially, indicating that they are different noisy measures of G .

Assume we observe symmetric matrices $H^{(1)}$ and $H^{(2)}$ satisfying (A1), (A2), (A3), and

(A4') Conditional on (G, X) , $H_{ij}^{(1)}$ and $H_{kl}^{(2)}$ are independent whenever $(i, j) \neq (k, l)$.

Note that $H^{(1)}$ and $H^{(2)}$ can each have their own, different missing link rates $p^{(1)}$ and $p^{(2)}$. Using either measure $H^{(1)}$ or $H^{(2)}$, we can construct a feasible structural form. That is, for $t = 1, 2$,

$$y = \frac{\lambda}{1-p^{(t)}} H^{(t)} y + X\beta + v^{(t)}, \text{ where } v^{(t)} = \varepsilon + \lambda \left[G - \frac{H^{(t)}}{1-p^{(t)}} \right] y. \quad (9)$$

Under (A1)-(A3) and (A4') and by an argument similar to Proposition 2, we can show that $H^{(2)}X$ satisfies exogenous conditions with regard to $v^{(1)}$ (see Appendix A for details):

$$E \left[(H^{(2)}X)' v^{(1)} \right] = 0.$$

By a symmetric argument, similar exogeneity holds for $H^{(1)}X$ and $v^{(2)}$. We can therefore use $H^{(1)}X$ as instruments in equation (9) with $t = 2$, and $H^{(2)}X$ as instruments in (9) with $t = 1$. In Section 4, we discuss how to construct 2SLS estimators using these multiple network measures.

3.4 Recovering peer effects and missing rates

To remove the augmentation bias and recover the peer effect λ from the 2SLS estimated λ^* , we must identify and estimate the unknown missing link rate p . Here we provide two different methods for identifying p under two different scenarios.

First, consider a scenario where the actual network G is symmetric (meaning links are undirected, so $G_{ij} = G_{ji}$ with probability one for all $i \neq j$), but suppose the sample reports a noisy *asymmetric* H (meaning that $H_{ij} \neq H_{ji}$ with some positive probability). For example, the sample may collect self-reported survey responses about undirected links,

with individual i reporting H_{ik} for $k \neq i$ and individual j reporting $H_{jk'}$ for $k' \neq j$.

In this scenario, we can construct a symmetric measure \tilde{H} given by elements $\tilde{H}_{ij} = \max\{H_{ij}, H_{ji}\}$. By construction, given our assumptions, if the missing link rate for H is p , then the missing link rate for \tilde{H} will be p^2 . Let $\psi(H) \in \mathbb{R}$ denote the average of all off-diagonal components in a network measure H . By the implication of randomly missing links in (5) and the linearity of $\psi(\cdot)$,

$$E[\psi(H)] = (1 - p)E[\psi(G)] \text{ and } E[\psi(\tilde{H})] = (1 - p^2)E[\psi(G)].$$

Hence we can identify the missing rate as $p = E[\psi(\tilde{H})]/E[\psi(H)] - 1$.

Next, consider a different scenario where the sample has two independent, noisy measures of the adjacency matrix, $H^{(1)}$ and $H^{(2)}$, with unknown missing rates $p^{(1)}$ and $p^{(2)}$ respectively. Construct a third measure $H^{(3)}_{ij} = \max\{H^{(1)}_{ij}, H^{(2)}_{ij}\}$. The implied missing rates in $H^{(3)}$ is $p^{(3)} = p^{(1)} \times p^{(2)}$. By equation (5) we have

$$E[H^{(t)}] = (1 - p^{(t)})E(G) \text{ for } t = 1, 2, 3.$$

By the linearity of ψ , we therefore get $E[\psi(H^{(t)})] = (1 - p^{(t)})E[\psi(G)]$, with $E[\psi(G)] \neq 0$, for $t = 1, 2, 3$. Hence we can identify the missing rates $p^{(1)}$ and $p^{(2)}$ by

$$p^{(1)} = \frac{E[\psi(H^{(3)})] - E[\psi(H^{(1)})]}{E[\psi(H^{(2)})]} \text{ and } p^{(2)} = \frac{E[\psi(H^{(3)})] - E[\psi(H^{(2)})]}{E[\psi(H^{(1)})]}.$$

Once the missing rates are recovered, we can use them to remove the augmentation bias in the 2SLS estimand in (6). Equivalently, we can use these rates to rescale the endogenous variable as $Hy/(1 - p)$ so that 2SLS can then estimate $(\lambda, \beta)'$ consistently.

In each of the above scenarios, the matrix we construct, either \tilde{H} or $H^{(3)}$, is under our assumptions a more accurate measure of G than the original H or $H^{(1)}$ and $H^{(2)}$, in the sense of having a lower rate of missing links. However, direct estimation using these constructed matrices in place of G would still be biased and inconsistent due to the missing links. The estimators we propose in the next section don't directly use these constructed

matrices (other than to estimate missing rates as above), but the estimators do make use of the information involved in such construction. Specifically, estimation in the first scenario make use of both H and its transpose that were used to construct \tilde{H} , while estimation in the second scenario make use of both $H^{(1)}$ and $H^{(2)}$ that were used to construct $H^{(3)}$.

4 Two-Step Estimation

The previous section provides two different identification strategies, based on either observing an asymmetric noisy adjacency matrix measure, or observing two noisy adjacency matrix measures. We now propose estimators based on each of these identification strategies. Consider a sample of S independent groups, indexed by $s = 1, 2, \dots, S$, with group s consisting of n_s members (later we consider extensions to a single growing network instead of many independent groups). For each group s , the sample reports an n_s -by-1 vector of individual outcomes y_s , an n_s -by- K matrix of explanatory variables X_s , and either an n_s -by- n_s noisy asymmetric network measure H_s , or two symmetric n_s -by- n_s noisy measures $H_s^{(1)}$ and $H_s^{(2)}$.

Consider the first scenario in Section 3.4, which has symmetric G_s and asymmetric H_s . We begin by estimating the missing rate p . Let \tilde{H}_s denote the symmetric measure with its (i, j) -th component constructed as $\tilde{H}_{s,ij} = \max\{H_{s,ij}, H_{s,ji}\}$. Define:

$$\psi_s \equiv \psi(H_s) \text{ and } \tilde{\psi}_s \equiv \psi(\tilde{H}_s).$$

We estimate the missing rate p by

$$\hat{p} = \frac{\frac{1}{S} \sum_{s=1}^S \tilde{\psi}_s}{\frac{1}{S} \sum_{s=1}^S \psi_s} - 1,$$

Because the estimator for the missing rate \hat{p} is a simple function of sample averages, we apply the Delta Method to derive its asymptotic properties. Assume $\frac{1}{S} \sum_{s=1}^S E[\psi(G_s)]$ converges to a finite constant as $S \rightarrow \infty$. With (A1) and (A2) holding for each group s , $\frac{1}{S} \sum_{s=1}^S E(\psi_s)$ and $\frac{1}{S} \sum_{s=1}^S E(\tilde{\psi}_s)$ also converge. Denote their limits by μ_ψ and $\mu_{\tilde{\psi}}$ respec-

tively. Let $\chi_s \equiv (\tilde{\psi}_s - E[\tilde{\psi}_s], \psi_s - E[\psi_s])'$ and $\frac{1}{S} \sum_{s=1}^S E(\chi_s \chi_s') \rightarrow \Sigma_\psi$, which we assume is finite, as $S \rightarrow \infty$. By applying the Delta Method, with $\mathcal{R} \equiv \left(\frac{1}{\mu_\psi}, -\frac{\mu_{\tilde{\psi}}}{\mu_\psi^2} \right)$,

$$\sqrt{S}(\hat{p} - p) \xrightarrow{d} \mathcal{N}(0, \mathcal{R}\Sigma_\psi\mathcal{R}').$$

Next, we use \hat{p} to adjust the 2SLS estimator, yielding consistent estimation of λ . To simplify derivation and notation, let $n_s = n$ for $s = 1, \dots, S$. The derivation for the case where n_s varies across the groups is similar, and only differs by requiring versions of the Law of Large Numbers and the Central Limit Theorem for independent and heterogeneous arrays indexed by s . Later, as in our empirical application, we allow for group size variation.

Consider the first scenario in Section 3.4, where the sample reports a single network measure H_s for each group s that is asymmetric. For each group s in the sample and a generic $\tilde{p} \in (0, 1)$, define

$$W_s(\tilde{p}) \equiv \left(\frac{1}{1-\tilde{p}} H_s y_s, X_s \right) \text{ and } Z_s \equiv (H_s' X_s, X_s).$$

Let Y denote an nS -by-1 vector that stacks y_s for $s \leq S$. Similarly, define $\mathbf{W}(\hat{p})$ as an nS -by- $(K+1)$ matrix that stacks $W_s(\hat{p})$ for $s \leq S$, and define \mathbf{Z} as an nS -by- $2K$ matrix that stacks Z_s for $s \leq S$. Our estimator for $\theta \equiv (\lambda, \beta)'$ is:

$$\hat{\theta} \equiv (\mathbf{A}'\mathbf{B}^{-1}\mathbf{A})^{-1} \mathbf{A}'\mathbf{B}^{-1} (\mathbf{Z}'Y), \quad (10)$$

where

$$\mathbf{A} \equiv \mathbf{Z}'\mathbf{W}(\hat{p}) \text{ and } \mathbf{B} \equiv \mathbf{Z}'\mathbf{Z}.$$

The next proposition characterizes the limit distribution of $\hat{\theta}$ as $S \rightarrow \infty$. Define $\Sigma_0 \equiv (A_0' B_0^{-1} A_0)^{-1} A_0' B_0^{-1}$ with $B_0 \equiv E(Z_s' Z_s)$ and $A_0 \equiv E[Z_s' W_s(p)]$, where p is the actual missing rate that generates the sample data. Let $\xi_s \equiv Z_s' v_s - F_0 \chi_s$, where v_s is the n -by-1 vector of composite errors in (6), and F_0 is a $2K$ -by-1 vector defined as:

$$F_0 \equiv E[Z_s' \nabla W_s(p) \theta] = \frac{\lambda}{(1-p)^2} E(Z_s' H_s y_s), \text{ from } \nabla W_s(p) \equiv \frac{dW_s(\tilde{p})}{d\tilde{p}} \Big|_{\tilde{p}=p} = \left(\frac{H_s y_s}{(1-p)^2}, 0 \right).$$

Intuitively, F_0 illustrates how the moment condition in 2SLS depends on the missing rate p , and $-F_0\chi_s$ is the adjustment in the influence function that accounts for the first-stage estimation error in \hat{p} .

Proposition 4. *Suppose (A1), (A2), (A3), and (A4) hold, and (IV-R) is satisfied with $Z \equiv (H'X, X)$. Then*

$$\sqrt{S} \left(\hat{\theta} - \theta \right) \xrightarrow{d} \mathcal{N}(0, \Sigma_0 E(\xi_s \xi_s') \Sigma_0'),$$

under the regularity conditions (REG) in Appendix B.

The conditions (REG), provided in Appendix B, are standard conditions that suffice to apply the Law of Large Numbers, the Central Limit Theorem, and the Delta Method.

Standard errors for $\hat{\theta}$ are calculated by replacing A_0 , B_0 , F_0 , and $E(\xi_s \xi_s')$ with their sample analogs:

$$\hat{\mathbf{A}} = \frac{1}{S} \sum_s \mathbf{z}'_s \mathbf{W}_s(\hat{p}), \hat{\mathbf{B}} = \frac{1}{S} \sum_s \mathbf{z}'_s \mathbf{z}_s, \hat{F} = \frac{1}{S} \frac{\hat{\lambda}}{(1-\hat{p})^2} \sum_s Z'_s H_s y_s, \hat{\xi}_s = Z'_s \hat{v}_s - \hat{F} \hat{\tau}_s,$$

where

$$\hat{v}_s = y_s - \mathbf{W}_s(\hat{p}) \hat{\theta}, \hat{\tau}_s = \left(\frac{1}{\bar{\psi}}, -\frac{\bar{\psi}}{(\bar{\psi})^2} \right) \left(\tilde{\psi}_s - \bar{\psi}, \psi_s - \bar{\psi} \right)',$$

with $\bar{\psi}, \tilde{\psi}$ being averages of $\psi_s, \tilde{\psi}_s$ over $s \leq S$.

We conclude this section by explaining how to apply a similar idea for estimation under the second scenario in Section 3.4. In this case, the sample reports for each group s two symmetric network measures with randomly missing links, $H_s^{(1)}$ and $H_s^{(2)}$, with missing rates $p^{(1)}$ and $p^{(2)}$ respectively. As we show in Section 3.3, this leads to two feasible structural forms, depending on which value of t we use in the expression:

$$y_s = W_s^{(t)} \theta + v_s^{(t)} \text{ for } t = 1, 2, \tag{11}$$

where $\theta \equiv (\lambda, \beta)'$, $W_s^{(t)} \equiv \left(\frac{H_s^{(t)} y_s}{1-p^{(t)}}, X_s \right)$, and $v_s^{(t)} \equiv \varepsilon_s + \lambda \left(G_s - \frac{H_s^{(t)}}{1-p^{(t)}} \right) y_s$. The exogenous instruments for these two systems are respectively $Z_s^{(1)} \equiv (H_s^{(2)} X_s, X_s)$ and $Z_s^{(2)} \equiv$

$(H_s^{(1)}X_s, X_s)$. Let's write:

$$\tilde{Z}_s \equiv \begin{pmatrix} Z_s^{(1)} & 0 \\ 0 & Z_s^{(2)} \end{pmatrix}; \tilde{y}_s \equiv \begin{pmatrix} y_s \\ y_s \end{pmatrix}; \tilde{W}_s \equiv \begin{pmatrix} W_s^{(1)} \\ W_s^{(2)} \end{pmatrix}.$$

Instrument exogeneity implies the following moments:

$$E \left[\tilde{Z}'_s (\tilde{y}_s - \tilde{W}_s \theta) \right] = 0.$$

This moment condition identifies θ , provided $E(\tilde{Z}'_s \tilde{W}_s)$ has full rank. Using arguments similar to Proposition 3 in Section 3.2, we can derive analogous sufficient conditions for this rank condition. We omit the details here for brevity.

We define a system, or stacked, two-stage least squares (S2SLS) estimator as follows. Let $\tilde{\mathbf{Z}}$ denote a $2nS$ -by- $4K$ matrix that is constructed by vertically stacking S matrices $(\tilde{Z}_s)_{s \leq S}$. Likewise construct a $2nS$ -by- $(K+1)$ matrix $\tilde{\mathbf{W}}$ by stacking $(\tilde{W}_s)_{s \leq S}$ (with $p^{(t)}$ estimated by $\hat{p}^{(t)}$) and a $2nS$ -by-1 vector $\tilde{\mathbf{y}}$ by stacking $(\tilde{y}_s)_{s \leq S}$. The S2SLS estimator is

$$\tilde{\theta} \equiv [\tilde{\mathbf{W}}' \tilde{\mathbf{Z}} (\tilde{\mathbf{Z}}' \tilde{\mathbf{Z}})^{-1} \tilde{\mathbf{Z}}' \tilde{\mathbf{W}}]^{-1} \tilde{\mathbf{W}}' \tilde{\mathbf{Z}} (\tilde{\mathbf{Z}}' \tilde{\mathbf{Z}})^{-1} \tilde{\mathbf{Z}}' \tilde{\mathbf{y}}. \quad (12)$$

This provides us with a single estimator that exploits both sets of instruments in the two structural forms in (11). Similar to $\hat{\theta}$ in (10), we can readily construct the standard error for $\tilde{\theta}$ that accounts for estimation error in $\hat{p}^{(1)}, \hat{p}^{(2)}$. We omit details here for brevity.

5 Extensions

We now extend the baseline method in Section 3 to more general settings with contextual effects, heterogeneous missing rates, and group fixed effects. In each case we focus on extending the ideas for constructive identification. Estimation in each of these cases follows from constructive identification arguments and the estimation steps in Section 4.

As before, to fix ideas and simplify notation, let group sizes $n_s = n$ be fixed throughout the remainder of this section. This allows us to suppress the group subscripts s .

5.1 Contextual effects

Suppose the structural form is:

$$y = \lambda Gy + X\beta + GX\gamma + \varepsilon,$$

where γ is a vector of contextual effects, which shows how individual outcomes are directly influenced by the characteristics of others linked to the individual. The reduced form is

$$y = M(X\beta + GX\gamma + \varepsilon),$$

where M is defined as in (4). The noisy structural form based on H is:

$$y = \lambda \frac{Hy}{1-p} + X\beta + \frac{HX}{1-p}\gamma + \eta,$$

where the composite error η is defined as

$$\eta \equiv \varepsilon - \lambda \left(\frac{H}{1-p} - G \right) y - \left(\frac{H}{1-p} - G \right) X\gamma.$$

Under the same conditions and by the same arguments as in the baseline case with no contextual effects (in Section 3.1), rescaling H by $1-p$ makes the new composite error η mean independent from (X, G) . We can similarly construct instruments using H as before. Our next proposition establishes these results. For generality, let $\zeta(X) \in \mathbb{R}^{n \times L}$ be any generic function of X with $L \geq K$.

Proposition 5. *Suppose (A1), (A2), and (A3) hold. Then $E(\eta|X, G) = 0$. If in addition (A4) holds, then $E\{[H'\zeta(X)]'\eta\} = 0$.*

This proposition implies that $H'\zeta(X)$ satisfies instrument exogeneity for generic functions of X . In fact, a stronger result holds under (A1)-(A4): $E(H\eta|G, X) = 0$. The intuition is the same as in Proposition 2. Thus we can apply 2SLS as before to consistently estimate $(\lambda, \beta', \gamma)'$ using $(H'X, X, H'\zeta(X))$ as instruments for $W \equiv \left(\frac{Hy}{1-p}, X, \frac{HX}{1-p} \right)$, provided appropriate rank conditions hold.

5.2 Heterogeneous missing rates

We now extend our methods to allow the missing link rate p to vary with individual characteristics X . To focus on the main idea, we return to the case with no contextual effects as in (6). The generalization to including contextual effects, using the results of the previous sub-section, is straight-forward.

Suppose we replace (A2) with the more general condition:

$$(A2') \ E(H_{ij}|G_{ij} = 1, X) = 1 - p_{ij}(X) \text{ and } E(H_{ij}|G_{ij} = 0, X) = 0 \ \forall i \neq j.$$

Under (A2'), $E(H|G, X) = Q \circ G$, where Q is an n -by- n matrix with its (i, j) -th component $Q_{ij} \equiv 1 - p_{ij}(X)$ and “ \circ ” denotes the Hadamard product. We suppress the dependence of Q on X for simplicity. By the Law of Iterated Expectation,

$$E(H|X) = Q \circ E(G|X).$$

To recover $p_{ij}(\cdot)$, we can apply a method similar to Section 3.4 by focusing on single links and conditioning on X . For example, consider the second scenario in Section 3.4 (the sample reports two noisy measures with missing rates $p_{ij}^{(1)}(X)$ and $p_{ij}^{(2)}(X)$ respectively). Under (A2'), $E\left(H_{ij}^{(t)}|X\right) = \left[1 - p_{ij}^{(t)}(X)\right] E(G_{ij}|X)$ for any $i \neq j$ and $t = 1, 2$. As before, we can construct a third measure $H_{ij}^{(3)} = \max\{H_{ij}^{(1)}, H_{ij}^{(2)}\}$ for each pair $i \neq j$, and then identify the missing rates as

$$p_{ij}^{(1)}(X) = \frac{E(H_{ij}^{(3)} - H_{ij}^{(1)}|X)}{E(H_{ij}^{(2)}|X)} \text{ and } p_{ij}^{(2)}(X) = \frac{E(H_{ij}^{(3)} - H_{ij}^{(2)}|X)}{E(H_{ij}^{(1)}|X)}.$$

In practice, we can avoid the curse of dimensionality in estimation by specifying the missing rates $p_{ij}(X)$ and link formation probability $E(G_{ij}|X)$ as functions of X_i and X_j alone.

With knowledge (or estimates) of $p(X) \equiv \{p_{ij}(X)\}_{i,j \leq n}$, we can use 2SLS to consistently estimate $(\lambda, \beta)'$. Let \tilde{Q} denote a “pointwise inverse” of Q , with the (i, j) -th entry being

$\tilde{Q}_{ij} \equiv 1/(1 - p_{ij})$. With $p(X)$ identified, we can transform the structural form in (6) as

$$y = \lambda \left(\tilde{Q} \circ H \right) y + X\beta + \varepsilon + \underbrace{\lambda[G - (\tilde{Q} \circ H)]}_{v^*} y.$$

Under (A2') and (A3),

$$\begin{aligned} E(v^*|G, X) &= \lambda \{ GE(y|G, X) - E \left[\left(\tilde{Q} \circ H \right) y \mid G, X \right] \} \\ &= \lambda [GMX\beta - \tilde{Q} \circ E(H|G, X)MX\beta] = \lambda(G - \tilde{Q} \circ Q \circ G)MX\beta = 0. \end{aligned} \quad (13)$$

Let $W^* \equiv ((\tilde{Q} \circ H) y, X)$ and $Z^* \equiv (\zeta(X), X)$ where $\zeta(X) \in \mathbb{R}^{n \times L}$ is a nonlinear function of X with $L \geq K$ (e.g., $\zeta(X) \equiv X \circ X$). It follows from (13) that $E(Z^{*'}v^*) = 0$. As long as $E(W^{*'}Z^*)$ and $E[Z^{*'}Z^*]$ both have full rank, then we can use 2SLS to consistently estimate λ and β .⁸

5.3 Group fixed effects

Suppose each group has an unobserved fixed effect α so that the structural form is:

$$y = \lambda Gy + X\beta + \alpha + \varepsilon.$$

Let \bar{G} denote an n -by- n matrix with identical rows, each of which equals the average of all rows in G . Define \bar{H} and \bar{X} analogously. Applying a *within* transformation (as with panel data fixed effects) using the group mean $\bar{y} \equiv \frac{1}{n} \sum_{i=1}^n y_i$, we eliminate α and get

$$y - \bar{y} = \frac{\lambda}{1-p} (H - \bar{H}) y + (X - \bar{X})\beta + v - \bar{v},$$

where

$$v - \bar{v} = \varepsilon - \bar{\varepsilon} + \lambda \left[G - \frac{H}{1-p} - \left(\bar{G} - \frac{\bar{H}}{1-p} \right) \right] y.$$

⁸With heterogeneous missing rates, $H'X$ does not satisfy instrument exogeneity, because $H(\tilde{Q} \circ H) \neq \tilde{Q} \circ H^2$ in general.

Because $\overline{G}, \overline{H}$ are linear functions of G, H respectively, the same argument as Lemma 1 in Section 3 applies to show that $E(v - \bar{v}|X, G) = 0$. We can therefore use 2SLS to estimate $(\lambda, \beta)'$ exactly as before, after applying the within transformation.

6 A Single Large Network

So far we have focused on cases where the sample consists of many small, fixed-sized groups, where no links exist between members of different groups.

In this section we show how our method can be applied to settings with interdependence between *all* individuals in a sample. Specifically, we consider two scenarios in which some forms of weak dependence exist between individuals that are “far enough” from each other. For both scenarios, our proposed 2SLS estimators, when pooling observations over a single large network in the sample, remain consistent and asymptotically normal.

6.1 Nearly block-diagonal (NBD) networks

In this section, we consider a scenario in which the sample can be partitioned into well-defined, *approximate groups*, which we henceforth refer to as “*blocks*”. As before, links within each block are dense (i.e., the probability of forming links between two individuals within the same block does *not* diminish as the sample size increases). But now, in addition, links between individuals from different blocks exist, but they’re sparse, so the probability of forming links across blocks diminishes as the number of blocks increases. Measurement issues arise because of two reasons. First, as before, links that exist within each block are missing randomly from the sample at a fixed rate. Second, those sparse cross-block links that now exist are never reported in the sample.

Formally, we partition the individuals in the sample into S blocks. Each of these blocks is indexed by $s \leq S$, and consists of n_s members, with $n_s \geq 3$ being finite integers. Links between individuals *within the same block* are reported in the sample, but are missing at a rate $p \in (0, 1)$ due to measurement errors. Links between individuals *across different blocks* are not reported in the sample. The sample size is $N \equiv \sum_{s=1}^S n_s$. Let G_N and H_N

denote true and reported N -by- N adjacency matrices that span the observed S blocks. To facilitate investigation of the asymptotic properties of our 2SLS estimators, let \tilde{G}_N be a hypothetical *block-diagonal approximation* of G_N , which perfectly reports all within-block links but drops all cross-block links. That is, for all individual i ,

$$\tilde{G}_{N,ij} = G_{N,ij} \text{ if } j \in s(i); \tilde{G}_{N,ij} = 0 \text{ if } j \notin s(i),$$

where $s(i)$ indicates the block that i belongs to. We maintain the following assumptions on the measurement error in H_N :

$$(N1) \ E(H_{N,ij}|\tilde{G}_N, X) = E(H_{N,ij}|\tilde{G}_{N,ij}, X) \ \forall i \neq j, \text{ and}$$

$$(N2) \ E(H_{N,ij}|\tilde{G}_{N,ij} = 1, X) = 1 - p \text{ and } E(H_{N,ij}|\tilde{G}_{N,ij} = 0, X) = 0 \ \forall i \neq j.$$

Furthermore, we maintain that the block-specific random arrays, $H_{N,s}$, $\tilde{G}_{N,s}$, $X_{N,s}$, $\epsilon_{N,s}$ (with $H_{N,s}$, $\tilde{G}_{N,s}$ being n_s -by- n_s matrices), are drawn independently across the blocks.

We provide conditions under which, in this setting of a single, large network, our 2SLS consistently estimates structural parameters up to augmentation bias, which as before is fixed by deflating by an estimate of $1-p$. Return to the model with no contextual effects, so that the structural form is

$$y_N = \lambda G_N y_N + X_N \beta + \epsilon_N, \tag{14}$$

where y_N , ϵ_N are N -by-1 vectors and X_N is N -by- K matrix of individual characteristics.

In Online Appendix A, we show the feasible version of this structural form using the noisy network measure is

$$y_N = \frac{\lambda}{1-p} H_N y_N + X_N \beta + v_N + u_N, \tag{15}$$

where $u_N \equiv \left(I_N - \lambda \frac{H_N}{1-p} \right) \left(I_N - \lambda \tilde{G}_N \right)^{-1} \lambda \Delta_N y_N$ with $\Delta_N \equiv G_N - \tilde{G}_N$, and

$$v_N \equiv \epsilon_N + \lambda \left(\tilde{G}_N - \frac{H_N}{1-p} \right) \tilde{y}_N \text{ with } \tilde{y}_N \equiv (I_N - \lambda \tilde{G}_N)^{-1} (X_N \beta + \epsilon_N).$$

Note that we decompose composite errors in (15) into u_N and v_N , which are both vectorizations of block-specific vectors $u_{N,s}, v_{N,s}$. As we explain below, v_N is a vectorization of $v_{N,s}$, which are independent across the blocks, whereas the components in u_N are correlated across the blocks because of interdependence between $y_{N,s}$ due to sparse links between the blocks. This difference requires us to apply separate tactics to characterize their contribution to the estimation errors in $\widehat{\theta}_a$.

This decomposition of the composite error is useful for illustrating two key steps for deriving the asymptotic result. To see this, recall the 2SLS estimator that uses $Z_N \equiv (H'_N X_N, X_N)$ as instruments for $R_N \equiv (H_N y_N, X_N)$ is:

$$\widehat{\theta}_a = (A'_N B_N^{-1} A_N)^{-1} A'_N B_N^{-1} Z'_N y_N.$$

where $A_N \equiv Z'_N R_N$ and $B_N \equiv Z'_N Z_N$. By definition,

$$\widehat{\theta}_a - \theta_a = (A'_N B_N^{-1} A_N)^{-1} A'_N B_N^{-1} Z'_N (v_N + u_N),$$

where $\theta_a \equiv \left(\frac{\lambda}{1-p}, \beta'\right)'$ with the subscript a being a reminder that this estimand has augmentation bias. Thus the asymptotic property of the estimator depends on that of $Z'_N v_N$ and $Z'_N u_N$, which we will investigate sequentially.

First, we characterize the order of $Z'_N v_N$, using the fact that $v_{N,s}$ are independent across blocks s . To see why such independence holds, recall that $H_{N,s}, \widetilde{G}_{N,s}, X_{N,s}, \epsilon_{N,s}$ are assumed independent across blocks s . By construct, \widetilde{G}_N, H_N and $(I - \lambda \widetilde{G}_N)^{-1}$ are all block-diagonal. Hence we can write \widetilde{y}_N as a vectorization of *independent*, hypothetical reduced forms. That is, $\widetilde{y}_N = \text{vec}([\widetilde{y}_{N,1}, \widetilde{y}_{N,2}, \dots, \widetilde{y}_{N,s}])$, where $\widetilde{y}_{N,s} = (I_s - \lambda \widetilde{G}_{N,s})^{-1} (X_{N,s} \beta + \epsilon_{N,s})$ are independent across s .⁹ It then follows that $v_{N,s} = \epsilon_{N,s} + \lambda \left(\widetilde{G}_{N,s} - \frac{H_{N,s}}{1-p}\right) \widetilde{y}_{N,s}$, and are independent across s .

We maintain exogeneity and independence conditions which are analogous to (A3) and

⁹We refer to \widetilde{y}_N as a *hypothetical* reduced form, because it is based on the block-diagonal approximation \widetilde{G}_N rather than the actual G_N .

(A4) for the case with small groups in Section 3:

$$(N3) \quad E(\varepsilon_{N,s} | X_{N,s}, G_{N,s}, H_{N,s}) = 0 \text{ for all } s;$$

$$(N4) \quad \text{Conditional on } (G_N, X_N), H_{N,ij} \perp H_{N,kl} \text{ for all } (i, j) \neq (k, l).$$

Under these conditions, $E(v_{N,s} | X_{N,s}, H_{N,s}) = 0$. The independence between $v_{N,s}$ mentioned above then allows us to apply the law of large numbers (Online Appendix Lemma A3) to show that

$$\frac{1}{S} Z'_N v_N = \frac{1}{S} \sum_s Z'_{N,s} v_{N,s} = O_p(S^{-1/2}).$$

Second, for analyzing the large-sample property of $Z'_N u_N$, we exploit the fact that it takes the form of $\mathcal{C}_N \Delta_N y_N$, where both \mathcal{C}_N and y_N are uniformly bounded under mild regularity conditions (Online Appendix Lemma A2). Hence the order of $\frac{1}{S} Z'_N u_N$ is bounded above by the expected number of missing links across the blocks, which are sparse in the following sense:

$$(S\text{-LOB}) \sum_{i=1}^N \sum_{j \notin s(i)} E(|\Delta_{N,ij}|) = O(S^\rho) \text{ for some } \rho < 1.$$

This condition holds if for individuals in each block s , cross-block links only exist with a finite number of nearby blocks, and if the probability for forming such links q_S diminishes as the sample size grows (that is, $q_S = O(S^{-\alpha})$ with $\alpha > 0$).¹⁰ Therefore, with \mathcal{C}_N and y_N bounded, we can establish that $\frac{1}{S} Z'_N u_N = O_p(S^{\rho-1})$ under (S-LOB). (See Online Appendix Lemma A1.) This sparsity condition also ensures A_N, B_N converges in probability to certain deterministic arrays in large samples. (See Online Appendix Lemma A3.) Regularity conditions used for deriving asymptotic properties of $\widehat{\theta}_a - \theta_a$ are collected and presented as Condition (S-REG) in Online Appendix A.

Putting all these pieces together, we show that a feasible 2SLS estimator, which uses a noisy measure H and ignores all links between different blocks, consistently estimates the (augmented) structural parameter $\theta_a \equiv \left(\frac{\lambda}{1-p}, \beta' \right)'$ at a rate that is governed by the order

¹⁰To see this, suppose all blocks have identical size $n < \infty$. Then the expected number of the cross-block links is $c \times S \times n(n-1) \times q_S = O(S^{1-\alpha})$, which satisfies (S-LOB).

of sparse, cross-block links. This result is formalized in the next proposition.

Proposition 6. *Suppose (N1), (N2), (N3) and (N4) hold. If Assumptions (S-LOB) and (S-REG) hold, then*

$$\widehat{\theta}_a - \theta_a = O_p(S^{-1/2} \vee S^{\rho-1}).$$

If in addition $\rho < 1/2$, then

$$\sqrt{S} \left(\widehat{\theta}_a - \theta_a \right) \xrightarrow{d} \mathcal{N}(0, \Omega),$$

where $\Omega \equiv (A'_0 B_0^{-1} A_0)^{-1} A'_0 B_0^{-1} \omega_0 B_0^{-1} A_0 (A'_0 B_0^{-1} A_0)^{-1}$ with A_0, B_0, ω_0 being non-stochastic arrays defined in Online Appendix A.

This proposition implies $\widehat{\theta}_a \xrightarrow{p} \theta_a$ because $\rho < 1$. Furthermore, if $\rho < 1/2$, the asymptotic distribution is determined by the leading term of $\frac{1}{\sqrt{S}} Z'_N u_N$, and hence matches the case of S independent small groups.

To estimate the missing rate p and remove the augmentation bias, one can apply the same method as the first step in Section 4, which remains valid because of independence of $H_{N,s}$ across the blocks $s = 1, 2, \dots, S$.

6.2 Networks with near-epoch dependence (NED)

Here we obtain another consistency result for a different scenario, in which the data consists of a single network that does *not* admit any definition of “approximate groups”, but does include some notion of “distance” between individuals on the network. The main working assumption in this case is that the dependence between two individuals weakens as the distance between them increases, which is reminiscent of the notion of weak dependence in time series models.

Using this primitive condition, we show that observed outcomes satisfy a notion of near-epoch dependence (NED) as used in Jenish and Prucha (2012). Hence a form of the law of large numbers and the central limit theorem can be applied to sample averages over individual outcomes and covariates. We also show that the adjusted-2SLS estimator, when

pooling observations over a single large network in the sample, converges in probability to the structural parameters, where the augmentation bias is removed as before, once missing rates are estimated. Details are in Online Appendix B and C of this paper.

7 Simulation

In this section we use monte carlo simulation to investigate the finite sample performance of our two-step 2SLS estimator in Section 4. Recall that the structural form of the data-generating process is:

$$y_s = \lambda G_s y_s + X_s \beta + \varepsilon_s, \quad s = 1, 2, \dots, S.$$

We fix each group size to be $n_s = 20$ in our simulation. In our design, each member i in each group s has two individual characteristics $X_{s,i} \in \mathbb{R}^2$, which are drawn independently across i and s . The first component $X_{s,i,1}$ is uniformly distributed over a finite support $\{-1, 1, 2\}$ while the second component $X_{s,i,2}$ is standard normal $N(0, 1)$. We consider three designs, corresponding to small, medium, and large peer effects, in which the true parameters are:

$$\lambda \in \{0.20, 0.35, 0.60\} \text{ while } (\beta_1, \beta_2) = (-1.5, 2).$$

The formulation of *undirected* links in the data-generating process is specified as follows. First, each individual sends invitations to two other individuals who are drawn randomly from the same group without replacement. An undirected link exists between two group members if either of them sends an invitation to the other. No links are formed across the groups. This generates each G matrix. Each H matrix is then constructed by dropping existing links randomly at the rate $p = 1/2$.

The size of a sample is defined as the number of independent groups in that sample. For each fixed sample size $S \in \{100, 400, 900\}$, we generate $T = 200$ samples. (Our simulated samples do not contain networks that are singular, which would violate regularity conditions.) By applying our two-step 2SLS estimator from Section 4 in each sample $t = 1, 2, \dots, T$, we record the empirical distribution of these estimates of $(\lambda, \beta_1, \beta_2)$. Table 1 below reports the average bias, sample variance, and mean-squared errors (MSEs) based

on this empirical distribution.

Table 1. Two-step 2SLS Estimator Performance in Simulated Samples

S		$\lambda=0.2$	β_1	β_2	$\lambda=0.35$	β_1	β_2	$\lambda=0.6$	β_1	β_2
100	avg. bias	0.000	0.014	0.002	0.009	0.055	-0.089	-0.303	0.734	-0.679
	variance	0.000	0.009	0.008	0.015	0.175	0.404	0.173	1.672	1.694
	m.s.e.	0.000	0.009	0.008	0.015	0.178	0.412	0.265	2.211	2.155
400	avg. bias	0.000	0.003	0.002	0.006	0.016	-0.037	-0.142	0.361	-0.235
	variance	0.000	0.002	0.002	0.002	0.033	0.083	0.176	0.780	0.606
	m.s.e.	0.000	0.002	0.002	0.002	0.033	0.084	0.196	0.910	0.661
900	avg. bias	0.000	0.001	0.000	0.005	0.004	-0.019	-0.056	0.162	-0.147
	variance	0.000	0.001	0.001	0.001	0.013	0.035	0.072	0.382	0.410
	m.s.e.	0.000	0.001	0.001	0.001	0.013	0.036	0.074	0.408	0.431

Table 1 shows that the mean-squared errors diminish as the sample sizes increase. For each parameter, the rate of decrease in MSE is roughly proportional to the rate of increase in the sample size. This offers evidence for the root- n convergence of our 2SLS estimator. It is also clear that estimator variance accounts for a major portion of the MSEs. For a fixed sample size and design, both the bias and variance of the peer effect λ are smaller than those for the individual effect (β_1, β_2) . We also note that, as the peer effects λ increases, the MSEs increase for all parameters. This might be related to the fact that the variance of the estimator depends on the variation of y_s , which is scaled by $(1 - \lambda G)^{-1}$.

8 Application: Microfinance Participation in India

We apply our method to study how peer effects influence household decisions to participate in a microfinance program in India. The sample was collected by Banerjee et al. (2013) using survey questionnaires from the State of Karnataka, India between 2006-2007. Banerjee et al. (2013) impute a social network structure in the sample by aggregating several network measures that were inferred from the survey responses. They studied how the

dissemination of information about a microfinance program, Bharatha Swamukti Samsathe, or *BSS*, depended on the network position of the households that were the first to be informed about the program. Banerjee et al. (2013) use a binary response model with social interactions to disentangle the effect of information diffusion from the peer effects, a.k.a. *endorsement* effects. In contrast, we use two of the multiple measures in Banerjee et al. (2013) as noisy proxies for an actual network, and apply our method to estimate peer effects in a linear social network model.

8.1 Institutional background and data

The sample was collected by Banerjee et al. (2013) through survey questionnaires from 43 villages in the State of Karnataka, India.¹¹ These villages are largely linguistically homogeneous but heterogeneous in terms of caste. The sample contains information about the socioeconomic status and some demographic characteristics of 9,598 households. On average, there were about 223 households in each village, with a minimum of 114, a maximum of 356, and a standard deviation of 56.2.

We merge the information from a full-scale household census and an individual-level survey in Banerjee et al. (2013). The household census gathered demographic information and data on a variety of amenities, such as roofing material, type of latrine, and quality of access to electric power. The individual survey was administered to a randomly selected sub-sample of villagers, which covered 46% of all households in the census. Individual questionnaires collected demographic information, such as age, caste and sub-caste, education, language, and having a ration card or not, but does not include explicit financial information. We merged the information about the head of household from the individual survey with the household information from the census. This yields a sample of 4,149 households. Table 2(a) reports summary statistics for the dependent variable ($y = 1$ if participates in the microfinance program) as well as a few continuous and binary explanatory variables. Summary statistics for additional categorical variables, such as religion, caste, property ownership, access to electricity, etc, are reported in Table 2(b).

¹¹The data are publicly available at: <http://economics.mit.edu/faculty/eduflo/social>.

Table 2(a): Summary of Dependent and Explanatory Variables

Variable	definition	obs.	mean	s.d.	min	max
<i>y</i>	dummy for participation	4149	0.1894	0.3919	0	1
<i>room</i>	number of rooms	4149	2.4389	1.3686	0	19
<i>bed</i>	number of beds	4149	0.9229	1.3840	0	24
<i>age</i>	age of household head	4149	46.057	11.734	20	95
<i>edu</i>	education of household head	4149	4.8383	4.5255	0	15
<i>lang</i>	whether to speak other language	4149	0.6799	0.4666	0	1
<i>male</i>	whether the hh head is male	4149	0.9161	0.2772	0	1
<i>leader</i>	whether it has a leader	4149	0.1393	0.3463	0	1
<i>shg</i>	whether in any saving group	4149	0.0513	0.2207	0	1
<i>sav</i>	whether to have a bank account	4148	0.3840	0.4864	0	1
<i>election</i>	whether to have an election card	4149	0.9525	0.2127	0	1
<i>ration</i>	whether to have a ration card	4149	0.9012	0.2985	0	1

Table 2(b): Summary of Category Variables

Variable	definition	obs.	per.	Variable	definition	obs.	per.
<i>religion</i>				<i>latrine</i>			
-	Hinduism	3943	95.04	-	Owned	1195	28.80
-	Islam	198	4.77	-	Common	20	0.48
-	Christianity	7	0.19	-	None	2934	70.72
<i>roof</i>				<i>own</i>	property ownership		
-	Thatch	82	1.98	-	Owned	3727	89.83
-	Tile	1388	33.45	-	Owned & shared	32	0.77
-	Stone	1172	28.25	-	Rented	390	9.40
-	Sheet	868	20.92				
-	RCC	475	11.45				
-	Other	164	3.95				
<i>electricity</i>	electricity provision			<i>caste</i>			
-	Private	2662	64.18	-	Scheduled caste	1139	27.54
-	Government	1243	29.97	-	Scheduled tribe	221	5.34
-	No power	243	5.86	-	OBC	2253	54.47
				-	General	523	12.65

The individual-level survey in Banerjee et al. (2013) also collected information about social interactions between households, such as (i) individuals whose homes the respondent visited, and (ii) individuals who visited the respondent’s home. Banerjee et al. (2013) construct graphs with undirected links by symmetrizing the data.¹² That is, the sample provided by Banerjee et al. (2013) contains two symmetric measures for the same latent network, based on the responses to (i) and (ii) respectively. These two measures, reported as “visitGo” and “visitCome” matrices in the sample and denoted as $H^{(1)}$ and $H^{(2)}$ in our notation, lend themselves to application of our method in Section 3.3.¹³

Table 3 reports the empirical distribution of the degrees of $H^{(1)}$ and $H^{(2)}$. Because these measures are symmetric, there is no distinction between the degrees of in-bound or out-bound links. We pool all households across 43 villages into a single, large network. There are no links between households from different villages in the sample, so the network structure is block-diagonal.

Table 3: Degree Distribution in Two Network Measures

Degree	0	1	2	3	4	5	6	7	8	9	10
$H^{(1)}$	2	21	110	227	357	505	526	546	506	379	269
$H^{(2)}$	4	24	112	245	384	522	534	577	491	386	255
Degree	11	12	13	14	15	16	17	18	19	20	≥ 21
$H^{(1)}$	224	145	90	74	54	33	27	15	9	6	24
$H^{(2)}$	179	137	102	59	46	28	22	13	9	3	17

Table 3 indicates large variation in the number of connections the households have. If there were no missing links in these reported measures, we would expect the two matrices $H^{(1)}$ and $H^{(2)}$ to be identical, and therefore have exactly the same degree distribution. The fact that they differ substantially is indicative of many missing links, possibly due to the respondents’ recall errors, or to differences in how they interpret the visiting question.

¹²Two households i and j are considered connected by an undirected link if an individual from either household mentioned the name of someone from the other household in response to the question in (1). Likewise, a second symmetric network measure is constructed based on responses to (2).

¹³Banerjee et al. (2013) aggregate responses from 12 questions, including (1) and (2), to construct a single symmetric network, that they assume equals, without any errors, an actual relevant adjacency matrix G . In contrast, we take a different approach by interpreting responses to questions (1) and (2) as two different noisy measures of a true underlying latent network.

8.2 Empirical strategy for estimating peer effects

We use the following specification for the feasible structural form:

$$y = \lambda \left(\frac{H^{(t)}}{1 - p^{(t)}} \right) y + X\beta + \text{villageFE} + v^{(t)} \text{ for } t = 1, 2, \quad (16)$$

where y is a binary variable indicating whether the household participated in the microfinance program (BSS), X is a matrix of household characteristics, and villageFE are village fixed effects. Definition and summary statistics of regressors in X are listed in Table 2. Note that (16) provides *two* different feasible structural forms (of the same underlying true structural model), corresponding to $t = 1, 2$ respectively.

To implement an adjusted-2SLS estimator, we first estimate the missing rates $p^{(t)}$ for $t = 1, 2$, and use them to rescale the endogenous regressors as in Section 4. Following Section 3.4, we construct $H^{(3)} = \max\{H^{(1)}, H^{(2)}\}$ and estimate the missing rates as

$$\hat{p}^{(1)} = \frac{\psi(H^{(3)}) - \psi(H^{(1)})}{\psi(H^{(2)})} = 0.1681, \text{ and } \hat{p}^{(2)} = \frac{\psi(H^{(3)}) - \psi(H^{(2)})}{\psi(H^{(1)})} = 0.1909,$$

where $\psi(H)$ is the mean of off-diagonal entries in H . We replace $p^{(1)}$ and $p^{(2)}$ in equation (16) with $\hat{p}^{(1)}$ and $\hat{p}^{(2)}$ respectively, and then apply the 2SLS estimators in Section 4.

The results are reported in Table 4. The columns of Table 4 are all 2SLS estimates, defined as follows:

Column (a) ignores missing links in $H^{(1)}$, and so treats $H^{(1)}$ as if it were the true adjacency matrix G , by putting (unscaled) $\lambda H^{(1)}y$ on the right-hand side, and using $H^{(1)}X$ as the instruments for $H^{(1)}y$ in 2SLS.

Column (b) estimates the structural form for $t = 1$ in (16), using $H^{(2)}X$ as instruments for $\left(\frac{H^{(1)}}{1 - \hat{p}^{(1)}} \right) y$ in adjusted 2SLS.

Column (c) is identical to Column (a), except for using $H^{(2)}$ instead of $H^{(1)}$ everywhere, and so treats $H^{(2)}$ as if it were the true matrix G for 2SLS estimation

Column (d) is identical to column (b), except for switching the roles of the matrices $H^{(1)}$ and $H^{(2)}$. So the feasible structural model in (16) is written in terms of $t = 2$, and

$H^{(1)}X$ is used as instruments for $\left(\frac{H^{(2)}}{1-\hat{\rho}^{(2)}}\right)y$.

Column (e) applies the S2SLS estimator defined in (12) in Section 4. This estimator combines (stacks) the 2SLS moments used in Columns (b) and (d) above, and so combines the moments generated by both of the feasible structural models and their associated IVs into a single estimator.

In summary, the estimators in (a) and (c) are what a researcher would do if he or she ignored the missing links problem and treated either $H^{(1)}$ or $H^{(2)}$, respectively, as if it were the true adjacency matrix G , applying the standard 2SLS estimator that is proposed in the literature. In contrast, the corresponding adjusted-2SLS estimators in (b), (d) and (e) are estimators that we propose to remove the augmentation bias in 2SLS resulting from missing links.¹⁴ Column (e) in particular combines the information used to construct the estimators in both columns (b) and (d), and so is our preferred estimator.

8.3 Empirical results

Table 4 reports that our adjusted 2SLS estimates for the peer effect $\hat{\lambda}$ are 0.0456 when using $H^{(1)}y$ in the structural form (column (b)), 0.0484 using $H^{(2)}y$ (column (d)), and 0.0461 using both measures and S2SLS (column (e)). These estimates are all significant at the 1% level, and the differences between them are small relative to the standard errors. These estimates imply the likelihood of a household to participate in the microfinance program is increased by about 4.6% when the household is linked to one more participating household on the network (note for this calculation that our model does not row-normalize the network measures). With the average participation rate being 18.9% in the sample, these estimates suggest that peer effects, called “endorsement effects” in Banerjee et al. (2013), are economically substantial.

¹⁴We need two noisy network measures in this particular context because the available reported measures are symmetric. As we show in Section 3.2, our method can also be used if the sample reports a single yet *asymmetric* noisy measure of the network.

Table 4: Two-stage Least Square Estimates

	(a)	(b)	(c)	(d)	(e)
r.h.s. endogeneity	$H^{(1)}y$	$\frac{H^{(1)}}{1-\hat{p}_1}y$	$H^{(2)}y$	$\frac{H^{(2)}}{1-\hat{p}_2}y$	$\frac{H^{(t)}}{1-\hat{p}}y$
IV used	$H^{(1)}X$	$H^{(2)}X$	$H^{(2)}X$	$H^{(1)}X$	Combined
$\hat{\lambda}$	0.0498*** (0.0076)	0.0456*** (0.0096)	0.0529*** (0.0092)	0.0484*** (0.0087)	0.0461*** (0.0075)
<i>leader</i>	0.0378** (0.0185)	0.0364** (0.0186)	0.0418** (0.0182)	0.0405** (0.0182)	0.0387** (0.0183)
<i>age</i>	-0.0016*** (0.0005)	-0.0017*** (0.0005)	-0.0016*** (0.0005)	-0.0017*** (0.0005)	-0.0017*** (0.0005)
<i>ration</i>	0.0441** (0.0201)	0.0435** (0.0201)	0.0423** (0.0195)	0.0413** (0.0194)	0.0426** (0.0197)
<i>electricity – gov</i>	0.0343** (0.0157)	0.0333** (0.0157)	0.0352** (0.0156)	0.0341** (0.0155)	0.0339** (0.0156)
<i>electricity – no</i>	0.0223 (0.0297)	0.0229 (0.0297)	0.0237 (0.0300)	0.0247 (0.0298)	0.0236 (0.0298)
<i>caste – tribe</i>	-0.0285 (0.0312)	-0.0272 (0.0309)	-0.0275 (0.0305)	-0.0257 (0.0300)	-0.0268 (0.0305)
<i>caste – obc</i>	-0.0520** (0.0217)	-0.0490** (0.0212)	-0.0486** (0.0215)	-0.0441*** (0.0206)	-0.0473*** (0.0210)
<i>caste – gen</i>	-0.0734*** (0.0239)	-0.0698*** (0.0242)	-0.0688*** (0.0241)	-0.0628** (0.0234)	-0.0673*** (0.0239)
<i>religion – Islam</i>	0.0980*** (0.0323)	0.0955*** (0.0323)	0.0893*** (0.0343)	0.0849*** (0.0344)	0.0910*** (0.0332)
<i>religion – Chri</i>	0.1434 (0.130)	0.1420 (0.1287)	0.1466 (0.1314)	0.1452 (0.1300)	0.1438 (0.1293)
<i>Controls</i>	✓	✓	✓	✓	✓
<i>VillageFE</i>	✓	✓	✓	✓	✓
R^2	0.1332	0.1345	0.1350	0.1365	0.1353
Obs	4134	4134	4134	4134	4134

Note: s.e. in parentheses. ***, **, and * indicate 1%, 5% and 10% significant.

Controls include *male, roof, room, bed, latrine, edu, lang, shg, sav, election, own*.

The signs of estimated marginal effects by individual or household characteristics are plausible. Column (e) suggests the head of household being a “leader” (e.g. a teacher, a leader of a self-help group, or a shopkeeper) increases the participation rate by around 3.9%. These households with “leaders” were the first ones to be informed about the program, and were asked to forward information about the microfinance program to other potentially interested villagers. These leaders had received first-hand, detailed information about the program from its administrator, which could be conducive to higher participation rates. Households with younger heads are more likely to participate, but the magnitude of this age effect is less substantial. Being 10 years younger increases the participation rate by 1.7%. Having a ration card increases the participation rate by around 4.3%. Compared to households using private electricity, households using government-supplied electricity have a 3.4% higher participation rate. These two factors indicate that, holding other factors equal, households in poorer economic conditions are more inclined to participate in the microfinance program.

Table 4 also shows that, if we had ignored the issue of missing links in network measures, and had done 2SLS using $H^{(t)}X$ as instruments for the (unscaled) endogenous peer outcomes $H^{(t)}y$, then the estimator would have been considerably biased upward. In (a), where we use $H^{(1)}X$ as instruments for $H^{(1)}y$, the estimate for λ is 0.0498. In comparison, in (b) where we correct for missing link bias by using $H^{(2)}X$ as instruments for $\frac{H^{(1)}y}{1-\hat{p}^{(1)}}$, then the estimated λ is 0.0456. The upward bias resulted from ignoring the missing links is about 9.2% (as $0.0498/0.0456=1.092$). Likewise, in (c) where we erroneously use $H^{(2)}X$ as instruments for $H^{(2)}y$, we get a proportionally almost the same upward bias in the peer effect estimate compared with the correct estimate in (d) (as $0.0529/0.0484=1.093$).

The over 9% upward bias in (a) and (c) is a manifestation of two factors at work. First, with missing links the instruments $H^{(t)}X$ are invalid because of the correlation between $H^{(t)}X$ and the composite errors $v^{(t)}$. Second, even if these instruments were valid, the augmentation bias, as defined in Section 3.1, would be present without rescaling the endogenous peer outcomes $H^{(t)}y$ by $1 - p^{(t)}$.

The magnitude of this upward bias is determined by the magnitude of $p^{(t)}$ and by the

correlation between the composite error and the invalid instruments. The microfinance survey data in Banerjee et al. (2013) is considered to have high quality social network information. In other empirical environments, we may expect even larger bias when missing links are not accounted for in estimation. The method we propose in this paper provides an easy remedy for this issue.

We conclude this section with some model validation results in Table 5, which shows how the predicted values of $E(y|X)$ fit with the sample data. The Probit and Logit models use the same set of regressors as in Table 4. We report the summary statistics of the fitted values $\widehat{E}(y|X)$ under different models. Columns (a) through (d) of Table 5 are the fitted values of the feasible structural models used in each of the corresponding columns in Table 4. Column (e) in Table 4 used two different feasible structural models to obtain S2SLS estimates. To make use of both for fitted values, in column (e) of table 5 we use the S2SLS estimates of (λ, β) and construct fitted values based on $\hat{\lambda} \frac{H^{(3)}}{1-p_1 p_2} y + X \hat{\beta} + \hat{F}E$, where $H^{(3)}$ is as defined in Section 3.4.

In all but one of the models in Table 5, the sample mean of the predicted participation probability $\widehat{E}(y|X)$ is 0.1894, which is equal to the sample mean of y in the 4,134 observations used in the regression. The standard deviation of the predicted participation probability varies across different models. Predictions of linear probability models (LPM), reported under the column of “OLS” and (a)-(e), are mostly within the unit interval $[0, 1]$. LPM predictions are strictly less than 1 for all observations in the sample; Only 2.95% to 5.49% of the households in the sample end up with negative LPM predictions. That is, about 95% all LPM predictions in the sample are indeed within the unit interval.

Based on $\widehat{E}(y|X)$, we use the indicator $I(\widehat{E}(y|X) > 0)$ to predict whether an individual participates in the microfinance program, and calculate prediction rates. Predictions in our linear social network models in columns (a)-(e) generally outperform the OLS, Probit and Logit models in terms of the percentage of correct predictions.

Table 5: Model Validation: Predicted Microfinance Participation

$\widehat{E}(y X)$	Probit	Logit	OLS	(a)	(b)	(c)	(d)	(e)
<i>mean</i>	0.1894	0.1894	0.1894	0.1894	0.1894	0.1894	0.1894	0.1881
<i>s.t.d</i>	0.1176	0.1181	0.1151	0.1339	0.1376	0.1356	0.1409	0.1337
<i>min</i>	0.0103	0.0166	-0.095	-0.104	-0.108	-0.127	-0.131	-0.110
<i>max</i>	0.7490	0.7673	0.6895	0.7807	0.8016	0.7279	0.7576	0.8036
<i>< 0</i>	0%	0%	2.95%	4.67%	4.98%	4.79%	5.49%	4.84%
$I\{\widehat{E}(y X) > 0.5\}$								
<i>underpredict</i> (1 to 0)	17.76%	17.66%	18.34%	17.34%	17.17%	17.34%	17.13%	17.30%
<i>overpredict</i> (0 to 1)	0.92%	1.11%	0.27%	0.87%	1.02%	0.85%	0.97%	0.80%
<i>correct</i>	81.33%	81.23%	81.40%	81.79%	81.81%	81.81%	81.91%	81.91%

9 Conclusion

This paper proposes adjusted-2SLS estimators that consistently estimate structural parameters, which include peer, individual, and contextual effects, in social network models when actual existing links are missing randomly from the sample. By rescaling the endogenous peer outcomes and applying new instruments constructed from noisy network measures, our estimators resolve the additional endogeneity issues caused by missing links. As an intermediate step of the method, we provide methods to estimate the rates at which links are missing from noisy measures of network links. We also show that ignoring missing links generally leads to augmentation bias, i.e., peer effect estimates are generally biased upward.

We apply our method to analyze the peer (endorsement) effects in households’ decisions to participate in a microfinance program in Indian villages, using the data collected by Banerjee et al. (2013). Consistent with our theoretical results, our empirical estimates show that ignoring the issue of missing links in the 2SLS estimation of the social network model leads to a substantial upward bias (over 9%) in the estimates of peer effects.

References

- Advani, A. and B. Malde (2018). Credibly identifying social effects: Accounting for network formation and measurement error. *Journal of Economic Surveys* 32(4), 1016–1044.
- Banerjee, A., A. G. Chandrasekhar, E. Duflo, and M. O. Jackson (2013). The diffusion of microfinance. *Science* 341(6144), 1236498.
- Blume, L. E., W. A. Brock, S. N. Durlauf, and Y. M. Ioannides (2011). Identification of social interactions. In *Handbook of social economics*, Volume 1, pp. 853–964. Elsevier.
- Boucher, V. and A. Houndetoungan (2020). *Estimating peer effects using partial network data*. Centre de recherche sur les risques les enjeux économiques et les politiques.
- Bramoullé, Y., H. Djebbari, and B. Fortin (2009). Identification of peer effects through social networks. *Journal of econometrics* 150(1), 41–55.
- Butts, C. T. (2003). Network inference, error, and informant (in) accuracy: a bayesian approach. *social networks* 25(2), 103–140.
- Chandrasekhar, A. and R. Lewis (2011). Econometrics of sampled networks. *Unpublished manuscript, MIT.[422]*.
- Goldsmith-Pinkham, P. and G. W. Imbens (2013). Social networks and the identification of peer effects. *Journal of Business & Economic Statistics* 31(3), 253–264.
- Graham, B. S. (2020). Network data. In *Handbook of Econometrics*, Volume 7, pp. 111–218. Elsevier.
- Griffith, A. (2021). Name your friends, but only five? the importance of censoring in peer effects estimates using social network data. *Journal of Labor Economics*.
- Hardy, M., R. M. Heath, W. Lee, and T. H. McCormick (2019). Estimating spillovers using imprecisely measured networks. *arXiv preprint arXiv:1904.00136*.
- Hu, Y. (2008). Identification and estimation of nonlinear models with misclassification error using instrumental variables: A general solution. *Journal of Econometrics* 144(1), 27–61.

- Jenish, N. and I. R. Prucha (2012). On spatial processes and asymptotic inference under near-epoch dependence. *Journal of econometrics* 170(1), 178–190.
- Kelejian, H. H. and I. R. Prucha (1998). A generalized spatial two-stage least squares procedure for estimating a spatial autoregressive model with autoregressive disturbances. *The Journal of Real Estate Finance and Economics* 17(1), 99–121.
- Lee, L.-F. (2007). Identification and estimation of econometric models with group interactions, contextual factors and fixed effects. *Journal of Econometrics* 140(2), 333–374.
- Lewbel, A. (2007). Estimation of average treatment effects with misclassification. *Econometrica* 75(2), 537–551.
- Lin, X. (2010). Identifying peer effects in student academic achievement by spatial autoregressive models with group unobservables. *Journal of Labor Economics* 28(4), 825–860.
- Liu, X. (2013). Estimation of a local-aggregate network model with sampled networks. *Economics Letters* 118(1), 243–246.
- Mahajan, A. (2006). Identification and estimation of regression models with misclassification. *Econometrica* 74(3), 631–665.
- Shalizi, C. R. and A. Rinaldo (2013). Consistency under sampling of exponential random graph models. *Annals of statistics* 41(2), 508.

Appendix

A. Identification proofs

Proof of Lemma 1. Under (A3), we have $E(Gy|X, G) = E[GM(X\beta + \varepsilon)|X, G] = GMX\beta$, and $E(Hy|X, G) = E[HME(X\beta + \varepsilon|X, G, H)|X, G] = E(H|G, X)MX\beta$.

Under (A1) and (A2), $E(H|G, X) = (1 - p)G$. It follows from the definition of v in (6) that $E(v|X, G) = 0$. □

Proof of Proposition 2. Under (A1), (A2), and (A4), the conditional mean of the (i, j) -th entry in H^2 is

$$\begin{aligned}
E[(H^2)_{ij}|G, X] &= E\left(\sum_{k \neq i, j} H_{ik} H_{kj} \middle| G, X\right) = \sum_{k \neq i, j} E(H_{ik} H_{kj} | G, X) \\
&= \sum_{k \neq i, j} E(H_{ik} | G_{ik}, X) E(H_{kj} | G_{kj}, X) = \sum_{k \neq i, j} (1-p)G_{ik}(1-p)G_{kj} \\
&= (1-p)^2 (G^2)_{ij}.
\end{aligned} \tag{17}$$

Besides, under (A1) and (A2),

$$E[HG|G, X] = E(H|G, X)G = (1-p)G^2. \tag{18}$$

It then follows that

$$\begin{aligned}
E[(H'X)'v|G, X] &= E(X'H\varepsilon|G, X) + \lambda E\left[X'H\left(G - \frac{H}{1-p}\right)y \middle| G, X\right] \\
&= \lambda E\left[X'H\left(G - \frac{H}{1-p}\right)MX\beta \middle| G, X\right] \\
&= \lambda X' \left(E(HG|G, X) - \frac{E(H^2|G, X)}{1-p}\right)MX\beta = 0,
\end{aligned}$$

where the first two equalities are due to (A3), and the last holds because of (17) and (18) under (A1), (A2), and (A4). \square

As we noted in Section 3.3, one can construct instruments from multiple symmetric measures for G , denoted by $H^{(1)}$ and $H^{(2)}$. Suppose $H^{(1)}$ and $H^{(2)}$ both satisfy (A1), (A2), (A3), and are independent in the sense of (A4'). Then one can construct feasible structural forms as in (9), and use $H^{(2)}X$ as instruments for $v^{(1)}$, and vice versa. To see why, note:

$$\begin{aligned}
E[(H^{(2)}H^{(1)})_{ij}|G, X] &= E\left(\sum_{k \neq i, j} H_{ik}^{(2)} H_{kj}^{(1)} \middle| G, X\right) \\
&= \sum_{k \neq i, j} E\left(H_{ik}^{(2)} H_{kj}^{(1)} \middle| G, X\right) = \sum_{k \neq i, j} E\left(H_{ik}^{(2)} \middle| G_{ik}, X\right) E\left(H_{kj}^{(1)} \middle| G_{kj}, X\right) \\
&= \sum_{k \neq i, j} (1-p^{(2)})G_{ik}(1-p^{(1)})G_{kj} = (1-p^{(2)})(1-p^{(1)}) (G^2)_{ij}.
\end{aligned} \tag{19}$$

Besides, under (A1) and (A2),

$$E[H^{(2)}G|G, X] = E(H^{(2)}|G, X)G = (1 - p^{(2)})G^2. \quad (20)$$

It then follows that

$$\begin{aligned} E[(H^{(2)}X)'v^{(1)}|G, X] &= E(X'H^{(2)}\varepsilon|G, X) + \lambda E\left[X'H^{(2)}\left(G - \frac{H^{(1)}}{1 - p^{(1)}}\right)y\middle|G, X\right] \\ &= \lambda E\left[X'H^{(2)}\left(G - \frac{H^{(1)}}{1 - p^{(1)}}\right)MX\beta\middle|G, X\right] \\ &= \lambda X'\left(E(H^{(2)}G|G, X) - \frac{E(H^{(2)}H^{(1)}|G, X)}{1 - p^{(1)}}\right)MX\beta = 0. \end{aligned}$$

where the first two equalities are due to (A3), and the last holds because of (19) and (20) under (A1), (A2), and (A4').

Proof of Proposition 3. Define the following K -by- K moments involving (G, X) :

$$\begin{aligned} B_1 &\equiv E(X'G^2MX), \quad B_2 \equiv E(X'GMX), \quad B_3 \equiv E(X'G^2X), \\ B_4 &\equiv E(X'GX), \quad B_5 \equiv E(X'X). \end{aligned}$$

Under (A1), (A2), (A3), and (A4),

$$\begin{aligned} E(Z'R) &= \begin{pmatrix} E(X'H^2y) & E(X'HX) \\ E(X'Hy) & E(X'X) \end{pmatrix} = \begin{pmatrix} E[X'H^2M(X\beta + \varepsilon)] & E(X'HX) \\ E[X'HM(X\beta + \varepsilon)] & E(X'X) \end{pmatrix} \\ &= \begin{pmatrix} (1 - p)^2E(X'G^2MX\beta) & (1 - p)E(X'GX) \\ (1 - p)E(X'GMX\beta) & E(X'X) \end{pmatrix} \equiv \begin{pmatrix} (1 - p)^2B_1\beta & (1 - p)B_4 \\ (1 - p)B_2\beta & B_5 \end{pmatrix}. \end{aligned}$$

Suppose the $2K$ -by- $(1 + K)$ matrix $E(Z'R)$ does not have full rank. By definition the $2K$ -by- $2K$ square matrix

$$\begin{pmatrix} (1 - p)^2B_1 & (1 - p)B_4 \\ (1 - p)B_2 & B_5 \end{pmatrix}$$

must be singular. This implies $[B_1, B_4; B_2, B_5]$ must also be singular because

$$\begin{aligned} \det \begin{pmatrix} (1-p)^2 B_1 & (1-p)B_4 \\ (1-p)B_2 & B_5 \end{pmatrix} &= \det(B_5) \det [(1-p)^2 B_1 - (1-p)^2 B_4 (B_5)^{-1} B_2] \\ &= (1-p)^{2K} \det(B_5) \det(B_1 - B_4 B_5^{-1} B_2) = (1-p)^{2K} \det \begin{pmatrix} B_1 & B_4 \\ B_2 & B_5 \end{pmatrix}. \end{aligned}$$

Therefore, non-singularity of $[B_1, B_4; B_2, B_5]$ implies that $E(Z'R)$ has full rank.

As $M - \lambda GM = I$, we have $GM = \frac{1}{\lambda}(M - I)$ and $G^2 M = \frac{1}{\lambda}(GM - G) = \frac{1}{\lambda^2}(M - I - \lambda G)$.

We can write

$$\begin{pmatrix} B_1 & B_4 \\ B_2 & B_5 \end{pmatrix} = \begin{pmatrix} \frac{1}{\lambda} E(X'(GM - G)X) & E(X'GX) \\ E(X'GMX) & E(X'X) \end{pmatrix},$$

with the 1st row plus the 2nd row $\ast(-\frac{1}{\lambda})$,

$$\begin{pmatrix} -\frac{1}{\lambda} E(X'GX) & E(X'GX) - \frac{1}{\lambda} E(X'X) \\ E(X'GMX) & E(X'X) \end{pmatrix},$$

with the 1st column plus the 2nd column $\ast(\frac{1}{\lambda})$,

$$\begin{pmatrix} -\frac{1}{\lambda^2} E(X'X) & E(X'GX) - \frac{1}{\lambda} E(X'X) \\ E(X'(GM + \frac{1}{\lambda}I)X) & E(X'X) \end{pmatrix} = \begin{pmatrix} -\frac{1}{\lambda^2} E(X'X) & -\frac{1}{\lambda} E(X'M^{-1}X) \\ \frac{1}{\lambda} E(X'MX) & E(X'X) \end{pmatrix}.$$

Hence, $\begin{pmatrix} B_1 & B_4 \\ B_2 & B_5 \end{pmatrix}$ is non-singular iff $\begin{pmatrix} E(X'X) & E(X'M^{-1}X) \\ E(X'MX) & E(X'X) \end{pmatrix}$ is non-singular.

By the same token, (A1), (A2), and (A4) imply that

$$\begin{aligned} E(Z'Z) &= \begin{pmatrix} E(X'H^2X) & E(X'HX) \\ E(X'HX) & E(X'X) \end{pmatrix} = \begin{pmatrix} (1-p)^2 E(X'G^2X) & (1-p)E(X'GX) \\ (1-p)E(X'GX) & E(X'X) \end{pmatrix} \\ &= \begin{pmatrix} (1-p)^2 B_3 & (1-p)B_4 \\ (1-p)B_4 & B_5 \end{pmatrix}. \end{aligned}$$

Similarly, the determinant of $E(Z'Z)$ is proportional to that of $[B_3, B_4; B_4, B_5]$. Therefore the non-singularity of $[B_3, B_4; B_4, B_5]$ implies $E(Z'Z)$ has full rank. \square

Proof of Proposition 5. Under (A3), we have

$$\begin{aligned} E(Gy|X, G) &= E[GM(X\beta + GX\gamma + \varepsilon)|X, G] = GM(X\beta + GX\gamma), \\ E(Hy|X, G) &= E[HME(X\beta + GX\gamma + \varepsilon|X, G, H)|X, G] = E(H|G, X)M(X\beta + GX\gamma). \end{aligned}$$

Under (A1) and (A2), $E(H|G, X) = (1 - p)G$. It then follows that $E(\eta|X, G) = 0$. Note

$$\begin{aligned} E[\zeta(X)'HHy|G, X] &= \zeta(X)'E(H^2|G, X)M(X\beta + GX\gamma); \\ E[\zeta(X)'HHX|G, X] &= \zeta(X)'E(H^2|G, X)X; \\ E[\zeta(X)'HGy|G, X] &= \zeta(X)'E(H|G, X)GM(X\beta + GX\gamma); \\ E[\zeta(X)'HGX|G, X] &= \zeta(X)'E(H|G, X)GX. \end{aligned}$$

As shown in the proof of Proposition 2, under (A4), $E(H^2|G, X) = (1 - p)^2G$. Because $E(H|G, X) = (1 - p)G$ under (A1) and (A2), this implies $E[\zeta(X)'H\eta] = 0$. \square

B. Asymptotic property of two-step Estimator

In this section we sketch a proof of asymptotic distribution for \hat{p} , $\hat{\lambda}$, and $\hat{\beta}$. We maintain the following regularity conditions:

(REG) $E(\psi_s) \neq 0$; $0 < p < 1$; $E(|Z'_s W_s(p)|) < \infty$, $E(|Z'_s Z_s|) < \infty$, $E(\|\xi_s\|^2) < \infty$ where ξ_s is defined below.

These conditions are needed for applying the law of large numbers, the central limit theorem, and the delta method below.

First off, by the central limit theorem,

$$\frac{1}{\sqrt{S}} \begin{pmatrix} \sum_s [\tilde{\psi}_s - E(\tilde{\psi}_s)] \\ \sum_s [\psi_s - E(\psi_s)] \end{pmatrix} \xrightarrow{d} \mathcal{N}(0, \Omega),$$

where Ω is the covariance matrix of $(\tilde{\psi}_s, \psi_s)'$. The delta method implies $\sqrt{S}(\hat{p} - p) \xrightarrow{d} \mathcal{N}(0, D\Omega D')$, where

$$D = \left(\frac{1}{E(\psi_s)}, -\frac{E(\tilde{\psi}_s)}{E(\psi_s)^2} \right).$$

The asymptotic linear presentation of \hat{p} is

$$\sqrt{S}(\hat{p} - p) = \frac{1}{\sqrt{S}} \sum_s \tau_s + o_p(1),$$

where $\tau_s \equiv D \times \left(\tilde{\psi}_s - E(\tilde{\psi}_s), \psi_s - E(\psi_s) \right)'$ with $E[\tau_s] = 0$.

Hence, $\sqrt{S}(\hat{p} - p) \xrightarrow{d} \mathcal{N}(0, E(\tau_s \tau_s'))$. Next, note that by construction,

$$\begin{aligned} \sqrt{S}(\hat{\theta} - \theta) &= \sqrt{S}(\mathbf{A}'\mathbf{B}^{-1}\mathbf{A})^{-1} \mathbf{A}'\mathbf{B}^{-1}\mathbf{Z}' [Y - \mathbf{W}(\hat{p})\theta] \\ &= (A'_0 B_0^{-1} A_0)^{-1} A'_0 B_0^{-1} \frac{1}{\sqrt{S}} \mathbf{Z}' [Y - \mathbf{W}(\hat{p})\theta] + o_p(1), \end{aligned} \quad (21)$$

where the second “=” holds since $\mathbf{A}/S \xrightarrow{p} A_0$, $\mathbf{B}/S \xrightarrow{p} B_0$ and $\frac{1}{\sqrt{S}} \mathbf{Z}' [Y - \mathbf{W}(\hat{p})\theta] = O_p(1)$.

Recall the definition from the text:

$$F_0 \equiv E[Z'_s \nabla W_s(p)\theta] = \frac{\lambda}{(1-p)^2} Z'_s H_s y_s, \text{ from } \nabla W_s(p) \equiv \frac{dW_s(\hat{p})}{d\hat{p}} \Big|_{\hat{p}=p} = \left(\frac{H_s y_s}{(1-p)^2}, 0 \right).$$

Let $\nabla \mathbf{W}(p)$ be nS -by- $(K+1)$ matrix that stacks $\nabla W_s(p)$ over $s \leq S$. Then

$$\begin{aligned} \frac{1}{\sqrt{S}} \mathbf{Z}' (Y - \mathbf{W}(\hat{p})\theta) &= \frac{1}{\sqrt{S}} \mathbf{Z}' (Y - \mathbf{W}(p)\theta) - \left(\frac{1}{S} \mathbf{Z}' \nabla \mathbf{W}(p)\theta \right) \sqrt{S}(\hat{p} - p) + o_p(1) \\ &= \frac{1}{\sqrt{S}} \sum_s Z'_s (y_s - W_s(p)\theta) - F_0 \left(\frac{1}{\sqrt{S}} \sum_s \tau_s \right) + o_p(1) \\ &= \frac{1}{\sqrt{S}} \sum_s \underbrace{(Z'_s v_s - F_0 \tau_s)}_{\xi_s} + o_p(1), \end{aligned} \quad (22)$$

The first equality follows from a Taylor approximation around the true missing rate p ; the second from $\left(\frac{1}{S} \mathbf{Z}' \nabla \mathbf{W}(p)\theta \right) \xrightarrow{p} E[Z'_s \nabla W_s(p)\theta]$ and from the asymptotic linear representation of the estimator \hat{p} ; the third from $y_s = W_s(p)\theta + v_s$. This proves the claim of limiting distribution of $\sqrt{S}(\hat{\theta} - \theta)$ in the text.

Supplemental Appendix:

Estimating Social Network Models with Missing Links

Arthur Lewbel, Xi Qu, and Xun Tang

December 23, 2022

A. Proofs for NBD networks

We begin by deriving the noisy, feasible structural form in (15). First off, note that the reduced form of y_N is:

$$\begin{aligned}
 y_N &= (I_N - \lambda G_N)^{-1}(X_N\beta + \varepsilon_N) \\
 &= (I_N - \lambda \tilde{G}_N)^{-1}(X_N\beta + \varepsilon_N) - \left[(I_N - \lambda \tilde{G}_N)^{-1} - (I_N - \lambda G_N)^{-1} \right] (X_N\beta + \varepsilon_N) \quad (1) \\
 &= \underbrace{(I_N - \lambda \tilde{G}_N)^{-1}(X_N\beta + \varepsilon_N)}_{\equiv \tilde{y}_N} + (I_N - \lambda \tilde{G}_N)^{-1} \underbrace{\lambda(G_N - \tilde{G}_N)}_{\equiv \Delta_N} \underbrace{(I_N - \lambda G_N)^{-1}(X_N\beta + \varepsilon_N)}_{=y_N}.
 \end{aligned}$$

where the third equality follows from the fact that $\mathcal{A}^{-1} - \mathcal{B}^{-1} = \mathcal{A}^{-1}(\mathcal{B} - \mathcal{A})\mathcal{B}^{-1}$ for invertible matrices \mathcal{A} , \mathcal{B} . Next, write (14) as

$$\begin{aligned}
 y_N &= \frac{\lambda}{1-p} H_N y_N + X_N\beta + \varepsilon_N + \lambda \left(\tilde{G}_N - \frac{H_N}{1-p} \right) y_N + \lambda \Delta_N y_N \\
 &= \frac{\lambda}{1-p} H_N y_N + X_N\beta + \varepsilon_N + \underbrace{\lambda \left(\tilde{G}_N - \frac{H_N}{1-p} \right) \tilde{y}_N}_{\equiv v_N} + \underbrace{\lambda^2 \left(\tilde{G}_N - \frac{H_N}{1-p} \right) (I_N - \lambda \tilde{G}_N)^{-1} \Delta_N y_N + \lambda \Delta_N y_N}_{\equiv u_N},
 \end{aligned}$$

where the second equality holds because we substitute y_N in $\lambda \left(\tilde{G}_N - \frac{H_N}{1-p} \right) y_N$ using the r.h.s. of (1). Furthermore, we can write

$$u_N = \left[\lambda \left(\tilde{G}_N - \frac{H_N}{1-p} \right) (I_N - \lambda \tilde{G}_N)^{-1} + I_N \right] \lambda \Delta_N y_N = \left(I_N - \lambda \frac{H_N}{1-p} \right) \left(I_N - \lambda \tilde{G}_N \right)^{-1} \lambda \Delta_N y_N.$$

This establishes (15).

Next, we introduce the technical regularity conditions needed for establishing the asymptotic properties in Proposition 6. Suppose $I_N - \lambda G_N$ and $I_N - \lambda \tilde{G}_N$ are invertible almost surely, and denote $M_N \equiv (I_N - \lambda G_N)^{-1}$, $\tilde{M}_N \equiv (I_N - \lambda \tilde{G}_N)^{-1}$. Let $\tilde{R}_{N,s} \equiv (H_{N,s} \tilde{M}_{N,s} X_{N,s}, X_{N,s})$.

- (S-REG)** (i) For all i , $\sup_i \left[\sum_j |M_{N,ij}| \right] < \infty$; $\sup_j E (|X_{N,j} \beta| + |\varepsilon_{N,j}| | \Delta_N) < \infty$;
 $\sup_j \left| \left(X'_N H_N^2 \tilde{M}_N \right)_{ij} \right| < \infty$ and $\sup_j \left| \left(X'_N H_N \tilde{M}_N \right)_{ij} \right| < \infty$ almost surely.
(ii) $(H_{N,s}, \tilde{G}_{N,s}, X_{N,s}, \varepsilon_{N,s})$ are independent across $s = 1, 2, \dots, S$.
(iii) There exist $\delta > 0$ s.t. for all s , $E \left[\|Z'_{N,s} \tilde{R}_{N,s}\|^{1+\delta} \right]$, $E \left[\|Z'_{N,s} H_{N,s} \tilde{M}_{N,s} \varepsilon_{N,s}\|^{1+\delta} \right]$, and $E \left(\|Z'_{N,s} Z_{N,s}\|^{1+\delta} \right)$ are uniformly bounded.
(iv) For some $\delta > 0$, $E \|Z'_{N,s} v_{N,s}\|^{2+\delta} < \Delta < \infty$ and $S^{-1} \sum_{s=1}^S \text{Var}(Z'_{N,s} v_{N,s}) > \delta' > 0$ for S sufficiently large.
(v) $\sup_j \left| \left[\left(I_N - \lambda \frac{H_N}{1-p} \right) \tilde{M}_N \right]_{ij} \right| < \infty$ for all i almost surely.
(vi) $\lim_{S \rightarrow \infty} \frac{1}{S} \sum_{s=1}^S E (Z'_{N,s} Z_{N,s})$ and $\lim_{S \rightarrow \infty} \frac{1}{S} \sum_{s=1}^S E \left(Z'_{N,s} \tilde{R}_{N,s} \right)$ exist and are non-singular.

Assumption (S-REG) collects regularity conditions needed for deriving the asymptotic properties of $\hat{\theta}_a - \theta_a$. Part (ii) implies that exogenous variables are drawn independently across the blocks. Part (i) and (v) introduce bound conditions on exogenous arrays in the model. These allow us to relate differences between y_N and its near-block diagonal approximation \tilde{y}_N to the order of difference between G_N and \tilde{G}_N . Parts (iii) and (iv) are boundedness conditions on population moments that ensure a law of large numbers and a central limit theorem apply to components of the estimator.

Lemma A1. *Let a_N, b_N be random vectors in \mathbb{R}^N . Suppose there exist constants $C_1, C_2 < \infty$ such that $\Pr\{\sup_{i \leq N} E(|a_i| | \Delta_N) \leq C_1\} = 1$ and $\Pr\{\sup_{j \leq N} E(|b_j| | \Delta_N) \leq C_2\} = 1$.*

Then Assumption S-LOB implies $\frac{1}{S}a'_N\Delta_N b_N = O_p(S^{\rho-1})$.

Proof of Lemma A1. From Assumption S-LOB, $\sum_i \sum_j E|\Delta_{N,ij}| = O(S^\rho)$ for some $\rho < 1$. By construction,

$$\begin{aligned} E\left(\left|\frac{1}{S}a'_N\Delta_N b_N\right|\right) &\leq \frac{1}{S}E\left[\sup_{i,j} E(|a_i b_j| \mid \Delta_N) \cdot \left(\sum_i \sum_j |\Delta_{N,ij}|\right)\right] \\ &\leq \frac{1}{S}E\left[C_1 C_2 \left(\sum_i \sum_j |\Delta_{N,ij}|\right)\right] = O(S^{\rho-1}). \end{aligned}$$

It then follows that $\frac{1}{S}a'_N\Delta_N b_N = O_p(S^{\rho-1})$. \square

Lemma A2. *Under the conditions in (S-REG)-(i), there exists a constant $C^* < \infty$ such that $\Pr\{\sup_{i \leq N} E(|y_i| \mid \Delta_N) \leq C^*\} = 1$ for all N .*

Proof of Lemma A2. Let $M_N \equiv (I_N - \lambda G_N)^{-1}$. For any matrix \mathcal{A} , let $\mathcal{A}_{(i)}$ denote its i -th row; and \mathcal{A}_{ij} denote its (i, j) -th component. It then follows from the reduced form that

$$\begin{aligned} \sup_{i \leq N} E(|y_{N,i}| \mid \Delta_N) &= \sup_i E\left(\left|\sum_j M_{N,ij}(X_{N,(j)}\beta + \varepsilon_j)\right| \mid \Delta_N\right) \\ &\leq \sup_i \left[\sum_j |M_{N,ij}|\right] \times \sup_j E(|X_{N,(j)}\beta| + |\varepsilon_{N,j}| \mid \Delta_N). \end{aligned}$$

Hence, there exists some constant $C^* < \infty$ with $\Pr\{\sup_i E(|y_i| \mid \Delta_N) \leq C^*\} = 1$. \square

Lemma A3. *Under the conditions in (S-REG), $\frac{1}{S}R'_N Z_N = A_0 + o_p(1)$, $\frac{1}{S}Z'_N Z_N = B_0 + o_p(1)$, and $\frac{1}{S}Z'_N v_N = O_p(S^{-1/2})$.*

Proof of Lemma A3. By definition, $\frac{1}{S}Z'_N Z_N = \frac{1}{S} \sum_{s=1}^S Z'_{N,s} Z_{N,s}$, with $Z_{N,s}$ independent across s due to (S-REG)-(ii). Then by (S-REG)-(iii) and the law of large numbers for independent and heterogeneously distributed observations (e.g., Corollary 3.9 in White (2001)), $\frac{1}{S}Z'_N Z_N = B_0 + o_p(1)$ where $B_0 \equiv \lim_{S \rightarrow \infty} \frac{1}{S} \sum_{s=1}^S E(Z'_{N,s} Z_{N,s})$. Next, note by construction and (1),

$$\frac{1}{S}Z'_N R_N = \frac{1}{S} \begin{pmatrix} X'_N H_N^2 \tilde{y}_N & X'_N H_N X_N \\ X'_N H_N \tilde{y}_N & X'_N X_N \end{pmatrix} + \frac{1}{S} \lambda \begin{pmatrix} X'_N H_N^2 \tilde{M}_N \Delta_N y_N & 0 \\ X'_N H_N \tilde{M}_N \Delta_N y_N & 0 \end{pmatrix}. \quad (2)$$

By (S-REG)-(i) and Lemma A2, y_N satisfies the condition on b_N in Lemma A1. It then follows from Lemma A1 that the *second* term on the right-hand side of (2) is $O_p(S^{\rho-1})$. Besides, the *first* term on the r.h.s. of (2) is

$$\frac{1}{S} \sum_{s=1}^S Z'_{N,s} \tilde{R}_{N,s} + \frac{1}{S} \sum_{s=1}^S \left(Z'_{N,s} H_{N,s} \tilde{M}_{N,s} \varepsilon_{N,s}, \mathbf{0} \right). \quad (3)$$

By (N3), $E \left(Z'_{N,s} H_{N,s} \tilde{M}_{N,s} \varepsilon_{N,s} \right) = 0$. It then follows from (S-REG)-(iii) that the expression in (3) is $A_0 + o_p(1)$, with $A_0 \equiv \lim_{S \rightarrow \infty} \frac{1}{S} \sum_{s=1}^S E \left(Z'_{N,s} \tilde{R}_{N,s} \right)$.

Next, note that by definition,

$$\frac{1}{S} Z'_N v_N = \frac{1}{S} \sum_{s=1}^S Z'_{N,s} \varepsilon_{N,s} + \lambda \frac{1}{S} \sum_{s=1}^S Z'_{N,s} \left(\tilde{G}_{N,s} - \frac{H_{N,s}}{1-p} \right) \tilde{y}_{N,s}. \quad (4)$$

By construction, $Z_{N,s}$, $\varepsilon_{N,s}$, $\tilde{G}_{N,s}$ and $H_{N,s}$ are independent across blocks $s = 1, 2, \dots, S$. Also, recall that $\tilde{y}_{N,s}$ is defined as $\tilde{y}_{N,s} \equiv (I_s - \lambda \tilde{G}_{N,s})^{-1} (X_{N,s} \beta + \varepsilon_{N,s})$, Hence $\tilde{y}_{N,s}$ is also independent across the blocks. Assumption (N3) implies $E(Z'_{N,s} \varepsilon_{N,s}) = 0$; Assumptions (N1) and (N2) imply

$$E \left(H_{N,s} | \tilde{G}_{N,s}, X_{N,s} \right) = (1-p) \tilde{G}_{N,s}.$$

Furthermore, the same argument as in the proof of Proposition 2 shows that under (N1), (N2), (N3) and (N4)

$$E \left(H_{N,s}^2 | \tilde{G}_{N,s}, X_{N,s} \right) = (1-p) E \left(H_{N,s} \tilde{G}_{N,s} | \tilde{G}_{N,s}, X_{N,s} \right),$$

so that

$$E \left[Z'_{N,s} \left(\tilde{G}_{N,s} - \frac{H_{N,s}}{1-p} \right) \tilde{y}_{N,s} \right] = 0.$$

It then follows from (S-REG)-(iv) and the Central Limit Theorem that $\frac{1}{S} Z'_N v_N = O_p(S^{-1/2})$. \square

Proof of Proposition 6. As shown in Lemma A3, $\frac{1}{S} R'_N Z_N = A_0 + o_p(1)$, $\frac{1}{S} Z'_N Z_N = B_0 + o_p(1)$, and $\frac{1}{S} Z'_N v_N = O_p(S^{-1/2})$ under (N1)-(N4), (S-LOB) and (S-REG). Furthermore, with (S-REG)-(v), Lemma A1 and Lemma A2 imply that $\frac{1}{S} Z'_N u_N = O_p(S^{\rho-1})$. When

$\rho < 1/2$, we have

$$\frac{1}{\sqrt{S}} Z'_N (u_N + v_N) \xrightarrow{d} \frac{1}{\sqrt{S}} Z'_N v_N \xrightarrow{d} \mathcal{N}(0, \omega_0),$$

where $\omega_0 = \lim_{S \rightarrow \infty} \frac{1}{S} \sum_{s=1}^S E (Z'_{N,s} v_{N,s} v'_{N,s} Z_{N,s})$. Hence,

$$\begin{aligned} \sqrt{S}(\hat{\theta}_a - \theta_a) &= (A'_0 B_0^{-1} A_0)^{-1} A'_0 B_0^{-1} \frac{1}{\sqrt{S}} Z'_N v_N + o_p(1) \\ &\xrightarrow{d} \mathcal{N}(0, (A'_0 B_0^{-1} A_0)^{-1} A'_0 B_0^{-1} \omega_0 B_0^{-1} A_0 (A'_0 B_0^{-1} A_0)^{-1}). \end{aligned}$$

□

B. Near-epoch Dependence (NED) in Large Networks

We consider a scenario in Section 6.2 of Lewbel, Qu and Tang (2022), where the sample contains a single large network with size N which does not admit any definition of approximate groups. Nonetheless, there is some notion of latent distance between individuals $i = 1, 2, \dots, N$ on the network. The distance and its associated space are formally defined in the following assumption.

(W0) The lattice $D \subset \mathbb{R}^{d_0}$, $d_0 \geq 1$, is infinitely countable. The location $l : \{1, \dots, N\} \rightarrow D_N \subset D$ is a mapping of individual i to its location $l(i) \in D_N \subset D \subset \mathbb{R}^{d_0}$. All elements in D are located at distances of at least $c_0 > 0$ from each other; w.l.o.g. let $c_0 = 1$, i.e., $\forall l(i), l(j) \in D : d_{ij} \geq 1$, where d_{ij} is the distance between individuals i and j .

This assumption was first used in Jenish and Prucha (2009, 2012), and later in Qu and Lee (2015), and Qu et al. (2017) to establish an *increasing domain* asymptotic environment, in which growth of the sample is ensured by an unbounded expansion of the sample space.

As before, denote the network structure, the individual outcomes and characteristics by $G_N \in \{0, 1\}^{N \times N}$, $y_N \in \mathbb{R}^{N \times 1}$, $X_N \in \mathbb{R}^{N \times K}$. To reiterate, the scenario we consider here in Section 6.2 differs from the case in Section 6.1 in that G_N is generally not near block-diagonal, and there is no clear group structure in H_N .

Our goal is to show that some notion of weak dependence (a.k.a. near-epoch dependence) exists between individual outcomes. Therefore, certain law of large numbers and

central limit theorem applies, so that the 2SLS estimand is the same as in the case with many independent groups with fixed sizes. While doing so, we allow for flexible network structures, and refrain from modeling the link formation process in G_N . Instead, we maintain the following requirement on the strength of links:

- (W1) (i) When $d_{ij} > c$, $G_{N,ij} = 0$, where c is a finite constant;
- (ii) $\sup_N \|\lambda G_N\|_\infty = \rho < 1$, where $\|G_N\|_\infty = \sup_i \sum_{j=1}^N |G_{N,ij}|$ is the row sum of G_N .

Assumption (W1) (i) states that there are no links between two individuals between whom the distance exceeds some threshold c . This distance and its associated space are determined by the link formation process. For example, links may only exist between individuals with similar ages and educational backgrounds. This assumption implies the matrix G_N is uniformly bounded in both row and column sums for all N , i.e., each individual can have only finite number of friends; (ii) restricts the strength of peer effects, i.e., $|\lambda| < 1/\sup_N \|G_N\|_\infty$, which is a sufficient condition for the invertibility and uniformly row and column boundedness of $M_N \equiv (I - \lambda G_N)^{-1}$.

We maintain the following exogeneity and independence conditions which are analogous to (A2), (A3) and (A4) for the case in Section 3. Note we add independence of $\varepsilon_{i,N}$ across i to replace to independence across small, well-defined groups.¹

$$(W2) E(H_{N,ij}|G_{N,ij} = 1, G_N, X_N) = 1 - p \text{ and } E(H_{ij}|G_{N,ij} = 0, G_N, X_N) = 0, \forall i \neq j ;$$

$$(W3) E(\varepsilon_N|X_N, G_N, H_N) = 0 \text{ and } \varepsilon_{i,N} \text{ is independent across all } i = 1, \dots, N;$$

$$(W4) \text{ Conditional on } (G_N, X_N), H_{N,ij} \perp H_{N,kl} \text{ for all } (i, j) \neq (k, l).$$

The structural model is the same as (14). A 2SLS estimator that uses $Z_N \equiv (H'_N X_N, X_N)$

¹This independence of $\varepsilon_{i,N}$ across i can be relaxed to the moving-average type of dependence, i.e., $\varepsilon_{i,N}$ is correlated with $\varepsilon_{j,N}$ only if i and j are within a finite distance.

as instruments for $R_N \equiv (H_N y_N, X_N)$ is:

$$\begin{aligned}\widehat{\theta}_a &= [R'_N Z_N (Z'_N Z_N)^{-1} Z'_N R_N]^{-1} R'_N Z_N (Z'_N Z_N)^{-1} Z'_N y_N \\ &= \theta_a + [R'_N Z_N (Z'_N Z_N)^{-1} Z'_N R_N]^{-1} R'_N Z_N (Z'_N Z_N)^{-1} Z'_N v_N,\end{aligned}$$

where $v_N = \varepsilon_N + \lambda \left(G_N - \frac{H_N}{1-p} \right) y_N$.

While analyzing the asymptotic properties of $\frac{1}{N} R'_N Z_N$, $\frac{1}{N} Z'_N Z_N$, and $\frac{1}{\sqrt{N}} Z'_N v_N$, we need to take into account the dependence between $y_{N,i}$ (and hence $v_{N,i}$) across individuals $i = 1, \dots, N$ through the reduced form $y_N = M_N(X_N \beta + \varepsilon_N)$. To do so, we exploit a law of large numbers (LLN) and a central limit theorem (CLT) under a notion of *near-epoch dependence* (NED), as introduced in Jenish and Prucha (2009) and Jenish and Prucha (2012) for spatial econometric models. In Online Appendix C, we first establish the NED of $R_{N,i}$, $Z_{N,i}$, and $v_{N,i}$ separately (Lemma C1). The intuition is that though $y_{N,i}$ (and $v_{N,i}$) are correlated across individuals, it can be approximated by $\varepsilon_{N,j}$'s where j is within certain distance from i conditional on (X_N, G_N) . Hence, if the latent distance between individuals i_1 and i_2 are larger than that distance, then y_{N,i_1} and y_{N,i_2} (as well as v_{N,i_1} and v_{N,i_2}) are almost independent conditional on (X_N, G_N) . Then, using the property that the product of NED terms still satisfy NED, we show the NED of $R'_{N,i} Z_{N,i}$, $Z_{N,i} Z'_{N,i}$, and $Z_{N,i} v_{N,i}$ (Lemma C2). Under (W2) (ii), the NED rates in $R'_{N,i} Z_{N,i}$, $Z_{N,i} Z'_{N,i}$, and $Z_{N,i} v_{N,i}$ satisfy the requirement needed for the LLNs and CLT. Hence, in Lemma C3, we show that

$$\begin{aligned}\frac{1}{N} R'_N Z_N - \frac{1}{N} E(R'_N Z_N | X_N, G_N) &= o_p(1); \quad \frac{1}{N} Z'_N Z_N - \frac{1}{N} E(Z'_N Z_N | X_N, G_N) = o_p(1); \\ \frac{1}{\sqrt{N}} Z'_N v_N &\xrightarrow{d} \mathcal{N}(0, \omega_c), \quad \text{where } \omega_c = \lim_{N \rightarrow \infty} \frac{1}{N} E(Z'_N v_N v'_N Z_N | X_N, G_N).\end{aligned}$$

Proposition B1. *Suppose (W0), (W1), (W2), (W3) and (W4) hold. If Assumption (W-REG) hold, then,*

$$\widehat{\theta}_a - \theta_a = o_p(1) \text{ and } \sqrt{N} \left(\widehat{\theta}_a - \theta_a \right) \xrightarrow{d} \mathcal{N}(0, \Omega_c)$$

where $\Omega_c \equiv (A'_c B_c^{-1} A_c)^{-1} A'_c B_c^{-1} \omega_c B_c^{-1} A_c (A'_c B_c^{-1} A_c)^{-1}$ with

$$A_c = \lim_{N \rightarrow \infty} \frac{1}{N} E(R'_N Z_N | G_N, X_N) \text{ and } B_c = \lim_{N \rightarrow \infty} \frac{1}{N} E(Z'_N Z_N | G_N, X_N).$$

Next, consider the estimation of p . From (W4), $H_{N,ij}$ is independent of each other conditional on (X_N, G_N) , so the conventional LLNs and CLT can be directly applied to

$$\frac{1}{N} \sum_i \sum_j H_{N,ij}.$$

Here, we rescale $\sum_i \sum_j H_{N,ij}$ using N , instead of N^2 , because $\sum_i \sum_j H_{N,ij} = O(N)$ from the uniform boundedness in the row and column sum of G_N . Thus we can estimate p using a procedure analogous to that in Section 3.

Notice in this section the asymptotic results are all conditional on (X_N, G_N) , so randomness comes from ε_N and H_N . If we further want to check the unconditional asymptotic distribution, we need to impose weak dependence in the data-generating process of (X_N, G_N) . For example, $X_{N,i}$ is only correlated with $X_{N,j}$ within a finite distance; $G_{N,ij}$ is only correlated with $G_{N,kl}$ within individuals k and l are within a finite distance of i and j .

C. Proofs for NED in Large Networks

Recall that the structural form is:

$$\begin{aligned} y_N &= \lambda G_N y_N + X_N \beta + \varepsilon_N \\ &= \frac{\lambda}{1-p} H_N y_N + X_N \beta + \varepsilon_N + \underbrace{\lambda \left(G_N - \frac{H_N}{1-p} \right) y_N}_{v_N}. \end{aligned}$$

A 2SLS estimator that uses $Z_N \equiv (H'_N X_N, X_N)$ as instruments for $R_N \equiv (H_N y_N, X_N)$ is:

$$\hat{\theta}_a = (A'_N B_N^{-1} A_N)^{-1} A'_N B_N^{-1} Z'_N y_N.$$

where $A_N \equiv Z'_N R_N$ and $B_N \equiv Z'_N Z_N$. By definition,

$$\widehat{\theta}_a - \theta_a = (A'_N B_N^{-1} A_N)^{-1} A'_N B_N^{-1} Z'_N v_N.$$

We collect all regularity conditions needed for asymptotic properties of $\widehat{\theta}_a - \theta_a$ in Assumption (W-REG).

Throughout Online Appendix C, we use the first subscripts to index the individuals, e.g. $y_{i,N}$, $\varepsilon_{i,N}$ (instead of $y_{N,i}$, $\varepsilon_{N,i}$ as in the text).

(W-REG) (i) $\lim_{N \rightarrow \infty} \frac{1}{N} E(Z'_N Z_N | G_N, X_N)$ and $\lim_{N \rightarrow \infty} \frac{1}{N} E(Z'_N R_N | G_N, X_N)$ exist and are non-singular;

(ii) $\sup_{i,N} |X_{i,N} \beta| < \infty$;

(iii) ε_N is independent of (G_N, H_N, X_N) with $\sup_{i,N} E(\varepsilon_{i,N}^4) < \infty$.

Parts (i) bounds population moments conditional on (G_N, X_N) , which is needed for a law of large numbers and a central limit theorem apply to components of the estimator. Part (ii) requires the uniform boundedness of elements in the regressor X_N , which is stronger than the usual linear regression because we allow for the correlation between network matrix G_N (and H_N) with X_N . If G_N (H_N) is independent of X_N , then part (ii) can be relaxed to boundedness of $\sup_{i,N} E|X_{i,N} \beta|^4$. Part (iii) introduces the independence between ε_N and (G_N, H_N, X_N) and the boundedness of its fourth moments, which is also stronger than the usual linear model. The independence can be relaxed to the conditional mean independence on the first fourth moments of ε_N .

Our central limit theorem of the 2SLS estimator is based on inference under NED. The following notion of NED for random fields is from Jenish and Prucha (2012).

Definition 1. For any random vector Y , $\|Y\|_p = [E(|Y|^p)]^{1/p}$ denotes its L_p -norm where $|Y|$ is the Euclidean norm of Y . Denote $\mathcal{F}_{i,N}(r)$ as a σ -field generated by the random vectors $\varsigma_{j,N}$'s located within the ball $B_i(r)$, which is a ball centered at the location i with a radius r in a d_0 -dimensional Euclidean space D .

Definition 2. (Near-epoch Dependence) Let $T = \{T_{i,N}, i \in D_N, N \geq 1\}$ and $\varsigma = \{\varsigma_{i,N}, i \in D_N, N \geq 1\}$ be random fields with $\|T_{i,N}\|_p < \infty$, $p \geq 1$, where $D_N \subset D$ and $|D_N| \rightarrow \infty$

as $N \rightarrow \infty$, and let $d = \{d_{i,N}, i \in D_N, N \geq 1\}$ be an array of finite positive constants. Then the random field T is said to be L_p -near-epoch dependent on the random field ς if $\|T_{i,N} - E(T_{i,N} | \mathcal{F}_{i,N}(r))\|_p \leq d_{i,n} \varphi(r)$ for some sequence $\varphi(r) \geq 0$ such that $\lim_{s \rightarrow \infty} \varphi(r) = 0$. The quantity $\varphi(r)$, which is, without loss of generality, assumed to be non-increasing, is called the NED coefficient (or rate), and the $d_{i,n}$'s are called NED scaling factors. The random field T is said to be L_p -NED on ς of size $-\alpha$ if $\varphi(r) = O(r^{-\alpha})$ for some $\mu > \alpha > 0$. Furthermore, if $\sup_N \sup_{i \in D_N} d_{i,N} < \infty$, then T is said to be uniformly L_p -NED on ς . If $\varphi(r) = O(\rho^r)$, where $0 < \rho < 1$, then T is called geometrically L_p -NED on ς .

To establish the consistency and asymptotic distribution of 2SLS estimator, we first explore the sparsity in G_N^{l+1} .

Claim 1. Under Assumption (W0) and (W1), the (i, j) -th entry of G_N^{l+1} or $H_N G_N^l$ or H_N^{l+1} is zero unless $j \in B_i(lc)$.

Proof of Claim 1. We can express

$$(G_N^{l+1})_{ij} = \sum_{j_1} \sum_{j_2} \cdots \sum_{j_l} G_{N,ij_1} G_{N,j_1j_2} \cdots G_{N,j_lj},$$

i.e., l steps from i to j . By the triangle inequality (because D_N is a lattice in a Euclidean space),

$$d_{ij} \leq d_{ij_1} + d_{j_1j_2} + \cdots + d_{j_lj}.$$

If $j \notin B_i(lc)$, i.e., $d_{ij} > lc$, then there is at least one distance out of $d_{ij_1}, d_{j_1j_2}, \dots, d_{j_lj}$ greater than c (say $d_{j_1j_2} > c$) and that step in G_N must be zero (i.e., $G_{N,j_1j_2} = 0$). Hence, $(G_N^{l+1})_{ij} \neq 0$ only if $j \in B_i(lc)$. The same argument applies to $H_N G_N^l$ and H_N^{l+1} . \square

Lemma C1. Under Assumptions (W0), (W1), (W2), (W3), and (W4), and (W-REG) (ii) and (iii), we have the L_4 -NED of $R_{i,N}$, $Z_{i,N}$, and $v_{i,N}$ with the NED rate $\varphi(r) = \rho^{r/c}$, where $\rho = \|\lambda G_N\|_\infty$ and c is the threshold distance that $G_{ij} = 0$ if $d_{ij} > c$.

Proof of Lemma C1. Let

$$\begin{aligned}
(H_N y_N)_i &= \sum_{j=1}^N (H_N M_N)_{ij} (X_{j,N} \beta + \varepsilon_{j,N}) = \sum_{j=1}^N \sum_{l=0}^{\infty} \lambda^l (H_N G_N^l)_{ij} (X_{j,N} \beta + \varepsilon_{j,N}) \\
&= \underbrace{\sum_{j=1}^N \sum_{l=0}^{r/c} \lambda^l (H_N G_N^l)_{ij} (X_{j,N} \beta + \varepsilon_{j,N})}_{t_{1i,N}} + \underbrace{\sum_{j=1}^N \sum_{l=r/c+1}^{\infty} \lambda^l (H_N G_N^l)_{ij} (X_{j,N} \beta + \varepsilon_{j,N})}_{t_{2i,N}}.
\end{aligned}$$

We will show the L_4 -NED for $t_{1i,N}$ and $t_{2i,N}$ on $\varepsilon_{i,N}$ separately. Let $\mathcal{F}_{i,N}(r)$ be a σ -field generated by $\varepsilon_{j,N}$'s located within the ball $B_i(r)$.

$$\begin{aligned}
& \|t_{1i,N} - E(t_{1i,N} | \mathcal{F}_{i,N}(r))\|_4 \\
&= \left\| \sum_{j=1}^N \sum_{l=0}^{r/c} \lambda^l [(H_N G_N^l)_{ij} (X_{j,N} \beta + \varepsilon_{j,N}) - E((H_N G_N^l)_{ij} (X_{j,N} \beta + \varepsilon_{j,N}) | \mathcal{F}_{i,N}(r))] \right\|_4 \\
&= \left\| \sum_{j \in B_i(lc)} \sum_{l=0}^{r/c} \lambda^l [(H_N G_N^l)_{ij} (X_{j,N} \beta + \varepsilon_{j,N}) - E((H_N G_N^l)_{ij} (X_{j,N} \beta + \varepsilon_{j,N}) | \mathcal{F}_{i,N}(r))] \right\|_4 = 0,
\end{aligned}$$

where the second “=” holds due to Claim 1 ($(H_N G_N^l)_{ij} \neq 0$ only if $j \in B_i(lc)$); the last “=” holds because when $l \leq r/c$, i.e., $r \geq lc$, all random variables involved in $(H_N G_N^l)_{ij} (X_{j,N} \beta + \varepsilon_{j,N})$ are contained by $\mathcal{F}_{i,N}(r)$ and hence, for $j \in B_i(lc)$ and $l \leq r/c$,

$$(H_N G_N^l)_{ij} (X_{j,N} \beta + \varepsilon_{j,N}) - E[(H_N G_N^l)_{ij} (X_{j,N} \beta + \varepsilon_{j,N}) | \mathcal{F}_{i,N}(r)] = 0.$$

As for $t_{2i,N}$,

$$\begin{aligned}
& \sup_{i,N} \|t_{2i,N} - E(t_{2i,N} | \mathcal{F}_{i,N}(r))\|_4 \\
&= \sup_{i,N} \left\| \sum_{j=1}^N \sum_{l=r/c+1}^{\infty} \lambda^l [(H_N G_N^l)_{ij} (X_{j,N} \beta + \varepsilon_{j,N}) - E((H_N G_N^l)_{ij} (X_{j,N} \beta + \varepsilon_{j,N}) | \mathcal{F}_{i,N}(r))] \right\|_4 \\
&\leq 2 \sup_{i,N} \left\| \sum_{j=1}^N \sum_{l=r/c+1}^{\infty} \lambda^l (H_N G_N^l)_{ij} (X_{j,N} \beta + \varepsilon_{j,N}) \right\|_4 \\
&\leq 2 \sum_{l=r/c+1}^{\infty} |\lambda|^l \sup_{i,N} \left(\left\| \sum_{j=1}^N (H_N G_N^l)_{ij} X_{j,N} \beta \right\|_4 + \left\| \sum_{j=1}^N (H_N G_N^l)_{ij} \varepsilon_{j,N} \right\|_4 \right) \\
&\leq 2 \sum_{l=r/c+1}^{\infty} |\lambda|^l \left(\sup_{j,N} |X_{j,N} \beta| + \sup_{j,N} \|\varepsilon_{j,N}\|_4 \right) \cdot \sup_{i,N} \left\| \sum_{j=1}^N |(H_N G_N^l)_{ij}| \right\|_4 \\
&\leq 2 \sum_{l=r/c+1}^{\infty} C |\lambda|^l \cdot \sup_N \|G_N\|_{\infty}^{l+1} \leq 2C \sum_{l=r/c+1}^{\infty} |\rho|^l \leq C^* \rho^{r/c},
\end{aligned}$$

where C and C^* are some generic large, but finite constants, the first “ \leq ” holds due to Minkowski and the conditional Jensen inequalities; the third “ \leq ” holds because $\varepsilon_{j,N}$ is independent of (G_N, H_N) ; the fourth “ \leq ” holds because $\sum_{j=1}^N |(H_N G_N^l)_{ij}| \leq \|G_N\|_{\infty}^{l+1}$.

Since $(H_N y_N)_i = t_{1i,N} + t_{2i,N}$, we have the L_4 -NED of $(H_N y_N)_i$ such that

$$\begin{aligned}
& \sup_{i,N} \|(H_N y_N)_i - E((H_N y_N)_i | \mathcal{F}_{i,N}(r))\|_4 \\
&\leq \sup_{i,N} \|t_{1i,N} - E(t_{1i,N} | \mathcal{F}_{i,N}(r))\|_4 + \sup_{i,N} \|t_{2i,N} - E(t_{2i,N} | \mathcal{F}_{i,N}(r))\|_4 \leq C^* \rho^{r/c}.
\end{aligned}$$

Same arguments can be applied to show

$$\sup_{i,N} \|(G_N y_N)_i - E((G_N y_N)_i | \mathcal{F}_{i,N}(r))\|_4 \leq C^* \rho^{r/c}.$$

Then, with

$$v_{i,N} = \frac{1}{1-p} (H_N y_N)_i - (G_N y_N)_i,$$

we have

$$\sup_{i,N} \|v_{i,N} - E(v_{i,N}|\mathcal{F}_{i,N}(r))\|_4 \leq C^* \rho^{r/c}.$$

Similar arguments can be applied to terms in Z_N and R_N , i.e.,

$$\sup_{i,N} \|R_{i,N} - E(R_{i,N}|\mathcal{F}_{i,N}(r))\|_4 \leq C^* \rho^{r/c} \text{ and } \sup_{i,N} \|Z_{i,N} - E(Z_{i,N}|\mathcal{F}_{i,N}(r))\|_4 \leq C^* \rho^{r/c}.$$

□

Then, by applying the following claim in Qu and Lee (2015) on the product of two NED terms, we can establish the L_2 -NED of $R_{i,N}Z'_{i,N}$, $Z_{i,N}Z'_{i,N}$, and $R_{i,N}v_{i,N}$.

Claim 2. *If $\|t_{1i,N} - E(t_{1i,N}|\mathcal{F}_{i,N}(r))\|_4 \leq C_1\varphi_1(r)$ and $\|t_{2i,N} - E(t_{2i,N}|\mathcal{F}_{i,N}(r))\|_4 \leq C_2\varphi_2(r)$, with $\max(\|t_{1i,N}\|_4, \|t_{2i,N}\|_4) \leq C$, then $\|t_{1i,N}t_{2i,N} - E(t_{1i,N}t_{2i,N}|\mathcal{F}_{i,N}(r))\|_2 \leq C(C_1 + C_2)\varphi(r)$, where $\varphi(r) = \max(\varphi_1(r), \varphi_2(r))$.*

Lemma C2. *Under Assumptions (W0), (W1), (W2), (W3), and (W4), and (W-REG) (ii) and (iii), we have the L_2 -NED of $R_{i,N}Z'_{i,N}$, $Z_{i,N}Z'_{i,N}$, and $R_{i,N}v_{i,N}$ with the NED rate $\varphi(r) = \rho^{r/c}$ the same as Lemma C1.*

Proof of Lemma C2. First, we show that $\max(\|R_{i,N}\|_4, \|Z_{i,N}\|_4, \|v_{i,N}\|_4) \leq C$. Again, the most complicated term is $(H_N y_N)_i$.

$$\begin{aligned} \sup_{i,N} \|(H_N y_N)_i\|_4 &= \sup_{i,N} \left\| \sum_{j=1}^N \sum_{l=0}^{\infty} \lambda^l (H_N G_N^l)_{ij} (X_{j,N} \beta + \varepsilon_{j,N}) \right\|_4 \\ &\leq \sup_{i,N} \sum_{l=0}^{\infty} |\lambda|^l \left(\left\| \sum_{j=1}^N (H_N G_N^l)_{ij} X_{j,N} \beta \right\|_4 + \left\| \sum_{j=1}^N (H_N G_N^l)_{ij} \varepsilon_{j,N} \right\|_4 \right) \\ &\leq \sum_{l=0}^{\infty} |\lambda|^l \left(\sup_{j,N} |X_{j,N} \beta| \cdot \sup_{j,N} \|\varepsilon_{j,N}\|_4 \right) \cdot \sup_{i,N} \left\| \sum_{j=1}^N |(H_N G_N^l)_{ij}| \right\|_4 \\ &\leq \sup_N C \sum_{l=0}^{\infty} |\lambda|^l \|G_N\|_{\infty}^l \leq C^* < \infty. \end{aligned}$$

Same arguments can be applied to other terms in $R_{i,N}$, $Z_{i,N}$, and $v_{i,N}$, so we have uniformly L_4 boundedness of $R_{i,N}$, $Z_{i,N}$, and $v_{i,N}$. By Cauchy-Schwarz inequality, we have uniformly

L_4 boundedness of $R_{i,N}Z'_{i,N}$, $Z_{i,N}Z'_{i,N}$, and $R_{i,N}v_{i,N}$ that

$$\sup_{i,N} \|R_{i,N}Z'_{i,N}\|_2 < \infty; \quad \sup_{i,N} \|Z_{i,N}Z'_{i,N}\|_2 < \infty; \quad \sup_{i,N} \|R_{i,N}v_{i,N}\|_2 < \infty.$$

Then, applying Claim 2, we have

$$\begin{aligned} \sup_{i,N} \|R_{i,N}Z'_{i,N} - E(R_{i,N}Z'_{i,N}|\mathcal{F}_{i,N}(r))\|_2 &\leq C^*\varphi(r); \\ \sup_{i,N} \|Z_{i,N}Z'_{i,N} - E(Z_{i,N}Z'_{i,N}|\mathcal{F}_{i,N}(r))\|_2 &\leq C^*\varphi(r); \\ \sup_{i,N} \|R_{i,N}v_{i,N} - E(R_{i,N}v_{i,N}|\mathcal{F}_{i,N}(r))\|_2 &\leq C^*\varphi(r). \end{aligned}$$

□

From Jenish and Prucha (2012), we have the following two claims for LLN and CLT under NED.

Claim 3. *Under Assumption (W0), if the random field $\{T_{i,N}, i \in D_N, N \geq 1\}$ is L_1 -NED, the base $\{\varsigma_{i,N}\}$'s are i.i.d., and $\{T_{i,N}\}$'s are uniformly L_p bounded for some $p > 1$, then $\frac{1}{N} \sum_{i=1}^N (T_{i,N} - ET_{i,N}) \xrightarrow{L_1} 0$, where “ $\xrightarrow{L_1}$ ” denotes convergence in L_1 -norm.*

Claim 4. *Let $\{T_{i,N}, i \in D_N, N \geq 1\}$ be a random field that is L_2 -NED on an i.i.d. random field ς . If Assumption (W1) and the following conditions are met:*

- (1) $\{T_{i,N}, i \in D_N, N \geq 1\}$ is uniformly $L_{2+\delta}$ -bounded for some $\delta > 0$,
 - (2) $\inf_N \frac{1}{N} \sigma_N^2 > 0$ where $\sigma_N^2 = \text{Var}(\sum_{i=1}^N T_{i,N})$,
 - (3) NED coefficients satisfy $\sum_{r=1}^{\infty} r^{d_0-1} \varphi(r) < \infty$,
 - (4) NED scaling factors satisfy $\sup_{N,i \in D} d_{i,N} < \infty$,
- then $\sigma_N^{-1} \sum_{i=1}^N (T_{i,N} - ET_{i,N}) \xrightarrow{d} N(0, 1)$.

Lemma C3. *Under Assumptions (W0), (W1), (W2), (W3), and (W4), and (W-REG), we have*

$$\begin{aligned} \frac{1}{N} R'_N Z_N - \frac{1}{N} E(R'_N Z_N | G_N, X_N) &= o_p(1); \quad \frac{1}{N} Z'_N Z_N - \frac{1}{N} E(Z'_N Z_N | G_N, X_N) = o_p(1); \\ \frac{1}{\sqrt{N}} Z'_N v_N &\xrightarrow{d} \mathcal{N}(0, \Sigma_c), \quad \text{where } \Sigma_c = \lim_{N \rightarrow \infty} \frac{1}{N} E(Z'_N v_N v'_N Z_N | G_N, X_N). \end{aligned}$$

Proof of Lemma C3. For terms in Z_N , we choose $H'_N X_N$ as an example, other terms are similar to show.

$$\begin{aligned}
& \frac{1}{N} \text{Var}[(H'_N X_N)' v_N | G_N, X_N] \\
&= \frac{1}{N} E \left(X'_N H_N [\varepsilon_N + \lambda(G_N - \frac{H_N}{1-p}) y_N] [\varepsilon_N + \lambda(G_N - \frac{H_N}{1-p}) y_N]' H'_N X_N | G_N, X_N \right) \\
&= \frac{\sigma_\varepsilon^2}{N} X'_N E \left([H_N + \lambda(H_N G_N - \frac{H_N^2}{1-p}) M_N] [H_N + \lambda(H_N G_N - \frac{H_N^2}{1-p}) M_N]' | G_N, X_N \right) X_N \\
&\quad + \frac{\lambda^2}{N} X'_N E \left([(H_N G_N - \frac{H_N^2}{1-p}) M_N (X_N \beta)] [(H_N G_N - \frac{H_N^2}{1-p}) M_N (X_N \beta)]' | G_N, X_N \right) X_N
\end{aligned}$$

In the special case of $\lambda = 0$, we need $\inf_N \frac{1}{N} X'_N E (H_N H'_N | G_N, X_N) X_N > 0$ to make sure Σ is non-singular. As

$$\begin{aligned}
& X'_N E (H_N H'_N | G_N, X_N) X_N \\
&= \sum_{i=1}^N \sum_{k=1}^N (1-p) G_{ik} X_{i,N} X'_{j,N} + \sum_{i=1}^N \sum_{j \neq i}^N \sum_{k=1}^N (1-p)^2 G_{ik} G_{jk} X_{i,N} X'_{j,N},
\end{aligned}$$

when $\lambda = 0$, we need $\inf_N \frac{1}{N} (1-p)^2 X'_N G_N G'_N X_N > 0$, i.e., $p \neq 1$ and $G'_N X_N$ has full rank of k , which is ensured by (W-REG) (i).

Since $|\rho| < 1$, the NED rate $\varphi(r) = |\rho|^{r/c} \rightarrow 0$ as $r/c \rightarrow \infty$. Furthermore, to establish the CLT, we need $\varphi(r)$ satisfies that

$$\sum_{r=1}^{\infty} r^{d_0-1} \varphi(r) = \sum_{r=1}^{\infty} r^{d_0-1} \rho^{r/c} < \infty.$$

When c is a fixed constant, $\rho^{r/c}$ dominates in $r^{d_0-1} \rho^{r/c}$ that

$$\lim_{r \rightarrow \infty} \frac{(r+1)^{d_0-1} \rho^{(r+1)/c}}{r^{d_0-1} \rho^{r/c}} = \lim_{r \rightarrow \infty} (1 + \frac{1}{r})^{d_0-1} \rho^{1/c} = \rho^{1/c} < 1.$$

Hence, $\sum_{r=1}^{\infty} r^{d_0-1} \rho^{r/c} < \infty$ holds. Then, all conditions in Claim 3 and 4 are satisfied, and the LLNs holds for $\frac{1}{N} R'_N Z_N$, $\frac{1}{N} Z'_N Z_N$, and $\frac{1}{N} Z'_N v_N$, and the CLT holds for $\frac{1}{\sqrt{N}} Z'_N v_N$. \square

Proof of Proposition B1. As shown in Lemma C3, $\frac{1}{N}R'_N Z_N = A_c + o_p(1)$, $\frac{1}{N}Z'_N Z_N = B_c + o_p(1)$, and $\frac{1}{\sqrt{N}}Z'_N v_N \xrightarrow{d} \mathcal{N}(0, \omega_c)$ under (W0)-(W4), and (W-REG). Hence,

$$\begin{aligned} \sqrt{N}(\widehat{\theta}_a - \theta_a) &= (A'_c B_c^{-1} A_c)^{-1} A'_c B_c^{-1} \frac{1}{\sqrt{N}} Z'_N v_N + o_p(1) \\ &\xrightarrow{d} \mathcal{N}(0, (A'_c B_c^{-1} A_c)^{-1} A'_c B_c^{-1} \omega_c B_c^{-1} A_c (A'_c B_c^{-1} A_c)^{-1}). \end{aligned}$$

□

References

- Jenish, N. and I. R. Prucha (2009). Central limit theorems and uniform laws of large numbers for arrays of random fields. *Journal of econometrics* 150(1), 86–98. pages 5, 7
- Jenish, N. and I. R. Prucha (2012). On spatial processes and asymptotic inference under near-epoch dependence. *Journal of econometrics* 170(1), 178–190. pages 5, 7, 9, 14
- Qu, X. and L.-f. Lee (2015). Estimating a spatial autoregressive model with an endogenous spatial weight matrix. *Journal of Econometrics* 184(2), 209–232. pages 5, 13
- Qu, X., L.-f. Lee, and J. Yu (2017). Qml estimation of spatial dynamic panel data models with endogenous time varying spatial weights matrices. *Journal of Econometrics* 197(2), 173–201. pages 5
- White, H. (2001). *Asymptotic theory for econometricians*. Academic press. pages 3