# Information Design for Social Learning on a Recommendation Platform

### Chen Lyu\*

This version: Jan 3, 2023

#### Abstract

A recommendation platform sequentially collects information about a new product revealed from past consumer trials and uses it to better guide later consumers. Because consumers do not internalize the value of information they bring to others, their incentive for trying out the product can be socially insufficient. Given such a challenge, I study how the platform can maximize the total consumer surplus generated on it by designing its recommendation policy. In a model with binary product quality and general trial-generated signals, I show that the optimal design features a sequence of time-specific thresholds, which vary in a U-shaped pattern over the product's life. At any time, the platform should recommend the product if its current belief about the product's quality being high is above the current threshold. My analysis also illustrates the potential usefulness of a Lagrangian duality approach for dynamic information design.

Key Words: information design, social learning, recommendation platform, Lagrangian duality

<sup>\*</sup>Peking University HSBC Business School, clyu@phbs.pku.edu.cn. I am grateful to Lones Smith, Marzena Rostek and Daniel Quint for valuable advice and comments. I also thank Emir Kamenica, Benjamin Bernard, Marek Weretka, Matteo Camboni, Nima Haghpanah, Axel Anderson, Miaojun Wang, Zhen Zhou as well as various workshop and seminar participants at CY Cergy Paris University, the University of Wisconsin-Madison, Zhejiang University and Renmin University for helpful discussions. All mistakes are my own.

### 1 Introduction

Recommendation platforms are quite popular in our daily lives. For examples, people rely on Goodreads for what to read, Netflix for what to watch, Yelp for where to eat and TripAdvisor for where to travel.<sup>[1](#page-1-0)</sup> To provide better recommendations, a common practice of these platforms is to induce a kind of social learning for new products. Namely, they collect information generated from early consumers' trials of a product (e.g., rating and reviews), and use it to better guide the later consumers. In this process, however, because individual consumers do not internalize the value of information they bring to others, their incentive for trying out the new product can be insufficient. This handicaps learning and can hinder the platforms from making better-informed recommendations.

I study how a platform facing the challenge above should design its dynamic recommendation policy, which can potentially "persuade" consumers towards more socially desirable trials for a new product. In the model, a sequence of short-lived consumers arrive over the (finite) lifetime of a product with unknown quality, which can be either high or low. Whenever a consumer consumes the product, a signal about its quality will be generated and privately observed by the platform.[2](#page-1-1) Unlike some existing studies surveyed later, I allow such consumption-generated signals to be general and non-conclusive. In each period, based on the signals previously received, the platform can guide the current consumer by providing a recommendation message. Knowing the message and her own arrival time, the consumer then makes her consumption decision in a Bayes-rational way. The platform's design problem is to find a dynamic recommendation policy, to which it can commit ex-ante, in order to maximize the total consumer surplus generated on it.[3](#page-1-2)

Notice that the platform's recommendations play a dual-role – they both decide how past information is used to guide the current consumer and decide whether new informa-tion will be generated for later use.<sup>[4](#page-1-3)</sup> Ideally, the platform should recommend the product for trial as long as this is socially desirable after taking into account the informational value, even if consumption is suboptimal for the current consumer based on the current information. With such a policy, however, the expected quality of some recommendations may be too low for the consumers to follow, which will render the design ineffective. An optimal policy therefore must choose when to recommend socially desirable but individually suboptimal consumption most efficiently, subject to the requirement that the

<span id="page-1-0"></span><sup>&</sup>lt;sup>1</sup>For some non-commercial examples, consider FDA for drug uses and Medicare Advantage Star Ratings for Medicare plan choices in the US.

<span id="page-1-1"></span><sup>&</sup>lt;sup>2</sup>The platform also receives an initial private signal about the product before any consumption, which reflects its internal research or data about previous performance of similar products.

<span id="page-1-3"></span><span id="page-1-2"></span><sup>3</sup>See Section [2.2](#page-7-0) for discussions on several assumptions of the model.

<sup>4</sup>Technically, the design problem is a decentralized bandit problem, where a principle (platform) faces a bandit process but cannot control it directly. Instead, the process is controlled by a sequence of shortlived agents (consumers). The principle privately monitors the process and can influence the agents' decisions only via providing information.

consumer in each period will be willing to follow the recommendation.

The need to convince consumers to follow recommendations induces a sequence of incentive-compatibility (abbr., IC) constraints – one for each consumer – in the dynamic design problem, which makes it a *constrained Markov decision process*.<sup>[5](#page-2-0)</sup> Solving such a problem is challenging because the standard dynamic programming technique cannot directly handle those constraints. I hence adopt a Lagrangian duality approach. It allows me to partially characterize the shadow values of those IC constraints and thereafter convert the optimization into an unconstrained one. To the best of my knowledge, this is the first paper solving a (non-degenerate) constrained Markov decision process that naturally arises from a dynamic information design problem.

I show that the optimal design features a sequence of time-specific thresholds, which generally vary in a U-shaped pattern over the product's life. At any time, the platform should recommend the product if its current belief of the product's quality being high is above the current threshold. Intuitively, this suggests that the platform should set a time-varying recommendation standard, which first goes down and then goes up as the product ages. Underlying this time-pattern is a tension between the platform's desire to create informational value for future consumers and its need to satisfy the current consumer's IC constraint. When the product is very young, the informational value is high due to a long remaining product lifespan, but consumers are "skeptical" about following recommendations because they know that even the platform has not acquired much information about the product yet. This implies a binding IC constraint and necessitates a picky censorship regarding when to recommend the product. As time passes, consumers become easier to convince as they expect that the platform may have gotten better informed by previous signals. The recommendation standard can hence be lowered.<sup>[6](#page-2-1)</sup> This continues until the standard has become sufficiently low such that trials with beliefs further below it are no longer worthwhile. Thereafter, the optimal threshold will gradually go up because the informational value of consumption dwindles as the product approaches its end of life. I note that without the consumer's IC constraints, the optimal threshold should always be increasing. The decreasing part of this U-shaped pattern thus exactly reflects how the consumer's incentive problem shapes the optimal design.

The result above implies an interesting prediction about the optimal recommendation dynamic: it can feature temporary recommendation suspensions following negative consumer feedback for young products. Specifically, following a negative feedback, the platform's belief of high quality can drop below the recommendation threshold, which suspends recommendation and learning. However, if we are in the early phase of the product's life where the threshold is declining, the threshold can fall below the belief again

<span id="page-2-0"></span><sup>&</sup>lt;sup>5</sup>See [Altman](#page-51-0) [\(1999\)](#page-51-0) for a textbook treatment to constrained Markov decision processes.

<span id="page-2-1"></span> $6$ See discussion about Figure [3](#page-16-0) in Section [4.1](#page-14-0) for a more concrete intuition behind this.

a few periods later, which restarts recommendation. In practice, it is well-documented that temporary recommendation suspensions can be used for punishing misconduct of a product supplier.[7](#page-3-0) My finding here suggests another motivation for taking such actions, which is to induce more efficient social learning about new products given the inadequacy of individual consumers' incentives.

My characterization of the optimal design also enables a couple of comparative statics. The first analysis considers how the design should be adjusted when consumption becomes more likely to yield non-neutral signals about the product's quality (e.g., due to better feedback elicitation designs). I show that the recommendation standards should be lowered uniformly over time when this happens. The second analysis incorporates random consumer arrivals. I show that when we have a thicker market where consumers arrive more frequently, the optimal policy should also become more generous in recommending the product at any age.

I note that the binary quality assumption will be relaxed in an extension in the appendix. Although a full characterization for the optimal design is not available there, my duality approach still helps to reveal certain properties of it. In particular, I show that the optimal policy features a partial-order monotone structure, which can be considered as a generalization of the threshold structure. I will discuss implications of this general result for algorithmic recommendation design following the extension.

Related literature – My study closely relates to [Kremer et al.](#page-52-0)  $(2014)$  and Che & Hörner [\(2018\)](#page-52-1), who also study the optimal recommendation design when early consumption has informational externality to later consumers. $\delta$  These papers have focused on special classes of consumption-generated signals. Specifically, the main model in [Kremer et al.](#page-52-0) [\(2014\)](#page-52-0) considers fully revealing signals, i.e., the underlying quality will be fully revealed after a single trial. This allows them to reduce the design problem into a decision problem about when to induce the first trial based on the platform's initial information.<sup>[9](#page-3-2)</sup> Che  $\&$ Hörner [\(2018\)](#page-52-1) considers a Poisson learning environment with binary quality levels in continuous time. They assume that at any time the platform has either received no news, or has received conclusive news that fully reveals the product's quality. The design problem then boils down into a deterministic control problem about the recommendation intensity following the history without any news arrival. Unlike these papers, my study accommodates general non-conclusive signals. My characterization of the optimal design is thus about whether to recommend the product in each period based on any current

<span id="page-3-1"></span><span id="page-3-0"></span><sup>7</sup>For a list of real-world examples, see Table 1 in [Liang et al.](#page-53-0) [\(2020\)](#page-53-0).

<sup>8</sup>[Lorecchio & Monte](#page-53-1) [\(2021\)](#page-53-1) also considers a setting where the designer records previous agents' feedbacks to guide later agents' decisions. However, their designer has state-independent payoff, rely on restricted communication rules, and only focuses on the long-run stationary equilibrium, which makes their paper distinct from mine.

<span id="page-3-2"></span><sup>&</sup>lt;sup>9</sup>More precisely, the initial information in their paper is about an alternative consumption option, which is always tried by the first consumer with its quality fully revealed since then.

belief of the platform, which goes beyond timing of the first trial or recommendation intensity without previous news. This in particular allows me to interpret my results as regarding the time-varying recommendation standards and necessitates the more general formulation of the problem.

An extension in [Kremer et al.](#page-52-0) [\(2014\)](#page-52-0) and a strand of subsequent algorithm-oriented research have studied environments more general than the main model of [Kremer et al.](#page-52-0) [\(2014\)](#page-52-0), which do allow for non-conclusive consumption-generated signals [\(Papanastasiou](#page-53-2) [et al.,](#page-53-2) [2018;](#page-53-2) [Mansour et al.,](#page-53-3) [2020\)](#page-53-3).<sup>[10](#page-4-0)</sup> The goal of this literature is to propose algorithms that can achieve better asymptotic performance as the number of consumers coming in sequence goes to infinity, which is often measured by the decay rate of per-consumer welfare loss compared to the full-information first-best benchmark. While such measurement reflects an important aspect of the design's performance, it ignores the early consumers' potential loss from trying a low-quality product within any finite time horizon, and can be insensitive to multiplicative changes in the total welfare loss.<sup>[11](#page-4-1)</sup> Hence, for the algorithms proposed in this literature, little is known about their finite-horizon efficiency, and little has been done to improve their non-asymptotic performance. My paper complements the literature by solving the optimal design in a stylized setting with fixed time horizon, which may serve as a performance benchmark for evaluating any algorithm and help to inspire new algorithms with a non-asymptotic focus.<sup>[12](#page-4-2)</sup>

Another growing literature also considers the optimal information provision by a platform to a sequence of short-lived agents [\(Glazer et al.,](#page-52-2) [2021;](#page-52-2) [Komiyama & Noda,](#page-52-3) [2021;](#page-52-3) Küçükgül et al., [2022\)](#page-52-4). In these papers, the agents are either endowed with or are able to acquire private signals, and a central task for the platform is to infer these private signals from the agents' decisions. These papers thus consider very different information sources of the platform and explore design concerns distinct from mine.

More generally, my paper belongs to the broad literature on information design [\(Ka](#page-52-5)[menica & Gentzkow,](#page-52-5) [2011;](#page-52-5) [Rayo & Segal,](#page-53-4) [2010\)](#page-53-4), and especially to studies on dynamic designs (e.g., [Ely,](#page-52-6) [2017;](#page-52-6) [Renault et al.,](#page-53-5) [2017;](#page-53-5) [Smolin,](#page-53-6) [2021;](#page-53-6) [Ely & Szydlowski,](#page-52-7) [2020;](#page-52-7) [Ball,](#page-51-1) [2019;](#page-51-1) [Orlov et al.,](#page-53-7) [2020;](#page-53-7) [Lorecchio,](#page-53-8) [2021\)](#page-53-8). One difference between many of the studies in this literature and mine is that I consider a designer whose private information flow is controlled by the receivers' decisions, rather than being exogenous. My analysis illus-

<span id="page-4-0"></span><sup>10</sup>Also see, for examples, [Bahar et al.](#page-51-2) [\(2015\)](#page-51-2), [Mansour et al.](#page-53-9) [\(2016\)](#page-53-9), [Chen et al.](#page-52-8) [\(2018\)](#page-52-8), [Immorlica et](#page-52-9) [al.](#page-52-9) [\(2019\)](#page-52-9) and [Bahar et al.](#page-51-3) [\(2021\)](#page-51-3) for a variety of extensions.

<span id="page-4-1"></span><sup>&</sup>lt;sup>11</sup>To see this, notice that an average loss function  $L(t)$  is considered to have the same decay rate in t as  $\alpha L(t)$  for any  $\alpha > 0$ .

<span id="page-4-2"></span><sup>12</sup>[Papanastasiou et al.](#page-53-2) [\(2018\)](#page-53-2) does investigate finite-horizon design in a particular setting, but in that setting the initial information is such that either no exploration can ever happen or consumer IC constraints are never binding, which makes the optimal design obvious. They also propose to formulate the designer's problem as a constrained Markov decision process in a more general setting, but concluded it to be computationally infeasible and did not derive analytical results from it except for giving a bound on how many belief states need to involve randomization in an optimal design.

trates how such a setting naturally leads to a constrained Markov decision process after simplification by the revelation principle of [Myerson](#page-53-10)  $(1986)$ ,<sup>[13](#page-5-0)</sup> and how the Lagrangian duality approach can be useful for solving it.<sup>[14](#page-5-1)</sup>

The paper is organized as follows. Section [2](#page-5-2) presents the main model. Section [3](#page-8-0) derives the optimal design. Section [4](#page-14-1) explores dynamic properties of the optimal design. Section [5](#page-18-0) considers comparative statics. Section [6](#page-20-0) provides additional discussions. Section [7](#page-22-0) concludes with some methodological remarks. Appendix [A](#page-23-0) considers the extension with general quality support. All proofs are provided in Appendix [B.](#page-25-0)

### <span id="page-5-2"></span>2 The Model

I first describe the model, and then discuss several underlying assumptions in Section [2.2.](#page-7-0)

#### <span id="page-5-4"></span>2.1 The Setting and the Design Problem

The model features a platform, a sequence of short-lived Bayes-rational consumers and a product. The product is launched in period 1 and will remain available for consumption over  $T < \infty$  periods. In each period  $t = 1, ..., T$ , a consumer arrives at the platform and decides whether to consume the product. I denote the consumer's decision as  $a_t \in \{0, 1\}$ with  $a_t = 1$  meaning consumption occurs. Without consuming the product, the consumer will receive her outside option, whose value is normalized to zero. If she consumes the product, the consumer's utility will be equal to  $\tilde{\theta}$ , which is a random variable taking values in  $\{\theta_L, \theta_H\}$ , with  $\theta_L < 0 < \theta_H$ .<sup>[15](#page-5-3)</sup> This  $\tilde{\theta}$  measures the underlying quality of the product, which is fixed over time but initially unknown. I assume that the platform and the consumers share a common prior  $p_0$  for  $\theta = \theta_H$ .

At the beginning of period 1, the platform receives a signal  $s_0$  about  $\tilde{\theta}$ , which reflects the platform's initial information based on, for example, its internal research or data about past performance of similar products. Subsequently, an additional signal will be generated to the platform whenever a consumer consumes the product. Let  $s_i$  denote the signal from the *i*'th consumption of the product. Conditional on  $\hat{\theta}$ , I assume that  $s_1, s_2, \ldots$  are i.i.d. and are independent from  $s_0$ .

In every period, the platform can compute its posterior belief about the product's

<span id="page-5-1"></span><span id="page-5-0"></span><sup>13</sup>I provide a more detailed remark on this in Section [7.](#page-22-0)

<sup>&</sup>lt;sup>14</sup>[Beutler & Ross](#page-51-5) [\(1985\)](#page-51-4) and Beutler & Ross [\(1986\)](#page-51-5) were the first to use a Lagrangian approach to study constrained Markov decision processes. The method is subsequently developed and applied in many mathematical and engineering papers (see, e.g., Section 1.2 of [Altman](#page-51-0) [\(1999\)](#page-51-0) for a brief survey). These studies typically only involve a few aggregate constraints corresponding to different design criteria. In contrast, my problem features one constraint for each period, which leads to a novel dynamic aspect of the problem.

<span id="page-5-3"></span><sup>&</sup>lt;sup>15</sup>As usual, one can interpret  $\tilde{\theta}$  as the consumer's expected utility from consumption given the product's true quality, and the actually realized utility can involve an ex-post idiosyncratic shock.

quality based on the previous signals received. I use  $p_t$  to denote the platform's belief about  $\tilde{\theta} = \theta_H$  at the beginning of period t. Let  $\mu_1$  denote the distribution of  $p_1 = \mathbb{P}(\tilde{\theta} =$  $\theta_H|s_0\rangle$ ; let  $G(\cdot|\cdot)$  denote the transition kernel of  $(p_t)_{t=1}^T$  following one's consumption; let  $D(\cdot|p)$  denote the Dirac measure at p. The process of  $(p_t)_{t=1}^T$  then follows the following transition rule:

<span id="page-6-3"></span><span id="page-6-2"></span>
$$
p_1 \sim \mu_1 \tag{1}
$$

$$
p_{t+1}|p_t, a_t \sim a_t G(\cdot | p_t) + (1 - a_t)D(\cdot | p_t)
$$
\n(2)

For later analyses, I define  $u(p) := p\theta_H + (1-p)\theta_L$ , and define  $\bar{p}$  to be the indifferent belief for consumers, i.e.,  $u(\bar{p}) = 0$ .

Before the realization of  $s_0$ , the platform can commit to an information transmission policy that decides what message to convey to the coming consumer in each period based on the information available at that time. I assume that the consumer can neither observe previous messages, nor observe decisions of earlier consumers.[16](#page-6-0) She only observes her arrival time and her own message, and then decides whether to consume the product.

The timeline of the environment is summarized as follows:

- 1. Before period 1, the platform (publicly) commits to an information transmission policy, and the product's quality  $\tilde{\theta}$  is secretly realized. Then, the platform privately receives its initial signal  $s_0$  about  $\hat{\theta}$ .
- 2. At the beginning of each period  $t = 1, ..., T$ , a consumer arrives and receives a message from the platform, which is generated according to the information transmission policy. She then decides whether to consume the product. If she is the  $n't$ h consumer who consumes the product, signal  $s_n$  will be generated to the platform. The economy then enters into the next period.

The Designer's Problem: I look for the information transmission policy that maximizes the total consumer surplus generated on the platform over the product's lifetime, i.e.,  $\sum_{t=1}^{T} \mathbb{E}[a_t \tilde{\theta}]$ . By the revelation principle [\(Myerson,](#page-53-10) [1986\)](#page-53-10), it suffices to consider incentive-compatible recommendation policies, which just decide whether to recommend the product for consumption in every period, subject to the requirement that a Bayesrational consumer will follow the recommendation. Since the belief  $p_t$  summarizes all the payoff-relevant information of the platform in period  $t$ , standard argument implies that we can further focus on randomized Markov policies with respect to the process of  $(p_t)_{t=1}^T$ .<sup>[17](#page-6-1)</sup> Formally, a *randomized Markov recommendation policy* is a sequence of measurable mappings  $\phi := (\phi_t : t = 1, ..., T)$ , where each  $\phi_t : [0,1] \to [0,1]$  decides the

<span id="page-6-0"></span> $16$ The model therefore mainly fits applications where the platform is using private recommendations sent to each individual consumer instead of using public recommendations.

<span id="page-6-1"></span><sup>&</sup>lt;sup>17</sup>A formal proof for this is available upon request.

probability of recommending the product at time t given any  $p_t \in [0,1]$ . For the rest of the paper, by "policy" I will be referring to a policy of this type.

I impose the following assumption on consumption-generated signals throughout:

<span id="page-7-4"></span>Assumption 1. Assume the following:

- (i)  $\mathbb{P}(\mathbb{E}[\tilde{\theta}|s_0] > 0) > 0;$
- (ii) For any  $i \geq 0$ , we have  $\mathbb{P}(s_i \in A | \tilde{\theta} = \theta_L) < \mathbb{P}(s_i \in A | \tilde{\theta} = \theta_H)$  for some (measurable) set A in the realization space of  $s_i$ ;
- (iii) For any  $i \geq 0$ , we have  $\mathbb{P}(s_i \in A | \tilde{\theta} = \theta_L) > 0 \Leftrightarrow \mathbb{P}(s_i \in A | \tilde{\theta} = \theta_H) > 0$  for any (measurable) set  $A$  in the realization space of  $s_i$ .

Condition (i) implies that it is optimal for the first consumer to consume given some realizations of the platform's initial information.[18](#page-7-1) Without such a condition, the first consumer will never want to consume the product, knowing which the second consumer will never consume either. Induction then implies that no consumption can ever happen under any design. Condition (i) rules out such a trivial scenario. Condition (ii) simply guarantees that the signals are indeed informative about  $\tilde{\theta}$ . Condition (iii) implies that no signal realization can conclusively reveal the quality level. It helps to simplify the exposition of certain proof, but is not essential for results in the paper.

#### <span id="page-7-0"></span>2.2 Discussion on Model Assumptions

1. Information v.s. monetary incentive. The model assumes that the platform cannot directly pay early consumers for trying out the new product. While it can work well in some applications, the use of monetary incentive may be problematic in others. In particular, if consumers are only attracted by the monetary incentive instead of the product itself, they may pretend to consume the product and leave some artificial feedback just to earn the money, especially when the product's pecuniary price is zero (e.g., digital contents on a subscribed platform).<sup>[19](#page-7-2)</sup> Moreover, monetary subsidy can sometimes attract a biased pool of consumers, whose feedback can be unhelpful or even misleading.<sup>[20](#page-7-3)</sup> I hence focus on the design of information instead of monetary incentives in this paper.

<span id="page-7-1"></span><sup>&</sup>lt;sup>18</sup>To guarantee this, one may instead impose the slightly weaker condition  $\mathbb{P}(\mathbb{E}[\tilde{\theta}|s_0] \geq 0) > 0$ . I impose the stronger condition for a technical reason when deriving the duality result.

<span id="page-7-2"></span><sup>&</sup>lt;sup>19</sup>For example, one may play a movie at background without watching it, and then fabricate some feedback to be paid. This kind of moral hazard problem makes it generally hard to buy information directly with monetary payment.

<span id="page-7-3"></span><sup>20</sup>For example, if we subsidize for trying a fine-dining restaurant, we may attract someone who will never eat there without the subsidy. Feedback from such consumers can be misleading as the product is actually targeting someone else.

- 2. Consumer information on product launch time. An important assumption of the model is that each consumer can observe when the product was launched. This is reasonable for many products like books, TV-series, restaurants or hotels, which typically have a public release or opening time. For some other products, however, the consumer may only have a rough idea about their ages. In such a case, my design setting can be considered as being robust to uncertainty about the consumer's exact information. Under the assumption that consumers can perfectly observe the product's launch time, the optimal design being derived will be incentive-compatible no matter what information consumers actually have about the product's age. It thus provides the best guaranteed performance.
- 3. Consumer information on earlier consumer arrivals. The model above has a feature that each consumer perfectly knows how many consumers have arrived before her, which is unrealistic. Fortunately, this is not truly a concern because my framework can be easily extended to incorporate random consumer arrivals. In such an extension, each consumer can make inference about how many consumers have arrived earlier based on her own arrival time, but cannot know the number exactly. All of my results will still hold. To ease notations, however, I do not explicitly introduce random consumer arrivals until Section [5.2,](#page-19-0) where I will examine how the arrival rate affects the optimal design.
- 4. Timing and consumer flow. Another feature of the model is that consumer arrivals are distributed evenly over time. Instead of treating this as a simplifying assumption, one can understand it in terms of how the model times  $t = 1, 2, \dots$  are defined. That is, the model time flow is arranged so that the consumers will arrive evenly over time. This implies that, for example, if consumers arrive more frequently in the first calendar month after the product's launch than in the second month, we should have denser model periods in the first month than in the second month. In this way, the model can capture product life-cycle with time-varying consumer flow flexibly.

## <span id="page-8-0"></span>3 Characterization for the Optimal Design

#### <span id="page-8-1"></span>3.1 The Constrained Markov Decision Process

Let  $\Phi$  denote the set of all (measurable) policies. Given any  $\phi \in \Phi$ , I use  $\mathbb{P}_{\phi}$  to denote the probability measure over events about  $((a_t)_{t=1}^T, (p_t)_{t=1}^T)$  provided that consumers follow the recommendations, and use  $\mathbb{E}_{\phi}$  to denote the corresponding expectation operator. Then, the incentive-compatibility (IC) constraint for a time-t consumer can be written as:[21](#page-9-0)

$$
\mathbb{P}_{\phi}(a_t = 1) > 0 \Rightarrow \mathbb{E}_{\phi}[u(p_t)|a_t = 1] \ge 0 \tag{3}
$$

<span id="page-9-4"></span>
$$
\mathbb{P}_{\phi}(a_t = 0) > 0 \Rightarrow \mathbb{E}_{\phi}[u(p_t)|a_t = 0] \le 0 \tag{4}
$$

where recall that  $u(p) := p\theta_H + (1-p)\theta_L$ . These respectively guarantee that the consumer will follow the recommendation when the product is recommended  $(a_t = 1)$  and when it is not  $(a_t = 0)$ . Since one's consumption of the product generally benefits later consumers by yielding information, the designer will never want to recommend  $a_t = 0$ when consumption is optimal for the current consumer. This implies that the second constraint above is non-restrictive for the designer and can thus be omitted. For the first constraint, we can more compactly write it as  $\mathbb{E}_{\phi}[u(p_t)|a_t=1]\mathbb{P}_{\phi}(a_t=1)\geq 0$ , which is equivalent to  $\mathbb{E}_{\phi}[a_tu(p_t)] \geq 0$ . Notice  $\mathbb{E}_{\phi}[a_tu(p_t)]$  is the expected surplus of the time-t consumer when she follows the recommendation under  $\phi$ . The designer's problem can hence be formulated  $\text{as}:^{22}$  $\text{as}:^{22}$  $\text{as}:^{22}$ 

<span id="page-9-3"></span>
$$
\max_{\phi \in \Phi} \left\{ \sum_{t=1}^{T} \mathbb{E}_{\phi}[a_t u(p_t)] \right\} \tag{5}
$$

<span id="page-9-2"></span>
$$
\text{s.t. } \mathbb{E}_{\phi}[a_t u(p_t)] \ge 0 \quad \forall t = 1, ..., T \tag{6}
$$

Due to the presence of the expectation operator, each constraint in [\(6\)](#page-9-2) is not just restricting the recommendation decision following a particular realization of  $p_t$ , but involves integration over the entire distribution of  $p_t$ . Such an aggregated IC constraint arises because the payoff-relevant process  $(p_t)_{t=1}^T$  is only observed by the platform but not by the consumer, who therefore must integrate over the equilibrium distribution of  $p_t$ when computing her posterior belief given any recommendation message. The presence of such constraints makes the problem a constrained Markov decision process, which cannot be directly handled by dynamic programming with  $p_t$  being the state variable. To overcome this difficulty, I will provide a dual characterization in Section [3.2,](#page-10-0) which allows me to partially reduce the problem into an unconstrained one.

The following lemma reveals the key properties of the belief process needed for later analyses.

<span id="page-9-5"></span>**Lemma 1.** The belief process  $(1) - (2)$  $(1) - (2)$  $(1) - (2)$  has the following properties: (P1)  $G(\cdot|p)$  as a measure-valued function of p is weakly continuous. (P2)  $\int_{p'} u(p') G(dp'|p) = u(p)$ . (P3)  $G(\cdot|p)$  increases in p in terms of first-order stochastic dominance.

<span id="page-9-1"></span><span id="page-9-0"></span><sup>21</sup>Notice  $\mathbb{E}_{\phi}[\tilde{\theta}|a_t] = \mathbb{E}_{\phi}[\mathbb{E}_{\phi}[\tilde{\theta}|p_t]|a_t] = \mathbb{E}_{\phi}[u(p_t)|a_t]$  since  $a_t$  is measurable w.r.t.  $p_t$ .

<sup>22</sup>[Papanastasiou et al.](#page-53-2) [\(2018\)](#page-53-2) first proposed the constrained Markov decision process formulation for this kind of design problem. However, they do not pursue much further analysis with it. See footnote [12](#page-4-2) for details.

(P4)  $G([0,p)|p)$  and  $G((p,1]|p)$  are strictly positive for any  $p \in (0,1)$ .  $(P5)$   $\mu_1((\bar{p}, 1]) > 0.$ 

Property (P1) means that small changes in the prior can only lead to small changes in the posterior, which is a technical result that guarantees the existence of the optimal design. Property (P2) is implied by the standard law of iterated expectation. Property (P3) is an inertia property of the belief process, which roughly says that a product looking more promising today is also more likely to look promising tomorrow. It will be important for showing the threshold structure of the optimal policy. Property (P4) is directly implied by the assumption that signals are informative, which guarantees that the belief process will not stay constant for sure following consumption. For property (P5), recall that  $\bar{p}$  is such that  $u(\bar{p}) = 0$ . The property is thus directly implied by Assumption [1\(](#page-7-4)i), which allows consumption and learning to occur with some probability in period 1. I note that these five properties are all that one needs to know about the belief process for later analyses, which abstract away from other details of the learning process.

The following result guarantees that the designer's problem is well defined.

<span id="page-10-3"></span>**Proposition 1.** There exists an optimal solution to the designer's problem  $(5) - (6)$  $(5) - (6)$  $(5) - (6)$ .

#### <span id="page-10-0"></span>3.2 The Dual Characterization

Given any vector of Lagrangian multipliers  $\lambda \in \mathbb{R}^T_+$  associated to the IC constraints, I define the Lagrangian function of the designer's problem as:

$$
\mathcal{L}(\phi; \lambda) = \sum_{t=1}^{T} \mathbb{E}_{\phi}[(1 + \lambda_t) a_t u(p_t)] \tag{7}
$$

Then, we have the following strong-duality result.

<span id="page-10-2"></span>**Lemma 2.** Let  $w^*$  denote the optimal value of the designer's problem. Then,

<span id="page-10-1"></span>
$$
w^* = \min_{\lambda \in \mathbb{R}_+^T} \sup_{\phi \in \Phi} \mathcal{L}(\phi; \lambda)
$$
\n(8)

where the minimum is achieved by some  $\lambda^*$ . Given any such  $\lambda^*$ , a policy  $\phi^*$  is optimal for the designer's problem if and only if:

- (*i*)  $\phi^* \in \arg \max_{\phi \in \Phi} \mathcal{L}(\phi; \lambda^*)$
- (*ii*)  $\lambda_t^* \mathbb{E}_{\phi^*}[a_t u(p_t)] = 0, \forall t = 1, ..., T$
- (iii)  $\mathbb{E}_{\phi^*}[a_t u(p_t)] \geq 0, \forall t = 1, ..., T$

To see how this result is helpful, notice that once we know a solution  $\lambda^*$  to the dual problem [\(8\)](#page-10-1), which intuitively measures the "shadow values" of the IC constraints,

the lemma implies that any optimal policy must solve  $\max_{\phi \in \Phi} \mathcal{L}(\phi; \lambda^*)$ , which is an unconstrained problem. The optimal design can then be characterized by studying this unconstrained problem with the standard dynamic programming approach.

The problem here, however, is that the value of  $\lambda^*$  is not available. Generally, deriving it requires one to either solve the min-max problem in [\(8\)](#page-10-1) or solve the fixed-point problem defined by conditions (i) – (iii) jointly for  $(\lambda^*, \phi^*)$ , both of which are difficult. Fortunately, as I show below, a property of  $\lambda^*$  can be directly derived from the dual problem, which turns out to enable a sharp characterization for the optimal policy.

#### <span id="page-11-2"></span>3.3 Main Structures of the Optimal Design

The following lemma is a key result derived from the dual problem [\(8\)](#page-10-1).

<span id="page-11-0"></span>**Lemma 3.** There exists  $\lambda^* \in \arg\min_{\lambda \in \mathbb{R}_+^T} \sup_{\phi \in \Phi} \mathcal{L}(\phi; \lambda)$  such that  $\lambda_t^* \geq \lambda_{t+1}^*$  for all  $t = 1, ..., T - 1.$ 

To see an intuition behind this result, assume that the dual problem has a unique solution  $\lambda^*$ . As usual, we can interpret  $\lambda_t^*$  as the shadow value of marginally relaxing the time-t IC constraint for the designer's problem. As time passes, two changes happen in the designer's problem. First, as information accumulates over time, we will be able to make recommendation selections more wisely. This makes it possible to obey the consumer's IC constraint with less sacrifice of socially desirable consumption. Second, as the remaining lifetime of the product gets shorter, the dynamic value from having additional myopically suboptimal consumption drops. These both suggest that relaxing later IC constraints is less helpful than relaxing the earlier ones. Hence, the associated shadow values should decrease over time. While this argument is intuitive, formalizing it can be difficult. I hence instead develop an "inter-change" argument for the lemma's proof. In particular, given any  $\lambda^*$  solving the dual problem, I show that if two adjacent components of it violate the time pattern, then interchanging them will lead to a new solution to the dual problem. Starting with any solution to the dual problem, one can thus construct a solution satisfying the time pattern by making such interchanges repeatedly.

An important implication of Lemma [3](#page-11-0) is that the optimal design will generally feature a two-phase structure. In the first phase,  $\lambda_t^* > 0$  and the IC constraints are thus binding; in the second phase,  $\lambda_t^* = 0$  and the IC constraints are thus essentially non-restrictive.<sup>[23](#page-11-1)</sup> As we will see later, this two-phase structure is critical in constructing the optimal design and has important implications on the dynamic pattern of optimal recommendations.

The time-pattern of  $\lambda^*$  found in Lemma [3](#page-11-0) also turns out to induce a simple solution structure for the Lagrangian function optimization  $\max_{\phi \in \Phi} \mathcal{L}(\phi; \lambda^*)$ . To state this result,

<span id="page-11-1"></span> $^{23}$ It is easy to see that the second phase includes at least the last period, since the optimal policy there will be myopically optimal. The first phase is non-empty as long as the prior on  $\hat{\theta}$  is not sufficiently favorable to support first-best learning.

I define threshold policies as follows:

**Definition 1.** A time-t policy  $\phi_t : [0,1] \to [0,1]$  is called a threshold time-t policy if there exists a threshold  $\eta_t \in [0, 1]$  such that  $p > \eta_t \Rightarrow \phi_t(p) = 1$  and  $p < \eta_t \Rightarrow \phi_t(p) = 0$ . A policy  $\phi$  is called a *threshold policy* if  $\phi_t$  is a threshold time-t policy for every t.

Namely, a threshold policy will recommend the product when the current belief of  $\tilde{\theta} = \theta_H$  is above a threshold, and will not recommend it when the belief is below the threshold. It can also involve arbitrary randomization at the threshold. By applying backward induction on the dynamic programming of  $\max_{\phi \in \Phi} \mathcal{L}(\phi; \lambda)$ , I show the following result:

<span id="page-12-2"></span>**Lemma 4.** Given any non-increasing sequence of multipliers  $(\lambda_t)_{t=1}^T$ , any solution to  $\max_{\phi \in \Phi} \mathcal{L}(\phi; \lambda)$  is almost surely equivalent to a threshold policy. Moreover,  $p_t > \bar{p} \Rightarrow$  $a_t = 1$  a.s. under such a policy.<sup>[24](#page-12-0)</sup>

Together with Lemma [3](#page-11-0) and the dual characterization for the optimal design, this directly implies the threshold structure of the optimal design:

<span id="page-12-3"></span>Proposition 2. Any optimal policy is almost surely equivalent to a threshold policy. Moreover,  $p_t > \bar{p} \Rightarrow a_t = 1$  a.s. under it for any  $t^{0.25}$  $t^{0.25}$  $t^{0.25}$ .

I note that although threshold policies are practically appealing, their optimality is not obvious in my setting. While the myopic value of consumption always increases in  $p_t$ , the dynamic informational value of it does not. Given the presence of IC constraints, even measuring such dynamic value is not straightforward, as one not only needs to consider the direct benefit to later consumers, but also needs to consider how better information may help to relax the IC constraints of later consumers and thereby facilitate more information generation from them. The duality approach I take partly resolves such difficulty by characterizing the shadow values of those IC constraints. Given the monotonicity property of  $(\lambda_t^*)_{t=1}^T$ , I show that when  $p_t$  is increased, the positive change in the myopic value of consumption always dominates the potentially indeterminate change in the dynamic informational value measured in the continuation problem of  $\max_{\phi \in \Phi} \mathcal{L}(\phi; \lambda^*)$ . The total value of consumption is thus always increasing in  $p_t$ , which implies the threshold structure of the optimal design.

Lemma [4](#page-12-2) above also helps to characterize the dictator's optimal policy, where by "dictator" I mean a social planner who can dictate consumers' decisions without obeying their IC constraints. Notice that if  $\lambda_t = 0$  for all t, the Lagrangian optimization max<sub> $\phi \in \Phi$ </sub>  $\mathcal{L}(\phi; \lambda)$ 

<span id="page-12-0"></span><sup>&</sup>lt;sup>24</sup>By saying  $A \Rightarrow B$  almost surely (a.s.), I mean that the event in which A happens but B does not happen is of zero probability.

<span id="page-12-1"></span> $25$ The second statement implies that the product is always recommended when consumption is optimal for the current consumer. Hence the IC constraint [\(4\)](#page-9-4) I ignored before is indeed satisfied.

is reduced to the dictator's problem. Lemma [4](#page-12-2) then implies that the dictator's problem also features a threshold solution. This solution will be used in the construction of the optimal design below.

#### <span id="page-13-1"></span>3.4 The Optimal Policy

Notation: For any vector indexed by time, I will use subscription " $> t$ " to indicate the sub-vector corresponding to time no earlier than t. For example,  $\phi_{\geq t}$  will denote the continuation policy since time t. Notations like  $\phi_{\geq t}$  and  $\phi_{\leq t}$  are similarly defined.

Based on the two-phase structure of the optimal design implied by Lemma [3](#page-11-0) and the threshold structure stated in Proposition [2,](#page-12-3) one can explicitly construct the optimal design using a forward induction algorithm. To do so, I define  $\phi^d$  to be the "most conservative" optimal policy for the dictator's problem (i.e., the designer's problem without IC constraints), whose details are provided in Appendix [B.5.](#page-38-0) When the dictator's problem admits multiple solutions,  $\phi^d$  is the most conservative one in the sense that it always breaks ties in favor of non-recommendation.

A candidate optimal threshold policy  $\phi^o$ , together with a cutoff time point  $\hat{t}$ , can be inductively defined as follows.

<span id="page-13-2"></span>**Definition 2.** A policy  $\phi^o$ , a sequence of distributions  $(\mu_t^o)_{t=1}^T$  over [0, 1] and a time point  $\hat{t} \in \{1, ..., T\}$  are defined with the following algorithm:

Start with  $t = 1$  and define  $\mu_1^o = \mu_1$ .

• Step 1: If  $\int_p [\phi_t^d(p)u(p)] \mu_t^o(dp) \geq 0$ , then end the algorithm while defining:

$$
-t=t;
$$

 $-\phi_{\geq t}^o = \phi_{\geq t}^d;$ 

- (μ<sup>o</sup><sub>s</sub>)<sub>s>t</sub> to be the marginal distributions of (p<sub>s</sub>)<sub>s>t</sub> under φ<sup>d</sup><sub>2</sub>t given p<sub>t</sub> ~ μ<sup>o</sup><sub>i</sub>. Otherwise, go to the next step.

- Step 2: Define  $\phi_t^o$  to be a threshold time-t policy such that:<sup>[26](#page-13-0)</sup>
	- (i)  $\phi_t^o(p) = 1$  for all  $p > \bar{p}$ ;
	- (ii)  $\int_p [\phi_t^o(p)u(p)] \mu_t^o(dp) = 0.$

Also define  $\mu_{t+1}^o$  to be the distribution of  $p_{t+1}$  under  $\phi_t^o$  given  $p_t \sim \mu_t^o$ . Then go back to step 1 with t replaced by  $t + 1$ .

Intuitively, beginning with the initial period, the algorithm will progressively check whether it is incentive-compatible to follow the dictator's policy in every period  $t$  given  $\phi_{\leq t}^o$  previously determined, i.e., whether  $\int_p [\phi_t^d(p)u(p)]\mu_t^o(dp) \geq 0$ . If so, the current period will be marked as period  $\hat{t}$ , and  $\phi^o$  will be set to follow  $\phi^d$  for all later periods; otherwise,  $\phi_t^o$  will be defined to have its threshold just high enough to satisfy the current consumer's

<span id="page-13-0"></span><sup>&</sup>lt;sup>26</sup>See Appendix [B.5.2](#page-38-1) for details of the construction of  $\phi_t^o$ .

<span id="page-14-2"></span>

Figure 1: Two-Phase Structure of the Optimal Policy  $\phi^o$ .

IC constraint, which is what conditions (i) and (ii) in step 2 guarantee. The policy  $\phi^o$  such defined will be in accordance with the two-phase structure of the optimal design found in Section [3.3,](#page-11-2) which is illustrated in Figure [1.](#page-14-2) In its early phase, the consumers' IC constraints will be binding and  $\phi^o$  will be just "picky" enough to satisfy those constraints as equalities. In the later phase, the IC constraints will become non-restrictive and  $\phi^c$ will follow the dictator's unconstrained optimal policy. The cutoff time  $\hat{t}$  is when  $\phi^o$  shifts from the first phase to the second, at which it starts following  $\phi^d$ .

The following proposition shows that  $\phi^o$  is indeed an optimal policy and fully characterizes any optimal design.

<span id="page-14-3"></span>**Proposition 3.** Any policy  $\phi^*$  is optimal for the designer's problem [\(5\)](#page-9-3) – [\(6\)](#page-9-2) if and only if: (i)  $\phi^*$  $\overrightarrow{a}$  agrees with  $\phi^o$  $\int_{\hat{\epsilon}}^{\hat{\epsilon}} e^{-\hat{\epsilon}}$  almost surely; (ii) given  $p_{\hat{t}} \sim \mu_{\hat{t}}^{\hat{\epsilon}}$  $\hat{t}$ ,  $\phi^*_{\hat{t}}$  $\sum_{i=1}^{\infty}$  is incentive-compatible and is optimal for the dictator's continuation problem starting from time  $\hat{t}$ . In particular,  $\phi^o$  is optimal.

The characterization in Proposition [3](#page-14-3) makes it convenient to explore dynamic features of the optimal design and study how it should be tailored to market details. I pursue these in the following sections.

### <span id="page-14-1"></span>4 Dynamic Properties of the Optimal Design

#### <span id="page-14-0"></span>4.1 Time Pattern of the Recommendation Standards

Threshold policies can be naturally interpreted as policies setting the minimum agespecific standards for a product to qualify for recommendations. Given the dynamic nature of the problem, it is conceivable that such a minimum standard should evolve over the product's life. I explore this time pattern below.

For the ease of exposition, I impose the following full-support and atomless assumption on the belief process.

<span id="page-14-4"></span>**Assumption 2.** The signals  $(s_0 \text{ and } \{s_i\}_{i\geq 1})$  are such that the marginal distributions of  $p_1, ..., p_T$  are atomless and have full support over [0, 1] under any policy.

<span id="page-15-1"></span>

Figure 2: The optimal threshold policy for a numerical example:  $T = 25$ ,  $\theta_L = -1$ ,  $\theta_H = 1$ ,  $p_0 = 0.2$ ,  $s_i | \theta \sim Normal(\theta, \sigma^2)$  with  $\sigma = 3$  for all  $i \ge 0$ . The cutoff time  $\hat{t} = 11$ .

The atomless assumption renders randomization at the threshold irrelevant, so we can solely focus on the thresholds themselves. The full-support assumption guarantees that any deviation in the recommendation threshold matters, which avoids the need to discuss off-path indeterminacy of the optimal policy. Assumption [2](#page-14-4) holds, in particular, if the log-likelihood ratios of the signals  $s_0, s_1, \ldots$  are continuous random variables with full support over  $\mathbb{R}^{27}$  $\mathbb{R}^{27}$  $\mathbb{R}^{27}$ 

Recall that  $\hat{t}$  is the first time when it is incentive-compatible for  $\phi^o$  to follow the dictator's optimal policy. The following result characterizes the time pattern of the optimal recommendation standards.

<span id="page-15-2"></span>**Proposition 4.** Under Assumption [2,](#page-14-4) the thresholds  $(\eta_t^*)_{t=1}^T$  of any optimal threshold policy satisfies: (a)  $\eta_{t-1}^* > \eta_t^*$  for all  $t \leq \hat{t} - 1$ ; (b)  $\eta_t^* < \eta_{t+1}^*$  for all  $t \geq \hat{t}$ . Moreover,  $\eta_t^* \leq \bar{p}$  for any t.

The proposition suggests that the optimal recommendation standard should first decrease and then increase over the product's life, which corresponds to the two phases with binding and non-binding IC constraints respectively. Figure [2](#page-15-1) presents a numerical example. Underlying this result is a tension between our desire to create dynamic informational value for future consumers and the need to fulfill the current consumer's IC constraint. In the early phase, the dynamic value is generally high since the product has a long future to go, but consumers are more "skeptical" about following recommendations because they know that even the platform has not acquired much information yet. This implies a binding IC constraint and necessitates a picky censorship over the beliefs eligible for recommendations. As time proceeds, consumers become easier to convince and the recommendation criterion can thus be relaxed. This continues until the

<span id="page-15-0"></span><sup>&</sup>lt;sup>27</sup>That is, for both  $i = 0$  and  $i \ge 1$ ,  $s_i$  admits density functions  $f_i^L$  and  $f_i^H$  conditional on  $\tilde{\theta} = \theta_L$  and  $\tilde{\theta} = \theta_H$  respectively such that  $\log \left( \frac{f_i^H(s_i)}{f^L(s_i)} \right)$  $f_i^{(t)}(s_i)$  is a continuous random variable with full support over  $\mathbb{R}$ .



<span id="page-16-0"></span>Figure 3: Explanation for decreasing recommendation thresholds before time  $\hat{t}$ . The dots represent possible realizations of  $p_t$  or  $p_{t+1}$ . The arrows indicate the evolution of these beliefs. The blue bars represent the optimal recommendation thresholds.

standard is already sufficiently low such that trials with beliefs further below it are no longer worthwhile given the remaining time of the product. The IC constraint then turns non-restrictive. Thereafter, the optimal design just follows the dictator's optimal policy, where the recommendation standard gradually goes up since the dynamic value from myopically suboptimal consumption shrinks as the product approaches its end of life.

Figure [3](#page-16-0) explains why the optimal threshold drops before time  $\hat{t}$  more precisely. The two dots on the left represent two possible realizations of  $p_t$ , and the blue bar between them represents the optimal threshold in period t. Since the green dot is above the threshold, it is associated with a recommendation and will hence split in a mean-preserving spread manner, which leads to realizations of  $p_{t+1}$  represented by the two new green dots in period  $t + 1$ . This reflects the new information generated by consumption. The black-dot belief in period  $t$  does not qualify for a recommendation and is thus carried over into period  $t + 1$ . Now, suppose the designer keeps the threshold unchanged over the two periods. Then in period  $t + 1$ , only the upper green dot (green dot 2) would qualify for a recommendation. Since this belief is more favorable than its predecessor (green dot 1), the consumer's IC constraint in period  $t + 1$  would turn slack. Intuitively, the better information in period  $t + 1$  would have induced a more favorable selection for the consumer if the threshold were kept the same as before. This leaves room for the designer to also include the black dot into the recommendation region. When  $t + 1 < \hat{t}$ , the designer indeed wants to do so since my previous characterization has shown that the consumer's IC constraint should keep binding before time  $\hat{t}$ . This implies that the threshold in period  $t + 1$  should be lowered.

The U-shaped pattern of recommendation thresholds has interesting implications for the optimal recommendation and learning dynamics, which I examine in the next subsection.

<span id="page-17-0"></span>

Figure 4: A realized path of recommendations under the optimal design in the numerical example of Figure [2.](#page-15-1) Blue bars represent the optimal thresholds. The crosses and circles track the platform's realized beliefs  $(p_t)_{t=1}^T$ , where green circles mean that recommendation is made and red crosses mean the opposite.

#### <span id="page-17-2"></span>4.2 Optimal Recommendation Dynamic

Figure [4](#page-17-0) demonstrates an example path of realized recommendations under the optimal policy in the numerical example of Figure  $2^{28}$  $2^{28}$  $2^{28}$  The sequence of blue bars still represents the optimal age-specific thresholds. The series of crosses and circles tracks a realized path of  $(p_t)_{t=1}^T$ , where green circles mean that the current belief is above the current threshold and the product is hence recommended, while red crosses mean the opposite.

The figure highlights an interesting property of the optimal recommendation dynamic – recommendation and learning can be temporarily suspended following negative feedback from the last consumption (e.g., the suspension in period 4). Such kind of suspension is beneficial because it allows us to support myopically suboptimal exploration for products looking more promising at the same age (i.e., with a  $p_t$  being higher but still less than  $\bar{p}$ ) without violating the IC constraint. However, the suspension does not need to last forever when further exploration is still socially desirable. As soon as the threshold for recommendation drops below the current belief, trials for the product should be resumed. Notice that such a restart of recommendations can only happen in the early phase of the product's life, where the IC constraints are binding and the recommendation threshold decreases over time. If it were in the later phase, any suspension of recommendation would be permanent.

In practice, temporary recommendation suspensions (or deprioritization) following negative consumer feedback are often used for punishing misconduct of a product sup-

<span id="page-17-1"></span><sup>&</sup>lt;sup>28</sup>The figure presents a case where recommendation is eventually abandoned, which is more likely to happen when the product's true quality is low. However, the property discussed below does not rely on this.

plier (e.g., a seller or content provider).<sup>[29](#page-18-1)</sup> My finding here suggests another motivation for taking such actions, which is to induce social exploration on new products in a more efficient way given the inadequacy of individual consumer's incentive. Compared to those used for punishing supplier misconduct, the temporary recommendation suspensions in my model have two distinct features. First, they can happen following negative feedback regarding the product's innate features instead of unsatisfactory behaviors of the supplier. Second, they only happen for young products. Presuming that platforms in practice are indeed trying to enhance social learning about new products by tailoring their recommender systems, these may serve as concrete predictions of the model that can be tested with real data on platform recommendations.

### <span id="page-18-0"></span>5 Comparative Statics

In this section, I provide two comparative statics regarding the optimal recommendation standards.

#### 5.1 Information Generation Rate

In many applications, having someone trying out a product is not guaranteed to generate meaningful information about the product's quality. The consumer may not leave feedback, $30$  or may give a piece of feedback that is too cursory to be authenticated.<sup>[31](#page-18-3)</sup> These will lead to little post-consumption information generation.

How should the optimal design be adjusted when consumption becomes more likely to yield information due to, for example, better feedback elicitation designs? To study this, I introduce an information generation rate into my model. Specifically, I assume that following one's consumption, the signal  $s_i$  being generated will be a compounded signal. It has probability  $\alpha \in (0, 1)$  to be an informative signal and has probability  $1 - \alpha$  to be uninformative, and the platform can tell which type the signal is.<sup>[32](#page-18-4)</sup> Let  $G^I(\cdot|\cdot)$  denote the transition kernel for the platform's belief following an informative signal. Given any  $\alpha$ , the transition kernel for  $p_t$  following one's consumption then becomes:

<span id="page-18-5"></span>
$$
G(\cdot|p) = \alpha G^{I}(\cdot|p) + (1 - \alpha)D(\cdot|p)
$$
\n(9)

<span id="page-18-2"></span><span id="page-18-1"></span><sup>&</sup>lt;sup>29</sup>See, e.g., Table 1 in [Liang et al.](#page-53-0)  $(2020)$  for a list of examples.

<sup>30</sup>I note that non-feedback is not necessarily uninformative. However, it will be so if there is little self-selection bias so that the feedback probability is uncorrelated with the product's quality, which may be reasonable to assume for some applications.

<span id="page-18-3"></span><sup>&</sup>lt;sup>31</sup>In particular, platforms like Amazon or Yelp often rely on textual analysis to filter out fake reviews. If a review is not material enough to pass such a test, it will be disregarded or attached with little weight in any recommendation algorithm. This is important to deter fake reviews.

<span id="page-18-4"></span> $32$ This compounded signal satisfies requirements in Assumption [1](#page-7-4) as long as the informative component is non-conclusive and indeed informative.

The following proposition provides the comparative statics result with respect to  $\alpha$ . For simplicity, I still impose the full-support and atomless assumption in Section [4.1.](#page-14-0)

<span id="page-19-2"></span>**Proposition 5.** Assume Assumption [2](#page-14-4) holds.<sup>[33](#page-19-1)</sup> Given any  $\alpha$ , let  $(\eta_t^*(\alpha))_{t=1}^T$  denote the thresholds of the optimal threshold policy. Then  $\eta_t^*(\alpha)$  weakly decreases in  $\alpha$  for all t.

The proposition suggests that if the information generation rate is improved, the optimal recommendation standard should be lowered for all product ages. There are two forces behind this change. First, a higher  $\alpha$  implies greater informational value from one's consumption, which motivates more exploration. Second, with a higher  $\alpha$ , information from past consumption is accumulated at a faster rate. This enables better-informed recommendations at any time, which makes consumers more willing to follow the recommendations ceteris paribus. We thus have room to lower the recommendation standards without violating the IC constraints. Together, these lead to looser recommendation criteria in the optimal design.

The formal proof of the proposition is technically involved because it requires comparing two controlled Markov processes corresponding to different  $\alpha$ . Central to the proof is a coupling argument, where I explicitly construct the belief processes under the optimal designs corresponding to different  $\alpha$  in the same probability space. This allows a direct comparison between them. I refer interested readers to Claim (d) in Appendix [B.7.](#page-44-0)

#### <span id="page-19-0"></span>5.2 Random Consumer Arrivals

As is mentioned earlier, my framework easily accommodates random consumer arrivals. I formally illustrate this below and examine how the optimal design should depend on the consumer arrival rate.

I now assume that a consumer arrives in each period with probability  $\rho \in (0,1)$ . The arrivals are independent over time and independent from other random objects in the model. Accordingly, I re-interpret  $a_t$  as the consumption decision (or the platform's recommendation) of an arriving consumer. Then, a signal of quality will be generated following period t if and only if  $a_t = 1$  and a consumer does arrive in that period. The transition rule of the belief process  $(p_t)_{t=1}^T$  then becomes:

$$
p_{t+1}|p_t, a_t \sim a_t [\rho G(\cdot | p_t) + (1 - \rho)D(\cdot | p_t)] + (1 - a_t)D(\cdot | p_t)
$$
\n(10)

Compared to the transition rule in [\(2\)](#page-6-3), the change is that  $G(\cdot|\cdot)$  is now replaced by  $\rho G(\cdot|\cdot) + (1 - \rho)D(\cdot|\cdot)$ . This will be the only change in the design environment.

Another change to the designer's problem is that we should now replace  $u(\cdot)$  in the designer's objective function [\(5\)](#page-9-3) with  $\rho u(\cdot)$ , which reflects the fact that a consumer arrives

<span id="page-19-1"></span><sup>&</sup>lt;sup>33</sup>This holds, in particular, if the log-likelihood ratios of  $s_0$  and subsequent *informative* signals are continuous random variables, and the log-likelihood ratio of  $s_0$  has full support over  $\mathbb R$ .

only with probability  $\rho$ . This does not matter for the optimization, however, since it only multiplies the objective function with a strictly positive scalar.

Because Lemma [1](#page-9-5) still holds with  $G(\cdot|\cdot)$  replaced by  $\rho G(\cdot|\cdot) + (1 - \rho)D(\cdot|\cdot)$ , all of my previous characterizations for the optimal design will remain valid. The following proposition reveals how the consumer arrival frequency matters for the optimal design.

<span id="page-20-2"></span>**Proposition 6.** Assume Assumption [2](#page-14-4) holds.<sup>[34](#page-20-1)</sup> Given any arrival rate  $\rho$ , let  $(\eta_t^*(\rho))_{t=1}^T$ denote the thresholds of the optimal threshold policy. Then  $\eta_t^*(\rho)$  weakly decreases in  $\rho$ for all t.

The proposition suggests that when we have a thicker market where consumers arrive more frequently, the recommendation standard for any product age should be lower. The intuition is again twofold. First, a higher arrival rate implies that more consumers are likely to come in the future. This increases the informational value from early consumption. Second, with a higher arrival rate, we will in expectation have more consumption and hence more signals accumulated before any given period. This supports more generous recommendations while obeying the IC constraints.

### <span id="page-20-0"></span>6 Additional Discussion

#### 6.1 Platform's incentive to maximize total consumer surplus

The designer's objective in my model is to maximize the total consumer surplus. One may wonder whether this is in line with a commercial platform's interest. I believe so in at least two scenarios. In the first one, the platform can directly extract consumer surplus by charging a subscription fee (e.g., Netflix). In the second one, the platform does not directly charge consumers, but wants to maximize the user base attracted by its recommendation service. For example, ads-financed search engines want to maximize the quality of their organic recommendations so that more people will use them and see their sponsored ads. In both scenarios, the platform's profit will be increasing in the total consumer surplus generated on it, maximizing which should thus be a primary concern in their recommendation algorithm design.

#### 6.2 Non-binary Quality Levels

One simplifying assumption in my main model is that the product's quality can only take binary values. As in many other papers on information design or social learning,

<span id="page-20-1"></span> $34$ This still holds, in particular, under the conditions in footnote [27,](#page-15-0) but now there is no need to require  $\log\left(\frac{f_i^H(s_i)}{f^L(s_i)}\right)$  $\frac{f_i^H(s_i)}{f_i^L(s_i)}$  ( $i \geq 1$ ) to have full support over  $\mathbb{R}$ .

this allows one to represent the evolving belief with a single-dimensional variable, which significantly eases the analysis.<sup>[35](#page-21-0)</sup>

In Appendix [A,](#page-23-0) I extend the model to allow general quality support. Although a full characterization of the optimal design is not available, the duality approach does help to extend certain structures of the optimal design to that general setting. In particular, I show that the optimal design still features a two-phase structure implied by Lemma [3.](#page-11-0) Moreover, the optimal policy should be more inclined to recommend the product when the platform's current belief about quality is higher in the likelihood-ratio order. This extends the threshold structure to the case with multi-dimensional beliefs. I will discuss how this result can be helpful for algorithmic recommendation design in the appendix.

#### 6.3 Comparison to Previous Studies

As has been mentioned in the introduction, my study is closely related to [Kremer et al.](#page-52-0)  $(2014)$  and Che & Hörner [\(2018\)](#page-52-1), who also study how platform recommendations can improve social learning efficiency when early consumption produces information for later consumers through the platform. Similar to my paper, they also reveal how past information accrued to the platform enables it to persuade more consumers into socially desirable explorations in the future. However, the three papers provide different characterizations for the optimal design.

Because [Kremer et al.](#page-52-0) [\(2014\)](#page-52-0) assumes fully revealing consumption-generated signals, their design is mainly about when to induce the first trial of the product. $36$  The main result is that a product that looks better based on the platform's initial information (in its quality relative to an alternative option) should receive the first trial earlier. Che  $\&$ Hörner  $(2018)$  assumes that the platform learns from conclusive news that fully reveals the product's quality upon its arrival. The design in their paper is hence about "how much" to recommend the product following the history without news arrival. They show that myopically suboptimal recommendation, given no news arrival, should gain increasing intensity as the product ages until being ceased at some point. In contrast, my study accommodates general non-conclusive consumption-generated signals. My prediction of the optimal design is therefore about whether to recommend the product in each period based on any current belief of the platform. This allows me to interpret my results as being about the time-varying recommendation standards. In particular, I have found that the optimal design features threshold policies with respect to the evolving belief, and the recommendation standard should vary in a U-shaped pattern as the product ages.<sup>[37](#page-21-2)</sup>

Allowing non-conclusive signals also enables richer predictions about the optimal rec-

<span id="page-21-0"></span><sup>&</sup>lt;sup>35</sup>See, e.g., section 2 in Hörner & Skrzypacz [\(2017\)](#page-52-10) for papers on social experimentation.

<span id="page-21-1"></span><sup>36</sup>After the first trial, quality is fully revealed and recommendations should be myopically optimal.

<span id="page-21-2"></span> $37$ With binary product quality, my result nests that in [Kremer et al.](#page-52-0) [\(2014\)](#page-52-0) as a special case.

ommendation dynamic. In particular, the phenomenon of temporary recommendation suspensions following negative consumer feedback in Section [4.2](#page-17-2) cannot exist with conclusive signals, since conclusive negative feedback should stop recommendation forever. Moreover, both of the previous papers suggest that exploration (i.e., myopically suboptimal trials) should stop after some middle age of the product. With general consumptiongenerated signals, however, Proposition [4](#page-15-2) implies that exploration can happen until the last period of the product's life, although the belief region for exploration gradually shrinks after some point.

Finally, my study has also provided a couple of comparative statics regarding the optimal recommendation standards in Section [5,](#page-18-0) which do not have counterparts in the previous papers.

## <span id="page-22-0"></span>7 Conclusion and Methodological Remarks

I have studied the optimal design of platform recommendations when early consumption of a product yields informational externality to later consumers. The optimal design is shown to feature threshold policies with the recommendation standard varying in a U-shaped pattern over the product's life. An interesting implication is that recommendations can be temporarily suspended following negative consumer feedback for young products, while any such suspension must be permanent for older products. My characterization also enables a couple of comparative statics. In particular, I have shown that the recommendation standard should be lower throughout the product's life when consumption is more likely to generate informative feedback or when consumers are arriving more frequently over time.

My model accommodates non-conclusive consumption-generated signals. Consequently, compared to the existing literature, it requires a more general formulation of the designer's problem as a constrained Markov decision process. I argue that such a mathematical formulation can naturally arise in dynamic information design problems where the designer's private information flow is controlled by the receivers' decisions. In such scenarios, if one focuses on direct mechanisms, then the design problem can be treated as a Markov decision process where the designer decides the receivers' actions and thereby controls his own information flow subject to the receivers' IC constraints. Since the receivers do not observe the designer's information, their IC constraints will involve taking expectations over it. This then leads to the aggregated constraints that cannot be directly handled in dynamic programming (like constraint  $(6)$ ). I expect that such a formulation and the Lagrangian duality approach I take can be generally useful for this kind of dynamic information design problems.

### <span id="page-23-0"></span>A Non-binary Quality Levels

In this appendix, I extend the model to allow for non-binary product quality and generalize certain characterizations of the optimal design.

#### A.1 The General Setting

Consider the same setting as in Section [2.1](#page-5-4) except that the support of  $\tilde{\theta}$  can be an arbitrary set  $\Theta \subset \mathbb{R}$  now. I assume that the joint distribution of  $\tilde{\theta}$  and the signals  $(s_0, s_1, \ldots)$  is such that the platform's posterior belief is always within a family of distributions  $\{Q_z\}_{z\in\mathbb{Z}}$ , where  $Z \subset \mathbb{R}^n$  is a countable parameter set.<sup>[38](#page-23-1)</sup> I assume that  $\{Q_z\}_{z \in Z}$  admits density functions  $\{q_z\}_{z\in\mathbb{Z}}$  with respect to some common dominating measure over  $\mathbb{R}$ , and  $q_z(\cdot) > 0$ on  $\Theta$  for all  $z \in \mathbb{Z}$ . Moreover, I impose the following assumption:

<span id="page-23-2"></span>Assumption A.1. Assume the following:

(i)  $\mathbb{P}(\mathbb{E}[\tilde{\theta}|s_0] > 0) > 0.$ 

(ii) For any  $i \geq 1$ ,  $s_i$  takes values in some set  $S \subset \mathbb{R}$ . Its distribution conditional on  $\theta$  admits a conditional density function  $\ell(\cdot|\cdot)$  (w.r.t. some dominating measure over  $\mathbb{R}$ ) such that:  $\ell(s|\theta) > 0$  for all  $s \in S$  and  $\theta \in \Theta$ ;  $\ell(\cdot|\theta)$  increases in  $\theta$  in the likelihood-ratio order.

Condition (i) plays the same role as its counterpart in Assumption [1.](#page-7-4) Condition (ii) implies that higher realizations of  $s_i$  suggest that the product is more likely to be of higher quality. This framework is general enough to incorporate many parametric learning models with congruent prior and signals (e.g., the Beta-Binomial model). Moreover, it accommodates any learning model with finite support of  $\tilde{\theta}$  that satisfies Assumption [A.1.](#page-23-2)

Let  $z_t \in Z$  denote the platform's belief parameter at the beginning of period t. Since  $z_t$  (or  $Q_{z_t}$ ) summarizes all information available to the platform at time t, we can focus on (randomized) Markov recommendation policies w.r.t.  $(z_t)_{t=1}^T$ . Formally, any policy of this type is a sequence of measurable mappings  $\phi := (\phi_t : t = 1, ..., T)$ , where each  $\phi_t: Z \to [0, 1]$  decides the probability of recommending the product at time t given any belief parameter  $z_t \in Z$ .

As in Section [5.2,](#page-19-0) I also allow i.i.d. random consumer arrivals and use  $\rho$  to denote the arrival probability.

<span id="page-23-1"></span> $38I$  assume Z to be countable for simplicity, which avoids the need to discuss certain measurability issues about the optimal policy.

#### A.2 Characterizations

Following similar arguments as those in Section [3.1](#page-8-1) and Section [5.2,](#page-19-0) the designer's problem can be formulated as follows:

$$
\max_{\phi \in \Phi} \left\{ \sum_{t=1}^{T} \mathbb{E}_{\phi}[a_t u(z_t)] \right\}
$$
\ns.t. 
$$
\mathbb{E}_{\phi}[a_t u(z_t)] \ge 0 \quad \forall t = 1, ..., T
$$
\n
$$
z_{t+1} | z_t, a_t \sim a_t [\rho G(\cdot; z_t) + (1 - \rho) D(\cdot; z_t)] + (1 - a_t) D(\cdot; z_t)
$$
\n
$$
z_1 \sim \mu_1
$$

where  $u(z_t) := \int_{\theta \in \Theta} \theta Q_{z_t}(d\theta)$  (i.e., the expected consumption surplus given belief  $Q_{z_t}$ ),  $G(\cdot;\cdot)$  is the transition kernel for  $z_t$  following one's consumption, and  $\mu_1$  is the distribution of  $z_1$ . Compared to the main model, the process of  $(z_t)_{t=1}^T$  now replaces the role of  $(p_t)_{t=1}^T$ . I define the Lagrangian function  $\mathcal{L}(\phi; \lambda)$  and the dual problem similarly as those for the main model. The following lemma follows easily from my assumptions and the definition of  $(z_t)_{t=1}^T$ .

<span id="page-24-2"></span>Lemma A.1. The belief (parameter) process has the following properties: (P1')  $[\rho G(\cdot; z) + (1 - \rho)D(\cdot; z)]$  as a measure-valued function of z is weakly continuous.<sup>[39](#page-24-0)</sup> (P2')  $\int_{z'} u(z') [\rho G(dz'; z) + (1 - \rho)D(dz'; z)] = u(z).$ (P5')  $\mu_1(\{z: u(z) > 0\}) > 0.$ 

These properties are the counterparts to properties (P1), (P2) and (P5) in Lemma [1.](#page-9-5) Because Lemmas [2](#page-10-2) and [3](#page-11-0) in the main text only rely on these properties in Lemma [1,](#page-9-5) they still hold in the current setting.<sup>[40](#page-24-1)</sup> In particular, we still have the following time pattern of Lagrangian multipliers derived from the dual problem:

<span id="page-24-3"></span>**Lemma A.2.** There exists  $\lambda^* \in \arg\min_{\lambda \in \mathbb{R}_+^T} \sup_{\phi \in \Phi} \mathcal{L}(\phi; \lambda)$  such that  $\lambda_t^* \geq \lambda_{t+1}^*$  for all  $t = 1, ..., T - 1.$ 

One particular implication of this lemma is that the multiplier will stay at zero once dropping to it. As in the main model, this implies that any optimal design features a two-phase structure. In the first phase, IC constraints are binding and hence the recommendation policy is just picky enough for the consumers to follow; in the second phase, the IC constraints become non-restrictive and the optimal design follows the optimal continuation policy of the dictator.

While I cannot fully pin down the optimal design, the proposition below reveals an important structure of it. Let  $\geq_{LR}$  denote dominance in the likelihood-ratio order.

<span id="page-24-1"></span><span id="page-24-0"></span> $39$ This is trivially true as  $Z$  is countable.

<sup>&</sup>lt;sup>40</sup>The proofs for them remain the same as before except that the role of  $(p_t)_{t=1}^T$  is now replaced by  $(z_t)_{t=1}^T$ .

<span id="page-25-4"></span>**Proposition A.1.** Assume the consumer arrival rate  $\rho \in (0,1)$ .<sup>[41](#page-25-1)</sup> Any optimal policy is almost surely equivalent to some policy  $\phi^*$  such that for all t: if  $Q_{z'} \geq_{LR} Q_z$  and  $\int_{\theta} \theta dQ_{z}(\theta) > \int_{\theta} \theta dQ_{z}(\theta)$ , then  $\phi_t^*(z) > 0 \Rightarrow \phi_t^*(z') = 1$ .

Intuitively, the proposition roughly suggests that any optimal policy should be more inclined to recommend the product when the current belief of quality is higher in the likelihood-ratio order. This naturally extends the threshold structure of the optimal design in my main model to the current setting, where the platform's belief is in a multidimensional space only endowed with a partial order.

Although the aforementioned property is intuitively appealing, it is actually violated by many recommendation algorithms proposed in papers that only focus on the asymptotic performance of the design. For example, the algorithm in [Mansour et al.](#page-53-3) [\(2020\)](#page-53-3) introduces randomized exploration to fulfill the consumers' IC constraints, which neces-sarily violates the property.<sup>[42](#page-25-2)</sup> My result suggests that modifying their algorithm to be more consistent with this property may help to improve the algorithm's finite-horizon performance. This may be an interesting topic for future algorithm-oriented research.

### <span id="page-25-0"></span>B Proofs

#### B.1 Proof for Lemma [1](#page-9-5)

Proof. Property (P2) is implied by the law of iterated expectation; property (P4) is obvious given Assumption [1\(](#page-7-4)ii) (i.e., the signals are not completely uninformative); property (P5) is directly implied by Assumption [1\(](#page-7-4)i). I show (P1) and (P3) below.

Fix any  $i \geq 1$ . Let S denote the signal realization space of  $s_i$ . Let  $f_L(\cdot)$  and  $f_H(\cdot)$ denote  $s_i$ 's conditional density functions conditional on  $\tilde{\theta} = \theta_L$  and  $\tilde{\theta} = \theta_H$  respectively, with respect to some dominating measure  $m$  over  $S$ . Without loss of generality, we can choose m s.t.  $f_L(s)$  and  $f_H(s)$  are not both equal to zero m-a.s. Since I assume no signal realization fully reveals the value of  $\tilde{\theta}$  (i.e., Assumption [1\(](#page-7-4)iii)), we also have  $f_L(s) \neq 0 \Leftrightarrow f_H(s) \neq 0$  m-a.s. Thus  $f_L(s)$  and  $f_H(s)$  are non-zero m-a.s. Define the log-likelihood ratio  $\ell_i = \log (f_H(s_i)/f_L(s_i))$ , and let  $J_L$  and  $J_H$  denote its distribution given  $\tilde{\theta} = \theta_L$  and  $\tilde{\theta} = \theta_H$  respectively.<sup>[43](#page-25-3)</sup>

We have the following observation:

**Claim** (a). For any a,  $J_H(a) = \int_{\ell \leq a} e^{\ell} dJ_L(\ell)$ .

<span id="page-25-1"></span><sup>&</sup>lt;sup>41</sup>Although I conjecture that the result should also hold with  $\rho = 1$ , my current proof requires  $\rho < 1$ to avoid some technical subtly.

<span id="page-25-2"></span> $^{42}$ The algorithm in [Mansour et al.](#page-53-3) [\(2020\)](#page-53-3) is not Markovian. Hence, more precisely, it is the randomized Markov policy outcome-equivalent to their algorithm that does not satisfy the property.

<span id="page-25-3"></span><sup>&</sup>lt;sup>43</sup>Such log-likelihood ratio representation of a signal has been previously used in Smith  $\&$  Tian [\(2018\)](#page-53-11).

*Proof for Claim (a).* The following equalities hold:

$$
\int_{\ell \le a} e^{\ell} dJ_{L}(\ell) \stackrel{\text{(1)}}{=} \mathbb{E}[\mathbb{1}_{\{\ell_{i} \le a\}} e^{\ell_{i}} | \tilde{\theta} = \theta_{L}] \stackrel{\text{(2)}}{=} \mathbb{E}[\mathbb{1}_{\left\{\log \frac{f_{H}(s_{i})}{f_{L}(s_{i})} \le a\right\}} \frac{f_{H}(s_{i})}{f_{L}(s_{i})} | \tilde{\theta} = \theta_{L}]
$$
\n
$$
\stackrel{\text{(3)}}{=} \int \mathbb{1}_{\left\{\log \frac{f_{H}(s)}{f_{L}(s)} \le a\right\}} \frac{f_{H}(s)}{f_{L}(s)} f_{L}(s) m(ds) \stackrel{\text{(4)}}{=} \int \mathbb{1}_{\left\{\log \frac{f_{H}(s)}{f_{L}(s)} \le a\right\}} f_{H}(s) m(ds)
$$
\n
$$
\stackrel{\text{(5)}}{=} \mathbb{E}[\mathbb{1}_{\{\ell_{i} \le a\}} | \tilde{\theta} = \theta_{H}] \stackrel{\text{(6)}}{=} J_{H}(a)
$$

where the first and the last equalities hold by the definitions of  $J_L$  and  $J_H$  respectively; the second equality holds by the definition of  $\ell_i$ ; the third and the fifth equalities hold by the definitions of  $f_L$  and  $f_H$  respectively; the fourth equality is a trivial identity.  $\Box$ 

Now, given any prior belief p about  $\tilde{\theta} = \theta_H$ , let  $\tilde{p}$  denote the posterior belief given  $s_i$ . Then by the Bayes rule we have:  $\log \frac{\tilde{p}}{1-\tilde{p}} = \log \frac{p}{1-p} + \ell_i$ . Let  $\mathbb{P}_p$  denote the probability measure given prior  $p$ . This then implies that

$$
\mathbb{P}_p(\tilde{p} \le x) = \mathbb{P}_p(\ell_i \le \log \frac{x}{1-x} - \log \frac{p}{1-p})
$$
\n(B.1)

$$
= pJ_H \left( \log \frac{x}{1-x} - \log \frac{p}{1-p} \right) + (1-p)J_L \left( \log \frac{x}{1-x} - \log \frac{p}{1-p} \right) \tag{B.2}
$$

<span id="page-26-2"></span><span id="page-26-0"></span>
$$
= \int_{\ell \le \log \frac{x}{1-x} - \log \frac{p}{1-p}} [pe^{\ell} + (1-p)] dJ_L(\ell)
$$
 (B.3)

where the last equality holds by Claim (a) above. Now, pick any  $p^* \in \mathbb{R}$  and a sequence of  $(p_n)_n \to p^*$ . When expression [\(B.3\)](#page-26-0) is continuous in x at  $x = x_0$  given  $p = p^*$ , obviously we must have  $J_L(\ell)$  being continuous at  $\ell = \log \frac{x_0}{1-x_0} - \log \frac{p^*}{1-p}$  $\frac{p^*}{1-p^*}$ , which further implies that the expression [\(B.3\)](#page-26-0) is continuous in p at  $p = p^*$  given  $x = x_0$ . Thus  $\mathbb{P}_{p^*}(\tilde{p} \leq x)$  being continuous in x at  $x = x_0$  implies  $\mathbb{P}_{p_n}(\tilde{p} \le x_0) \to \mathbb{P}_{p^*}(\tilde{p} \le x_0)$  as  $n \to \infty$ . Therefore, the distribution of  $\tilde{p}$  given prior p is weakly continuous in p. This proves the weak continuity condition for  $G(\cdot|p)$  in (P1).

To check property (P3), we need the following observation:

## **Claim** (b). For any a,  $\int_{\ell \leq a} e^{\ell} dJ_{L}(\ell) \leq \int_{\ell \leq a} dJ_{L}(\ell)$ .

*Proof for Claim (b)*. Notice by Claim (a) above,  $e^{\ell}dJ_{L}(\ell)$  just equals to  $dJ_{H}(\ell)$ . We can thus treat both  $e^{\ell}dJ_{L}(\ell)$  and  $dJ_{L}(\ell)$  as probability measures over  $\mathbb{R}$ , with densities  $e^{\ell}$ and 1 respectively w.r.t. the dominating measure  $dJ<sub>L</sub>(\ell)$ . Since  $e^{\ell}$  is increasing in  $\ell$ , we then have  $e^{\ell}dJ_L(\ell)$  dominating  $dJ_L(\ell)$  in the likelihood-ratio order.<sup>[44](#page-26-1)</sup> This further implies dominance in first-order stochastic dominance and thus  $\int_{\ell \leq a} e^{\ell} dJ_L(\ell) \leq \int_{\ell \leq a} dJ_L(\ell)$ .  $\Box$ 

<span id="page-26-1"></span><sup>&</sup>lt;sup>44</sup>See, e.g., section 1.4 in Müller & Stoyan [\(2002\)](#page-53-12) for an introduction to such order.

Now, pick any  $p_a$  and  $p_b$  s.t.  $p_a < p_b$ , we have

$$
\int_{\ell \le \log \frac{x}{1-x} - \log \frac{p_b}{1-p_b}} [p_b e^{\ell} + (1-p_b)] dJ_L(\ell) \le \int_{\ell \le \log \frac{x}{1-x} - \log \frac{p_a}{1-p_a}} [p_b e^{\ell} + (1-p_b)] dJ_L(\ell)
$$
  

$$
\le \int_{\ell \le \log \frac{x}{1-x} - \log \frac{p_a}{1-p_a}} [p_a e^{\ell} + (1-p_a)] dJ_L(\ell)
$$

where the second inequality holds due to Claim (b). Together with equations  $(B.1)$ – $(B.3)$ , this implies that  $\mathbb{P}_p(\tilde{p} \leq x)$  is weakly decreasing in p for any x. Thus we have the property of (P3).

Q.E.D.

#### B.2 Proof for Proposition [1](#page-10-3)

*Proof.* The proof basically applies Lemma  $1(iv)$  in [Feinberg & Piunovskiy](#page-52-11) [\(2000\)](#page-52-11) to my setting. Specifically, define

$$
\mathcal{V} = \{ v \in \mathbb{R}^{T+1} : \exists \phi \in \Phi \text{ s.t. } v_t = \mathbb{E}_{\phi}[a_t u(p_t)] \,\forall t = 1, ...T \text{ and } v_{T+1} = \sum_{t=1}^T \mathbb{E}_{\phi}[a_t u(p_t)] \}
$$

Notice for each admissible policy  $\phi$ , the first T arguments of the corresponding vector v are the values of the IC constraints and the  $(T+1)$ 'th argument of v is just the total surplus in the designer's objective. Lemma  $1(iv)$  in [Feinberg & Piunovskiy](#page-52-11) [\(2000\)](#page-52-11) implies that V is a compact set. This further implies that the set  $\mathcal{V} \cap \{v \in \mathbb{R}^{T+1} : v_t \geq 0 \forall t = 1, ...T\}$ is compact, and thus when we maximize over its  $(T + 1)$ 'th dimension, the supremum is achievable. By the definition of  $\mathcal{V}$ , this is equivalent to that the designer's problem has its supremum achieved by some  $\phi$ .

Now, it suffices to check that the four conditions of Lemma 1 in [Feinberg & Piunovskiy](#page-52-11) [\(2000\)](#page-52-11) are indeed satisfied in my setting. Condition 1 holds because my state space [0, 1] is closed, and the set of feasible actions  $A = \{0, 1\}$  is finite and does not vary in time and state. Conditions 2 and 4 hold because the flow payoffs in my setting are bounded and continuous in the pair of action and state, and is non-zero for only finitely many periods. For Condition 3, we just need to show the transition probability  $aG(\cdot|p) + (1 - a)D(\cdot|p)$ is weakly-continuous in  $(a, p) \in \{0, 1\} \times [0, 1]$ . With  $\{0, 1\}$  endowed with the discrete topology, it suffices to check weak continuity in p when  $a = 1$  and  $a = 0$  separately. These are respectively implied by the weak continuity of  $G(\cdot|p)$  (Property (P[1\)](#page-9-5) in Lemma 1) and  $D(\cdot|p)$  in  $p^{45}$  $p^{45}$  $p^{45}$ 

#### B.3 Proof for Lemma [2](#page-10-2)

Proof. To use the Lagrangian duality theorem, I first transform the designer's problem into a linear program. Throughout, I fix the initial belief state distribution  $\mu_1$ . Given any policy  $\phi \in \Phi$ , let  $m_t^{\phi}$  denote the distribution of  $(a_t, p_t)$  under it.<sup>[46](#page-28-1)</sup>

Let M denote the set of all sequences of such distributions under some  $\phi$ , i.e.,  $\mathcal{M} =$  $\{(m_t^{\phi})\}$  $(t)_{t=1}^T: \phi \in \Phi$ . A (standard) characterization for this set is that  $(m_t)_{t=1}^T \in \mathcal{M}$  if and only if:

<span id="page-28-3"></span><span id="page-28-2"></span>
$$
m_1(\{0,1\} \times B) = \mu_1(B) \tag{B.4}
$$

$$
m_{t+1}(\{0,1\} \times B) = \int_{p \in [0,1]} \sum_{a \in \{0,1\}} [aG(B|p) + (1-a)D(B|p)]m_t(a, dp) \,\forall t = 1, ..., T-1
$$
\n(B.5)

for any  $B \in \mathcal{B}_{[0,1]}$  (Borel  $\sigma$ -field of  $[0,1]$ ). I use  $\mathcal M$  to denote the set of  $(m_t)_{t=1}^T$  satisfying these conditions. The fact that  $\mathcal{M} \subset \widehat{\mathcal{M}}$  is obvious since any  $(m_t^{\phi})$  $(t)_{t=1}^{\phi}$  must be consistent with  $\mu_1$  and the transition probabilities, and thus satisfies [\(B.4\)](#page-28-2) and [\(B.5\)](#page-28-3).

To see  $\widehat{\mathcal{M}} \subset \mathcal{M}$ , pick any  $(m_t^*)_{t=1}^T \in \widehat{\mathcal{M}}$ . Let  $\phi^*$  be a (randomized) Markov policy such that  $\phi_t^*$  is just the conditional probability mass function of  $a_t$  given  $p_t$  under  $m_t^*$ . Formally, for any  $m_t$ , treat  $m_t(a, dp)$   $(a = 1, 2)$  as a measure over [0, 1] s.t.  $m_t(a, B) =$  $m_t({a} \times B)$ ,  $\forall B \in \mathcal{B}_{[0,1]}$ . Then  $\phi^*$  is defined as (an arbitrary version of) the Radon-Nikodym derivative of  $m_t^*(1, dp)$  w.r.t.  $m_t^*(0, dp) + m_t^*(1, dp)$ . (Notice  $m_t^*(0, dp) + m_t^*(1, dp)$ is just the marginal distribution of  $m_t^*$  over [0, 1] and the Radon-Nikodym derivative is by definition measurable.) Then, we can show  $\phi^*$  implements  $(m_t^*)_{t=1}^T$  by induction in t. Let  $m_t^{\phi^*}$  denote the joint distribution of  $(a_t, p_t)$  under  $\phi^*$  for any t. For  $t = 1$ , we have for all  $B \in \mathcal{B}_{[0,1]}$ :

$$
m_1^{\phi^*}(\{1\} \times B) = \int_{p \in B} \phi_1^*(p) \mu_1(dp) = \int_{p \in B} \phi_1^*(p) [m_1^*(0, dp) + m_1^*(1, dp)]
$$
  
= 
$$
\int_{p \in B} m_1^*(1, dp) = m_1^*(\{1\} \times B)
$$

<span id="page-28-0"></span><sup>&</sup>lt;sup>45</sup>To see  $D(\cdot|p)$  is weakly-continuous in p, notice its cdf is just  $\mathbb{1}_{\{x\geq p\}}$ . Given any sequence  $(p_n)_n \to p^*$ , we have  $1_{\{x\geq p_n\}} \to 1_{\{x\geq p^*\}}$  for any  $x \neq p^*$ . The weak-continuity is thus implied (see, e.g., Section 3.2) in [Durrett](#page-52-12) [\(2019\)](#page-52-12) for conditions of weak-continuity).

<span id="page-28-1"></span><sup>&</sup>lt;sup>46</sup>As is standard, we can construct the underlying measurable space for the process as  $({0, 1} \times [0, 1])^T$ with the Borel  $\sigma$ -field, and treat the corresponding random variables as identity mappings ({0,1}  $\times$  $[0,1]^T \to (\{0,1\} \times [0,1])^T$ . Thus we can treat any distribution for those random variables equivalently as a measure over the underlying measurable space, which is typically how I interpret those distributions.

where the second equality holds by condition [\(B.4\)](#page-28-2) and the third equality holds by the definition of  $\phi^*$ . Since  $m_1^{\phi^*}$  $j_1^{\phi^*}(\{1\} \times B) + m_1^{\phi^*}$  $y_1^{\phi^*}(\{0\}\times B) = \mu_1(B) = m_1^*(\{1\}\times B) + m_1^*(\{0\}\times B),$ we also have  $m_1^{\phi^*}$  $\phi^*$  ({0} × B) =  $m_1^*(\{0\} \times B)$ . Thus  $m_1^{\phi^*} = m_1^*$ .

Now, assume  $m_t^{\phi^*} = m_t^*$  and consider the result for  $t + 1$ . Because condition [\(B.5\)](#page-28-3) holds for  $m_{t+1}^*$ , we know that the marginal distribution over [0, 1] under  $m_{t+1}^*$  given  $m_t^*$  is determined by the same rule as that determines the marginal distribution over [0, 1] under  $m_{t+1}^{\phi^*}$  given  $m_t^{\phi^*}$  $u_t^{\phi^*}$ . Thus  $m_t^* = m_t^{\phi^*}$  $\varphi^*$  implies  $m^*_{t+1}(\{0,1\}\times B) = m^{\varphi^*}_{t+1}(\{0,1\}\times B)$ , ∀B ∈  $\mathcal{B}_{[0,1]}$ . This further implies:

$$
m_{t+1}^{\phi^*}(\{1\} \times B) = \int_{p \in B} \phi_{t+1}^*(p) [m_{t+1}^{\phi^*}(0, dp) + m_{t+1}^{\phi^*}(1, dp)]
$$
  
= 
$$
\int_{p \in B} \phi_{t+1}^*(p) [m_{t+1}^*(0, dp) + m_{t+1}^*(1, dp)] = m_{t+1}^*(\{1\} \times B)
$$

where the second equality holds because the two measures are equal as mentioned right above and the third equality holds by the definition of  $\phi^*$ . Together with  $m^*_{t+1}(\{0,1\} \times$  $B) = m_{t+1}^{\phi^*}(\{0,1\} \times B)$ , this also implies  $m_{t+1}^{\phi^*}(\{0\} \times B) = m_{t+1}^*(\{0\} \times B)$ . Therefore,  $m_{t+1}^* = m_{t+1}^{\phi^*}$ . This completes the induction proof for showing that  $m^*$  is implemented with  $\phi^*$ .

The above discussion has shown  $\mathcal{M} = \widehat{\mathcal{M}}$ . We can thus rewrite the designer's problem as

$$
\max_{(m_t)_{t=1}^T \in \widehat{\mathcal{M}}} \Big\{ \sum_{t=1}^T \Big[ \int_{p \in [0,1]} \sum_{a \in \{0,1\}} a u(p) m_t(a, dp) \Big] \Big\}
$$
  
s.t. 
$$
\int_{p \in [0,1]} \sum_{a \in \{0,1\}} a u(p) m_t(a, dp) \ge 0 \quad \forall t = 1, ..., T
$$

Since conditions [\(B.4\)](#page-28-2) and [\(B.5\)](#page-28-3) are affine in  $(m_t)_{t=1}^T$ , the set  $\mathcal M$  is a convex subset of {signed Borel measures on  $\{0,1\} \times [0,1]$ <sup>T</sup>. The optimization above is thus a linear program over this convex set  $\widehat{\mathcal{M}}$ .

Let  $\widehat{\mathcal{L}}((m_t)_{t=1}^T; \lambda) := \sum_{t=1}^T (1+\lambda_t) \big[ \int_{p \in [0,1]} \sum_{a \in \{0,1\}} au(p) m_t(a, dp) \big],$  i.e., the Lagrangian function associated to the linear program. Since  $u(\cdot)$  is bounded by Lemma [1](#page-9-5) and  $T < \infty$ , the optimal value  $w^*$  is finite. Standard Lagrangian duality (e.g., Theorem 1 in Section 8.6 of [Luenberger](#page-53-13)  $(1997)$  then implies:<sup>[47](#page-29-0)</sup>

$$
w^* = \min_{\lambda \in \mathbb{R}_+^T} \sup_{(m_t)_{t=1}^T \in \widehat{\mathcal{M}}} \widehat{\mathcal{L}}((m_t)_{t=1}^T; \lambda)
$$

<span id="page-29-0"></span><sup>&</sup>lt;sup>47</sup>Theorem 1 in Section 8.6 of [Luenberger](#page-53-13) [\(1997\)](#page-53-13) does not directly state the sufficiency of conditions (i) – (iii) for optimality. However, this is obvious as those conditions together imply  $(m_t^*)_{t=1}^T$  is feasible and  $\sum_{t=1}^T \left[ \int_{p \in [0,1]} \sum_{a \in \{0,1\}} au(p) m_t^*(a, dp) \right] = \sum_{t=1}^T (1 + \lambda_t) \left[ \int_{p \in [0,1]} \sum_{a \in \{0,1\}} au(p) m_t^*(a, dp) \right] = w^*$ .

where the minimum is achieved by some non-negative  $\lambda^*$ . Given any such  $\lambda^*$ ,  $(m_t^*)_{t=1}^T \in$  $\widehat{\mathcal{M}}$  solves the linear program if and only if:

- (i)  $(m_t^*)_{t=1}^T \in \arg \max_{(m_t)_{t=1}^T \in \widehat{\mathcal{M}}} \mathcal{L}((m_t)_{t=1}^T; \lambda^*)$
- (ii)  $\lambda_t^* \int_{p \in [0,1]} \sum_{a \in \{0,1\}} au(p) m_t^*(a, dp) = 0, \forall t = 1, ..., T$
- (iii)  $\int_{p \in [0,1]} \sum_{a \in \{0,1\}} au(p) m_t^*(a, dp) \geq 0, \forall t = 1, ..., T$

To check the corresponding Slater's condition, notice by properties (P2) and (P5) in Lemma [1,](#page-9-5) the consumer's surplus will be strictly positive at all  $t$  under the myopically optimal policy, and thus all IC constraints can hold strictly.

Finally, notice  $\mathcal{M} = \mathcal{M}$  just means that  $(m_t)_{t=1}^T \in \mathcal{M}$  if and only if it is induced by some  $\phi \in \Phi$ . Thus the above results are equivalent to the statements in the lemma. Q.E.D.

#### <span id="page-30-3"></span>B.4 Analyses and Proofs for Section [3.3](#page-11-2)

#### B.4.1 Preliminaries

Towards using the duality result, I start with examining the Lagrangian function optimization max<sub> $\phi \in \Phi$ </sub>  $\mathcal{L}(\phi; \lambda)$  given any generic multiplier  $\lambda \in \mathbb{R}^T_+$ . For this *unconstrained* Markov decision problem, let  $V_{\lambda}(\cdot,t)$  be the value function at time t, which is inductively defined with the Bellman equation:

$$
V_{\lambda}(\cdot, T+1) \equiv 0 \tag{B.6}
$$

<span id="page-30-1"></span>
$$
V_{\lambda}(p,t) = \max\left\{(1+\lambda_t)u(p) + \int_{p'} V_{\lambda}(p',t+1)G(dp'|p), V_{\lambda}(p,t+1)\right\} \forall t = 1,...,T
$$
\n(B.7)

where the two arguments in the maximization correspond to the values with and without time-t consumption of the product respectively. I define  $H_{\lambda}(p, t)$  to be the difference between these two values, i.e.,

<span id="page-30-2"></span>
$$
H_{\lambda}(p,t) := (1 + \lambda_{t})u(p) + \int_{p'} V_{\lambda}(p', t+1)G(dp'|p) - V_{\lambda}(p, t+1) \,\forall t = 1, ..., T \quad (B.8)
$$

Intuitively,  $V_{\lambda}(p, t)$  is the continuation value for the Lagrangian optimization at time t given  $p_t = p$ ;  $H_\lambda(p, t)$  measures the net benefit from inducing the time-t consumption of the product given  $p_t = p$ . A preliminary result needed later is that  $H_\lambda(\cdot, t)$  is continuous.

#### <span id="page-30-0"></span>**Lemma B.1.**  $H_{\lambda}(\cdot, t)$  is continuous for any t.

*Proof.* Since  $u(\cdot)$  is continuous by definition, we just need to show  $V_{\lambda}(p, t + 1)$  and  $\int_{p'} V_\lambda(p', t + 1) G(dp'|p)$  are continuous in p. When  $t = T$ , these hold by the definition of  $V_\lambda(\cdot, T+1)$ . Given that they hold for time t, by the Bellman equation we know

 $V_{\lambda}(p, t)$  is also continuous in p. Furthermore, because  $G(dp'|p)$  is weakly continuous in p and  $V_{\lambda}(p, t)$  is bounded due to the boundedness of  $u(\cdot)$ , this also implies the continuity of  $\int_{p'} V_\lambda(p',t)G(dp'|p)$  in p.<sup>[48](#page-31-0)</sup> The proof is thus completed by (backward) induction in t.  $Q.E.D.$ 

We have the standard dynamic programming result:

<span id="page-31-2"></span>**Lemma B.2.** Given any multiplier  $\lambda$  and initial belief state distribution  $\mu_1$ , we have:

- (a)  $\max_{\phi \in \Phi} \mathcal{L}(\phi; \lambda) = \int_p V_{\lambda}(p, 1) \mu_1(dp);$
- (b) A policy  $\phi^{\lambda} \in \arg \max_{\phi \in \Phi} \mathcal{L}(\phi; \lambda)$  if and only if  $H_{\lambda}(p_t, t) > 0 \Rightarrow a_t = 1$  and  $H_{\lambda}(p_t, t) < 0 \Rightarrow a_t = 0$  almost surely under it.

*Proof.* Pick any  $\phi \in \Phi$ . Notice that by the definition of  $V_{\lambda}$ , we have

$$
V_{\lambda}(p_t, t) \ge \phi_t(p_t) \Big[ (1 + \lambda_t) u(p_t) + \int_{p'} V_{\lambda}(p', t+1) G(dp'|p_t) \Big] + (1 - \phi_t(p_t)) V_{\lambda}(p_t, t+1)
$$
  

$$
\stackrel{\text{a.s.}}{=} \mathbb{E}_{\phi} [(1 + \lambda_t) a_t u(p_t) |p_t] + \mathbb{E}_{\phi} [V_{\lambda}(p_{t+1}, t+1) |p_t] \tag{B.9}
$$

where the inequality holds as equality if and only if  $H_{\lambda}(p_t, t) > 0 \Rightarrow \phi_t(p_t) = 1$  and  $H_{\lambda}(p_t, t) < 0 \Rightarrow \phi_t(p_t) = 0.$ 

<span id="page-31-1"></span>Using this repeatedly, we have:

$$
\int_{p} V_{\lambda}(p,1)\mu_{1}(dp) = \mathbb{E}_{\phi}[V_{\lambda}(p_{1},1)] \geq \mathbb{E}_{\phi}[(1+\lambda_{1})a_{1}u(p_{1})] + \mathbb{E}_{\phi}[V_{\lambda}(p_{2},2)]
$$
\n
$$
\geq \mathbb{E}_{\phi}[(1+\lambda_{1})a_{1}u(p_{1})] + \mathbb{E}_{\phi}[(1+\lambda_{2})a_{2}u(p_{2})] + \mathbb{E}_{\phi}[V_{\lambda}(p_{3},3)]
$$
\n
$$
\dots \geq \mathbb{E}_{\phi}[(1+\lambda_{1})a_{1}u(p_{1})] + \dots + \mathbb{E}_{\phi}[(1+\lambda_{T})a_{T}u(p_{T})] + \underbrace{\mathbb{E}_{\phi}[V_{\lambda}(p_{T+1},T+1)]}_{=0}
$$

Notice the last line is just  $\mathcal{L}(\phi; \lambda)$ . This shows  $\int_p V_\lambda(p, 1) \mu_1(dp) \ge \sup_{\phi \in \Phi} \mathcal{L}(\phi; \lambda)$ . Moreover, notice that all these inequalities hold as equalities if and only if the inequality in  $(B.9)$  holds as equality for all t almost surely under  $\phi$ . As is mentioned earlier, this is equivalent to  $H_{\lambda}(p_t, t) > 0 \Rightarrow \phi_t(p_t) = 1$  and  $H_{\lambda}(p_t, t) < 0 \Rightarrow \phi_t(p_t) = 0$  almost surely under  $\phi$ . If there is indeed a measurable  $\phi$  satisfying these properties, then we have  $\int_p V_\lambda(p,1)\mu_1(dp) = \sup_{\phi \in \Phi} \mathcal{L}(\phi;\lambda)$ , the supremum is achieved by such policy, and any other admissible policy is optimal if and only if it also satisfies these properties. Therefore, to prove the lemma, it now suffices to show that there is indeed a measurable policy satisfying  $H_{\lambda}(p_t, t) > 0 \Rightarrow \phi_t(p_t) = 1$  and  $H_{\lambda}(p_t, t) < 0 \Rightarrow \phi_t(p_t) = 0$  almost surely. We can construct such policy by defining  $\phi_t(p) = \mathbb{1}_{\{H_\lambda(p,t)\geq 0\}}$ , where  $\mathbbm{1}$  is the indicator function. It is indeed measurable since  $H_{\lambda}(\cdot, t)$  is continuous by Lemma [B.1.](#page-30-0)  $Q.E.D.$ 

I now provide some basic properties for the value function  $V_{\lambda}$ .

<span id="page-31-0"></span><sup>&</sup>lt;sup>48</sup>Pick any  $(p_n)_n \to p^*$ , the weak continuity implies  $G(dp'|p_n) \stackrel{w}{\to} G(dp'|p^*)$ , which further implies  $\int f(p')G(dp'|p_n) \to \int f(p')G(dp'|p^*)$  for any bounded and continuous function f.

<span id="page-32-0"></span>**Lemma B.3.** Given any  $\lambda \in \mathbb{R}^T_+$ , we have: (a)  $V_{\lambda}(p, t)$  is (weakly) increasing in p; (b)  $\int_{p'} V_{\lambda}(p',t) G(dp'|p) \geq V_{\lambda}(p,t)$  for any pair of  $(p,t)$ .

Proof. The results can be shown by backward induction in t using the Bellman equation. Since  $V_{\lambda}(\cdot, T+1) \equiv 0$ , both properties hold trivially for  $t = T + 1$ .

Now, assume property (a) holds for  $V_{\lambda}(\cdot, t + 1)$  $V_{\lambda}(\cdot, t + 1)$  $V_{\lambda}(\cdot, t + 1)$ . Then property (P3) in Lemma 1 implies that  $\int_{p'} V_\lambda(p', t + 1) G(dp'|p)$  increases in p. Together with the monotonicity of  $u(\cdot)$  and the fact that  $\lambda_t \geq 0$ , we know the RHS of the Bellman equation [\(B.7\)](#page-30-1) is increasing in p. This shows property (a) also holds for  $V_{\lambda}(\cdot, t)$  and concludes the proof for part (a).

For property (b), assuming it holds for all periods later than  $t$ , we need to show  $\int_{p'} V_\lambda(p',t) G(dp'|p) \geq V_\lambda(p,t)$ . Substituting the Bellman equation in, we know this is equivalent to:

$$
\int_{p'} \max \left\{ (1 + \lambda_t) u(p') + \int_{p''} V_{\lambda}(p'', t + 1) G(dp''|p'), V_{\lambda}(p', t + 1) \right\} G(dp'|p)
$$
  
 
$$
\geq \max \left\{ (1 + \lambda_t) u(p) + \int_{p'} V_{\lambda}(p', t + 1) G(dp'|p), V_{\lambda}(p, t + 1) \right\}
$$

It then suffices to check:

$$
\int_{p'} \left( (1 + \lambda_t) u(p') + \int_{p''} V_{\lambda}(p'', t + 1) G(dp''|p') \right) G(dp'|p)
$$
  
\n
$$
\geq (1 + \lambda_t) u(p) + \int_{p'} V_{\lambda}(p', t + 1) G(dp'|p)
$$
  
\nand 
$$
\int_{p'} V_{\lambda}(p', t + 1) G(dp'|p) \geq V_{\lambda}(p, t + 1)
$$

The second of these inequalities is directly implied by the induction hypothesis. To check the first one, notice by property (P2) in Lemma [1,](#page-9-5) we have  $\int_{p'}(1+\lambda_t)u(p')G(dp'|p)$  $(1 + \lambda_t)u(p)$ . Moreover, by the induction hypothesis we have  $\int_{p''} V_\lambda(p'', t + 1)G(dp''|p') \ge$  $V_{\lambda}(p', t + 1)$  for any p', and thus  $\int_{p'} \int_{p''} V_{\lambda}(p'', t + 1) G(dp''|p') G(dp'|p) \ge \int_{p'} V_{\lambda}(p', t + 1)$  $1)G(dp'|p)$ . These together imply the first inequality above. Thus property (b) holds for period t. This completes the proof by induction.

 $Q.E.D.$ 

#### B.4.2 Proof for Lemma [3](#page-11-0)

Proof. Pick any  $\lambda' \in \mathbb{R}_+^T$  such that  $\lambda'_\tau < \lambda'_{\tau+1}$  for some  $\tau$ . Define  $\lambda''$  to be equal to  $\lambda'$ except for terms of time  $\tau$  and  $\tau + 1$ , which are defined as:  $\lambda''_{\tau} = \lambda'_{\tau+1}$  and  $\lambda''_{\tau+1} = \lambda'_{\tau}$ . The key to the proof is the following observation:

**Claim**. Let  $\lambda'$  and  $\lambda''$  be defined as above. Then  $\sup_{\phi \in \Phi} \mathcal{L}(\phi; \lambda') \geq \sup_{\phi \in \Phi} \mathcal{L}(\phi; \lambda'')$ .

Proof for the Claim. Since  $\lambda'$  and  $\lambda''$  agree for  $t < \tau$ , any policy will lead to the same flow payoffs for periods before  $\tau$  for both  $\mathcal{L}(\phi; \lambda')$  and  $\mathcal{L}(\phi; \lambda'')$ . It thus suffices to show  $V_{\lambda'}(p,\tau) \geq V_{\lambda''}(p,\tau)$  given any p.

Since  $\lambda'$  and  $\lambda''$  also agree for  $t \geq \tau + 2$ , we have  $V_{\lambda'}(\cdot, \tau + 2) = V_{\lambda''}(\cdot, \tau + 2)$ . I can thus let  $V_*(\cdot, \tau+2)$  to denote both of them, i.e.,  $V_*(\cdot, \tau+2) := V_{\lambda}(\cdot, \tau+2)$  (=  $V_{\lambda}(\cdot, \tau+2)$ ). Also let  $(y_1, y_2) := (\lambda'_\tau, \lambda'_{\tau+1})$ . Then  $y_1 < y_2$  and  $(\lambda''_\tau, \lambda''_{\tau+1}) = (y_2, y_1)$ . By the Bellman equation  $(B.7)$ , we have:

$$
V_{\lambda''}(p,\tau) = \max\left\{ (1+y_2)u(p) + \int_{p'} V_{\lambda''}(p',\tau+1)G(dp'|p), V_{\lambda''}(p,\tau+1) \right\}
$$
(B.10)

Consider the following two cases:

• Case 1: the maximum in equation [\(B.10\)](#page-33-0) is achieved with  $a<sub>\tau</sub> = 0$  (i.e., no consumption).

In this case, we have:

<span id="page-33-0"></span>
$$
V_{\lambda''}(p,\tau) = V_{\lambda''}(p,\tau+1)
$$
  
= max  $\left\{ (1 + y_1)u(p) + \int_{p'} V_*(p',\tau+2)G(dp'|p), V_*(p,\tau+2) \right\}$   
 $\leq$  max  $\left\{ (1 + y_1)u(p) + \int_{p'} V_{\lambda'}(p',\tau+1)G(dp'|p), V_{\lambda'}(p,\tau+1) \right\}$   
=  $V_{\lambda'}(p,\tau)$ 

where the second equality holds by the Bellman equation for  $V_{\lambda''}(p, \tau+1)$ ; the inequality holds because the Bellman equation for  $V_{\lambda}(p, \tau+1)$  implies that  $V_*(p, \tau+2) \leq V_{\lambda}(p, \tau+2)$ 1) for any p; the last equality is just the Bellman equation for  $V_{\lambda}^{\prime}(p,\tau)$ .

• Case 2: the maximum in equation [\(B.10\)](#page-33-0) is achieved with  $a<sub>\tau</sub> = 1$ . In this case, we have:

$$
V_{\lambda''}(p,\tau) = (1+y_2)u(p) + \int_{p'} V_{\lambda''}(p',\tau+1)G(dp'|p)
$$
\n(B.11)

<span id="page-33-1"></span>
$$
\leq (1+y_1)u(p) + \int_{p'} \Big[ \underbrace{(y_2-y_1)u(p') + V_{\lambda''}(p',\tau+1)}_{=:M(p')} \Big] G(dp'|p) \tag{B.12}
$$

where the inequality holds because  $y_2 > y_1$  and  $u(p) = \int_{p'} u(p') G(dp'|p)$  by property (P2) in Lemma [1.](#page-9-5) Using the Bellman equation for  $V_{\lambda''}(p', \tau + 1)$ , we know the term  $M(p')$  satisfies:

$$
M(p') = (y_2 - y_1)u(p') + \max \left\{ (1 + y_1)u(p') + \int_{p''} V_*(p'', \tau + 2)G(dp''|p'), V_*(p', \tau + 2) \right\}
$$
  
= max  $\left\{ (1 + y_2)u(p') + \int_{p''} V_*(p'', \tau + 2)G(dp''|p'), V_*(p', \tau + 2) + (y_2 - y_1)u(p') \right\}$   
(B.13)

Now, consider properties of  $M(p')$  in two different scenarios about  $p'$ :

- Scenario 1:  $u(p') > 0$ .

Notice Lemma [B.3\(](#page-32-0)b) implies  $\int_{p''} V_*(p'', \tau + 2) G(dp''|p') \ge V_*(p', \tau + 2)$ . When  $u(p') > 0$ , we thus have  $(1 + y_2)u(p') + \int_{p''} V_*(p'', \tau + 2)G(dp''|p') \ge V_*(p', \tau + 2) +$  $(y_2-y_1)u(p')$ . Therefore, [\(B.13\)](#page-34-0) implies:

<span id="page-34-0"></span>
$$
M(p') = (1 + y_2)u(p') + \int_{p''} V_*(p'', \tau + 2)G(dp''|p')
$$
  
= max  $\left\{ (1 + y_2)u(p') + \int_{p''} V_*(p'', \tau + 2)G(dp''|p'), V_*(p', \tau + 2) \right\}$   
=  $V_{\lambda'}(p', \tau + 1)$ 

where the second equality holds since the first argument in the bracket is larger when  $\int_{p''} V_*(p'', \tau + 2) G(dp''|p') \ge V_*(p', \tau + 2)$  and  $u(p') > 0$ .  $-$  Scenario 2:  $u(p') \leq 0$ .

In this case, we have  $(y_2 - y_1)u(p') \leq 0$ . [\(B.13\)](#page-34-0) thus implies:

$$
M(p') \le \max\left\{ (1 + y_2)u(p') + \int_{p''} V_*(p'', \tau + 2)G(dp''|p'), V_*(p', \tau + 2) \right\}
$$
  
=  $V_{\lambda'}(p', \tau + 1)$ 

where the equality holds by the Bellman equation.

In both scenarios, we always have  $M(p') \leq V_{\lambda}(p', \tau + 1)$ . Together with inequality [\(B.12\)](#page-33-1), this implies that  $V_{\lambda''}(p, \tau) \le (1 + y_1)u(p) + \int_{p'} V_{\lambda'}(p', \tau + 1)G(dp'|p) \le V_{\lambda'}(p, \tau)$ , where the second inequality is due to the Bellman equation for  $V_{\lambda'}(p, \tau)$ .

In sum,  $V_{\lambda''}(p, \tau) \leq V_{\lambda}(p, \tau)$  in both cases, which completes the proof for the claim.

□

I now go back to the main proof for Lemma [3.](#page-11-0) Pick any  $\lambda^0 \in \arg\min_{\lambda \in \mathbb{R}_+^T} \sup_{\phi \in \Phi} \mathcal{L}(\phi; \lambda)$ (whose existence is guaranteed by Lemma [2\)](#page-10-2). If  $\lambda_t^0$  is already non-increasing in t, we are done; if  $\lambda_t^0 < \lambda_{t+1}^0$  for some t, then the claim above implies that by interchanging terms  $\lambda_t^0$  and  $\lambda_{t+1}^0$ , we will get a new multiplier still in  $\arg\min_{\lambda \in \mathbb{R}_+^T} \sup_{\phi \in \Phi} \mathcal{L}(\phi; \lambda)$ . By repeatedly making such interchanges, we can then derive a multiplier  $\lambda^* \in \argmin_{\lambda \in \mathbb{R}_+^T} \sup_{\phi \in \Phi} \mathcal{L}(\phi; \lambda)$ 

such that  $\lambda_t^* \geq \lambda_{t+1}^*$ ,  $\forall t$ .<sup>[49](#page-35-0)</sup>

Next, I provide some properties of  $H_{\lambda}$ , which are key to the proof for Lemma [4.](#page-12-2)

<span id="page-35-1"></span>**Lemma B.4.**  $H_{\lambda}$  satisfies the following properties:

- (a)  $p > \overline{p} \Rightarrow H_{\lambda}(p, t) > 0$ ,  $\forall t$ ;
- (b) If  $\lambda_t$  is non-increasing in t, then  $H_{\lambda}(p,t)$  is (weakly) increasing in p for any t;
- (c) If  $\lambda_t$  is non-increasing in t, then for any x, y s.t.  $x < y$  and  $H_\lambda(y, t) \leq 0$ , we have  $H_{\lambda}(x, t) < H_{\lambda}(y, t)$  (thus  $H_{\lambda}(\cdot, t) = 0$  has at most one root);
- (d) If  $\lambda_t = \lambda_{t+1}$ , then  $H_\lambda(p, t) \leq 0 \Rightarrow H_\lambda(p, t+1) < 0$ .

*Proof.* Part (a): Because  $\int_{p'} V_{\lambda}(p', t+1)G(dp'|p) - V_{\lambda}(p, t+1) \geq 0$  according to Lemma [B.3,](#page-32-0) part (a) is directly implied by the definition of  $H_{\lambda}$ .

**Part (b):** I prove (b) by backward induction in t. By definition,  $H_{\lambda}(p,T) = (1+\lambda_T)u(p)$ is strictly increasing in p and thus the monotonicity property holds for  $H_{\lambda}(p,T)$ . Now, assuming it holds for  $H_{\lambda}(p, t + 1)$ , I show it also holds for  $H_{\lambda}(p, t)$ . Notice the following

- <span id="page-35-0"></span><sup>49</sup>More specifically, starting with  $n = 0$ , we can run the following algorithm:
	- 1. Let  $\tau_n = \inf\{t : \lambda_t^n < \lambda_{t+1}^n\}$ . If  $\tau_n = +\infty$ , end the algorithm and out-put  $\lambda^n$ ; otherwise, go to the next step.
	- 2. Let  $s = \max\left\{0, \sup\{t < \tau_n : \lambda_t^n \geq \lambda_{\tau_n+1}^n\}\right\}$  and define

$$
\lambda_t^{n+1} = \begin{cases}\n\lambda_{\tau_n+1}^n & \text{if } t = s+1; \\
\lambda_{t-1}^n & \text{if } t = s+2, ..., \tau_n+1; \\
\lambda_t^n & \text{elsewhere}\n\end{cases}
$$

Then, repeat the procedures with n replaced by  $n + 1$ .

Intuitively, in step 2 of the algorithm we advance the first term in  $\lambda^n$  greater than its predecessor to an earlier position such that the first  $\tau_n + 1$  terms will be in descending order. It is then easy to see that this algorithm will end in finite time and the vector it delivers will be non-increasing over t. Moreover, to derive  $\lambda^{n+1}$  from  $\lambda^n$  in step 2, one can just interchange the  $(\tau_n + 1)$ 'th term with the  $\tau_n$ 'th term, then interchange the (new)  $\tau_n$ 'th term with the  $(\tau_n - 1)$ 'th term,..., and finally interchange the (new)  $(s + 2)$ 'th term with the  $(s + 1)$ 'th term. In each of these steps, we interchange two adjacent terms with the latter greater than the former. By the claim proved above, this keeps each of the (intermediate) vector within  $\arg\min_{\lambda\in\mathbb{R}_+^T} \sup_{\phi\in\Phi} \mathcal{L}(\phi;\lambda)$ . Thus the multiplier we derive in the end remains in  $\arg \min_{\lambda \in \mathbb{R}_+^T} \sup_{\phi \in \Phi} \mathcal{L}(\phi; \lambda)$ . This completes the proof.

equations hold:

$$
H_{\lambda}(p,t) = (1 + \lambda_{t})u(p) + \int_{p'} V_{\lambda}(p', t+1)G(dp'|p) - V_{\lambda}(p, t+1)
$$
  
\n
$$
= (1 + \lambda_{t})u(p) + \int_{p'} \left( \max\{H_{\lambda}(p', t+1), 0\} + V_{\lambda}(p', t+2) \right)G(dp'|p)
$$
  
\n
$$
- \left( \max\{H_{\lambda}(p, t+1), 0\} + V_{\lambda}(p, t+2) \right)
$$
  
\n
$$
= (1 + \lambda_{t})u(p) + \int_{p'} V_{\lambda}(p', t+2)G(dp'|p) - V_{\lambda}(p, t+2)
$$
  
\n
$$
+ \int_{p'} \max\{H_{\lambda}(p', t+1), 0\}G(dp'|p) - \max\{H_{\lambda}(p, t+1), 0\}
$$
  
\n
$$
= (1 + \lambda_{t})u(p) - (1 + \lambda_{t+1})u(p) + H_{\lambda}(p, t+1)
$$
  
\n
$$
+ \int_{p'} \max\{H_{\lambda}(p', t+1), 0\}G(dp'|p) - \max\{H_{\lambda}(p, t+1), 0\}
$$
  
\n
$$
= (\lambda_{t} - \lambda_{t+1})u(p) + \min\{H_{\lambda}(p, t+1), 0\} + \int_{p'} \max\{H_{\lambda}(p', t+1), 0\}G(dp'|p)
$$
  
\n(B.14)

<span id="page-36-0"></span>where the second equality is because  $V_{\lambda}(p, t + 1) = \max\{H_{\lambda}(p, t + 1), 0\} + V_{\lambda}(p, t + 2)$ according to the Bellman equation [\(B.7\)](#page-30-1); the first and the fourth equalities are directly implied by the definition of  $H_{\lambda}$ ; the other two are just trivial identities. Recall that:  $\lambda_t \geq \lambda_{t+1}$  by assumption;  $H_\lambda(p, t+1)$  weakly increases in p by the induction hypothesis; and  $G(\cdot|p)$  increases in first-order stochastic dominance in p by property (P3) of Lemma [1.](#page-9-5) These imply that all of the three terms in the last expression are (weakly) increasing in p. Thus  $H_{\lambda}(p,t)$  is (weakly) increasing in p. This completes the proof for (b).

**Part (c):** I still prove by induction. The result holds obviously for  $H_{\lambda}(p,T) = (1 +$  $\lambda_T$ )u(p). Now, assuming it holds for period  $t + 1$ , I show it also holds for period t. In particular, with any  $x < y$  in [0, 1], we want to show  $H_{\lambda}(y, t) \leq 0 \Rightarrow H_{\lambda}(x, t) < H_{\lambda}(y, t)$ . Given result (b) and equation [\(B.14\)](#page-36-0) derived above,  $H_{\lambda}(x, t) < H_{\lambda}(y, t)$  obviously holds when  $\lambda_t > \lambda_{t+1}$ , since  $u(p) = \theta_H p + \theta_L(1-p)$  is strictly increasing in p. It thus suffices to assume  $\lambda_t = \lambda_{t+1}$ . In this case, I have the following observation:

**Claim**. If  $\lambda_t = \lambda_{t+1}$  and  $H_\lambda(y, t) \leq 0$ , then  $H_\lambda(y, t+1) \leq 0$ .

Proof for the claim. Given  $\lambda_t = \lambda_{t+1}$  and  $H_\lambda(y, t) \leq 0$ , suppose  $H_\lambda(y, t+1) > 0$ . Then equation [\(B.14\)](#page-36-0) implies that  $\int_{p'} \max\{H_\lambda(p', t+1), 0\} G(dp'|y) = H_\lambda(y, t) \leq 0$ . However, by the law of iterated expectation condition on the belief process (property (P2) in Lemma [1\)](#page-9-5), we must have  $G([y, 1]|y) > 0$ . Moreover, by the monotonicity of  $H_{\lambda}(\cdot, t+1)$ proved in part (b), we know  $H_{\lambda}(p', t + 1) \geq H_{\lambda}(y, t + 1)$  for any  $p' \geq y$ . Together with the hypothesis  $H_{\lambda}(y, t + 1) > 0$ , these then imply  $\int_{p'} \max\{H_{\lambda}(p', t + 1), 0\} G(dp'|y) > 0$ , which contradicts with the previous conclusion. Thus we must have  $H_{\lambda}(y, t+1) \leq 0$ .  $\Box$ 

Now, go back to the main proof for part (c). Notice when  $\lambda_t = \lambda_{t+1}$  and  $H_\lambda(y, t) \leq 0$ , the following holds:

$$
H_{\lambda}(y,t) = \min\{H_{\lambda}(y,t+1),0\} + \int_{p'} \max\{H_{\lambda}(p',t+1),0\} G(dp'|y)
$$
  
=  $H_{\lambda}(y,t+1) + \int_{p'} \max\{H_{\lambda}(p',t+1),0\} G(dp'|y)$   
>  $H_{\lambda}(x,t+1) + \int_{p'} \max\{H_{\lambda}(p',t+1),0\} G(dp'|x) = H_{\lambda}(x,t)$ 

The first equality is just by equation [\(B.14\)](#page-36-0) with  $\lambda_t = \lambda_{t+1}$ . The second equality holds because the claim proved above implies  $H_{\lambda}(y, t + 1) \leq 0$ . The strict inequality holds because:  $H_{\lambda}(y, t + 1) \leq 0$  further implies  $H_{\lambda}(x, t + 1) < H_{\lambda}(y, t + 1)$  by the induction hypothesis;  $G(\cdot|y)$  first order stochastic dominates  $G(\cdot|x)$ ; and  $H_{\lambda}(\cdot, t+1)$  is increasing by part (b). The last equality holds also by equation [\(B.14\)](#page-36-0) and the fact that  $H_{\lambda}(x, t+1) < 0$ . This completes the proof by induction.

**Part (d):** Suppose  $\lambda_t = \lambda_{t+1}$ ,  $H_{\lambda}(p, t) \leq 0$ , but  $H_{\lambda}(p, t+1) \geq 0$ . Equation [B.14](#page-36-0) would imply

$$
H_{\lambda}(p,t) = \int_{p'} \max\{H_{\lambda}(p',t+1),0\} G(dp'|p)
$$

I now argue that the RHS above must be strictly positive. Notice  $H_{\lambda}(p, t) \leq 0$  obviously imply  $p < 1$  $p < 1$ . Property (P4) in Lemma 1 then implies that  $G((p, 1||p) > 0$ . Moreover, since  $H_{\lambda}(p, t + 1) \ge 0$ , parts (b) and (c) proved earlier imply  $H_{\lambda}(p', t + 1) > 0$  for any  $p' > p$ . These together imply  $\int_{p'} \max\{H_\lambda(p', t+1), 0\} G(dp'|p) > 0$ . This contradicts with  $H_{\lambda}(p, t) \leq 0$  given the equation above. Thus the result in part (d) holds.

 $Q.E.D.$ 

#### B.4.4 Proof for Lemma [4](#page-12-2)

The proof for Lemma [4](#page-12-2) easily follows from Lemma [B.4.](#page-35-1)

*Proof.* For each t, I construct the threshold  $\eta_t$  as follows:

- Case 1:  $\{p : H_{\lambda}(p, t) = 0\} = \emptyset$ . In this case, define  $\eta_t = \inf\{p : H_\lambda(p, t) > 0\}.$
- Case 2:  $\{p: H_\lambda(p, t) = 0\} \neq \emptyset$ . In this case, Lemma [B.4\(](#page-35-1)c) implies that  $\{p : H_\lambda(p, t) = 0\}$  contains a single element. Define  $\eta_t$  to be this element.

By the monotonicity property of  $H_{\lambda}(\cdot, t)$  in Lemma [B.4\(](#page-35-1)b), in both cases we have  $p_t >$  $\eta_t \Rightarrow H_\lambda(p_t, t) > 0$  and  $p_t < \eta_t \Rightarrow H_\lambda(p_t, t) < 0$ . Together with Lemma [B.2\(](#page-31-2)b), this implies that under any solution to  $\max_{\phi \in \Phi} \mathcal{L}(\phi; \lambda)$ , we have  $p_t > \eta_t \Rightarrow a_t = 1$  a.s. and  $p_t < \eta_t \Rightarrow a_t = 0$  a.s. Hence any solution is almost surely equivalent to a threshold policy with thresholds being  $(\eta_t)_{t=1}^T$ .

Moreover, notice Lemma [B.4\(](#page-35-1)a) implies  $u(p_t) > 0 \Rightarrow H_{\lambda}(p_t, t) > 0$ . Together with Lemma [B.2\(](#page-31-2)b), this then implies  $u(p_t) > 0 \Rightarrow a_t = 1$  a.s. under any optimal solution to  $\max_{\phi \in \Phi} \mathcal{L}(\phi; \lambda).$ 

$$
Q.E.D.
$$

#### <span id="page-38-0"></span>B.5 Proofs for Section [3.4](#page-13-1)

#### <span id="page-38-2"></span> $\textbf{B.5.1} \quad \textbf{Definition of}\ \phi^d$

As is mentioned in the main text, I define  $\phi^d$  as the "most conservative" optimal policy for the dictator's problem. Formally, for any  $t = 1, ..., T$ :

$$
\phi_t^d(p) := \begin{cases} 1 & \text{if } H_0(p, t) > 0; \\ 0 & \text{otherwise} \end{cases}
$$
 (B.15)

where  $H_0$  is as defined in [\(B.8\)](#page-30-2) in Appendix [B.4](#page-30-3) with  $\lambda = 0$ . By Lemma [B.2,](#page-31-2) it is easy to see that we not only have  $\phi^d$  being optimal for the dictator's problem, but also have  $\phi^d_{\geq t}$  to be optimal for the dictator's continuation problem starting from time  $t$  regardless of the belief distribution at t. Moreover,  $\phi^d$  (or  $\phi^d_{\geq t}$ ) is conservative in the sense that it breaks any tie in favor of non-recommendation (i.e.,  $H_0(p, t) = 0 \Rightarrow \phi_t^d(p) = 0$ ). This makes it most favorable to the current consumer among all dictator's optimal (continuation) policies.

#### <span id="page-38-1"></span>B.5.2 Details in the Construction of  $\phi^o$  and a Uniqueness Property

I first construct a threshold time-t policy  $\phi_t^o$  satisfying the requirements in step 2 of the algorithm in Definition [2.](#page-13-2) Given  $\mu_t^o$ , define the policy's threshold as

$$
\eta_t^o = \inf\{x \in [0, 1] : \int_{p > x} u(p) \mu_t^o(dp) > 0\}
$$

and define the recommendation probability at the threshold as

$$
\phi_t^o(\eta_t^o) = -\frac{\int_{p>\eta_t^o} u(p)\mu_t^o(dp)}{u(\eta_t^o)\mu_t^o(\{\eta_t^o\})}
$$

When the denominator above is zero, I define  $\phi_t^o(\eta_t^o) = 0$  for simplicity. Now, I check the  $\phi^o_t$  such defined indeed satisfies the desired properties.

*Claim.*  $\phi_t^o$  above is well-defined and satisfies the properties in step 2 of the algorithm.

*Proof.* First notice properties (P2) and (P5) in Lemma [1](#page-9-5) together imply  $\mu_t^o((\bar{p}, 1]) > 0$ and thus  $\int_{p>\bar{p}} u(p) \mu_t^o(dp) > 0$ . This implies  $\eta_t^o \leq \bar{p}$  and thus the first property is satisfied.

Now, I argue that  $\int_{p>\eta_t^o} u(p)\mu_t^o(dp) \geq 0$ . To see this, notice by the definition of  $\eta_t^o$ , there exists a sequence  $\{x_n\} \downarrow \eta_t^o$  such that  $\int_{p>x_n} u(p) \mu_t^o(dp) > 0$  for all n. Since  $\mathbb{1}_{\{p>n\}} \to \mathbb{1}_{\{p>\eta_i^o\}}$  and  $u(\cdot)$  is bounded, we must have  $\int_{p>\eta_i^o} u(p) \mu_i^o(dp) \geq 0$  by the dominated convergence theorem.

Next, I argue that  $\int_{p \ge \eta_t^o} u(p) \mu_t^o(dp) \le 0$ . As the algorithm has not been ended in step 1, we must have  $\int_{p\in[0,1]} u(p) \mu_t^o(dp) < 0^{.50}$  $\int_{p\in[0,1]} u(p) \mu_t^o(dp) < 0^{.50}$  $\int_{p\in[0,1]} u(p) \mu_t^o(dp) < 0^{.50}$  Thus the argument is true when  $\eta_t^o = 0$ . When  $\eta_t^o > 0$ , notice by the definition of  $\eta_t^o$ , there exists a sequence  $\{x_n\} \uparrow \eta_t^o$  such that  $\int_{p>x_n} u(p)\mu_t^o(dp) \leq 0$  for all n. Since  $\mathbb{1}_{\{p>x_n\}} \to \mathbb{1}_{\{p\geq \eta_t^o\}}$ , by the dominated convergence theorem we then must have  $\int_{p \ge \eta_t^o} u(p) \mu_t^o(dp) \le 0.$ 

Now, consider two cases:

- Case 1:  $\mu_t^o(\{\eta_t^o\}) = 0$ . In this case, the above arguments imply  $\int_{p > \eta_t^o} u(p) \mu_t^o(dp) = 0$ . Therefore the second property is satisfied.
- Case 2:  $\mu_t^o(\{\eta_t^o\}) > 0$ . In this case, the above arguments imply  $\int_{p > \eta_t^o} u(p) \mu_t^o(dp) \ge 0$ t and  $\int_{p>\eta_t^o} u(p)\mu_t^o(dp) + u(\eta_t^o)\mu_t^o(\{\eta_t^o\}) \leq 0$ . These imply  $0 \geq \frac{\int_{p>\eta_t^o} u(p)\mu_t^o(dp)}{u(\eta_t^o)\mu_t^o(\{\eta_t^o\})}$  $\frac{\frac{p>\eta_t^{\alpha}}{w(\eta_t^o)\mu_t^o(\{\eta_t^o\})}}{u(\eta_t^o)\mu_t^o(\{\eta_t^o\})} \geq$ -1 whenever  $u(\eta_t^o)\mu_t^o(\{\eta_t^o\}) \neq 0$ . Thus  $\phi_t^o(\eta_t^o)$  defined above is a valid probability. By the definition of  $\phi_t^o(\eta_t^o)$ , we have  $\int_p \phi_t^o(p)u(p)\mu_t^o(dp) = \int_{p>\eta_t^o} u(p)\mu_t^o(dp) +$  $\phi_t^o(\eta_t^o)u(\eta_t^o)\mu_t^o(\{\eta_t^o\})=0.$  Thus  $\phi_t^o$  satisfies the second desired property.

$$
Q.E.D.
$$

Due to the possible existence of off-path belief states, there can also be other forms of  $\phi_t$  satisfying the desired properties in step 2 of the algorithm. However, the following lemma implies that any such policy must  $\mu_t^o$ -a.e. agree with  $\phi_t^o$ .

<span id="page-39-2"></span>**Lemma B.5.** Given any probability measure  $\mu$  over [0, 1] such that  $\mu((\bar{p},1]) > 0$ , <sup>[51](#page-39-1)</sup> any threshold time-t policies satisfying  $p > \bar{p} \Rightarrow \phi_t(p) = 1$  ( $\mu$ -a.e.) and  $\int_p \phi_t(p) u(p) \mu(dp) = 0$ must agree  $\mu$ -a.e.

*Proof.* For  $i = 1, 2$ , let  $\phi_t^i$  be a threshold time-t policy with threshold  $\eta_t^i$ , which satisfies  $\int_p \phi_t^i(p)u(p)\mu(dp) = 0$  and  $p > \bar{p} \Rightarrow \phi_t^i(p) = 1$  ( $\mu$ -a.e.). Without loss of generality, assume  $\eta^1 \leq \eta^2$ . Notice under the assumption  $\mu((\bar{p},1]) > 0$ ,  $\int_p \phi_t^i(p)u(p)\mu(dp) = 0$  implies that we must have  $\phi_t^i(p) > 0$  for some p with  $u(p) < 0$ . Thus the threshold structure implies  $\phi_t^i(p) = 1$  for all p such that  $u(p) \geq 0$ , which holds for both  $i = 1, 2$ . Now, notice we have:

$$
0 = \int_p \phi_t^1(p)u(p)\mu(dp) - \int_p \phi_t^2(p)u(p)\mu(dp) = \int_{p:u(p)<0} (\phi_t^1(p) - \phi_t^2(p))u(p)\mu(dp)
$$

<span id="page-39-1"></span><span id="page-39-0"></span><sup>&</sup>lt;sup>50</sup>Notice  $\phi_t^d(p) = 1$  for any  $p > \bar{p}$  by the definition of  $\phi^d$  and Lemma [B.4\(](#page-35-1)a).

 $51$  $51$ Notice properties (P2) and (P5) in Lemma 1 together imply that this holds for time-t belief distribution  $\mu_t$  under any policy.

If  $\eta^1 < \eta^2$ , then  $\phi_t^1(p) \geq \phi_t^2(p)$  for all p because of the threshold structure. Supposing the policies do not agree  $\mu$ -a.e., which can only happen when  $u(p) < 0$ , then  $\phi_t^1(p) > \phi_t^2(p)$ for a positive  $\mu$ -measure set of p with  $u(p) < 0$ . This implies that the last expression above is strictly negative, which is a contradiction.

If  $\eta^1 = \eta^2 =: \eta$ , then the two policies can only differ at  $p = \eta$  with  $u(\eta) < 0$ . Supposing they do not agree  $\mu$ -a.e, we must have  $\mu({\{\eta\}}) > 0$  and  $\phi_t^1(\eta) \neq \phi_t^2(\eta)$ . These imply that the last expression above equals to  $[\phi_t^1(\eta) - \phi_t^2(\eta)]u(\eta)\mu(\{\eta\}) \neq 0$ , which is a contradiction. Thus the policies must agree  $\mu$ -a.e.  $Q.E.D.$ 

#### B.5.3 Proof for Proposition [3](#page-14-3) and Related Results

We need to first prove some properties of  $\phi^d$ .

<span id="page-40-0"></span>**Lemma B.6.** Set  $\{p : \phi_t^d(p) = 1\}$  shrinks in set inclusion order as t increases.

*Proof.* Since by construction  $\phi_t^d(p) = 1$  if and only if  $H_0(p, t) > 0$ , the result is directly implied by Lemma  $B.4(d)$  $B.4(d)$  in Section  $B.4$ .  $Q.E.D.$ 

This observation leads to the following result:

<span id="page-40-1"></span>**Lemma B.7.** Given any time-t belief distribution  $\mu_t$ , we have  $\int_p \phi_t^d(p) u(p) \mu_t(dp) \leq$  $\int_p \phi_{t+1}^d(p) u(p) \mu_{t+1}(dp)$ , where  $\mu_{t+1}$  is the period  $t+1$  belief distribution under  $\phi_t^d$  given  $\mu_t$ . (That is, the consumer's expected payoff is weakly higher in period  $t + 1$  than in period t under  $\phi^d$ .)

Proof. The result is proved by the following arguments:

$$
\int_{p'} \phi_t^d(p') u(p') \mu_t(dp')
$$
\n
$$
= \int_p \int_{p'} u(p') \phi_{t+1}^d(p') [\phi_t^d(p) G(dp'|p) + (1 - \phi_t^d(p)) D(dp'|p)] \mu_t(dp)
$$
\n
$$
= \int_p \int_{p'} u(p') \phi_{t+1}^d(p') G(dp'|p) \phi_t^d(p) \mu_t(dp) \ge \int_p u(p) \phi_t^d(p) \mu_t(dp)
$$

The first equality holds by the transition rule for  $p$ . The second equality holds because  $\phi_t^d(p) = 0 \Rightarrow \phi_{t+1}^d(p) = 0$  by Lemma [B.6,](#page-40-0) and thus  $\int_{p'} u(p') \phi_{t+1}^d(p') (1 - \phi_t^d(p)) D(dp'|p) =$  $u(p)\phi_{t+1}^d(p)(1-\phi_t^d(p))=0.$  The last inequality holds because  $\int_{p'} u(p') \phi_{t+1}^d(p') G(dp'|p) \geq \int_{p'} u(p') G(dp'|p) = u(p)$ , where the "≥" is due to  $u(p') > 0 \Rightarrow$  $\phi_{t+1}^d(p') = 1$  and the "=" is implied by property (P2) in Lemma [1.](#page-9-5)  $Q.E.D.$ 

To ease notation, let  $IC_t$  denote the IC constraint for time-t consumer. Repeated use of Lemma [B.7](#page-40-1) implies that given any  $\mu_t$ , if  $\phi_{\geq t}^d$  satisfies IC<sub>t</sub>, then it satisfies all later IC's. We are now ready to prove the proposition.

**Proof for Proposition [3](#page-14-3).** First consider the "only if" part. Let  $\phi^{opt}$  be any optimal policy for the designer and let  $\mu_t^{opt}$  be the distribution of  $p_t$  under it for any t. Due to Proposition [2,](#page-12-3) I can assume  $\phi^{opt}$  is a threshold policy without loss of generality. Let  $\lambda^*$ be a Lagrangian multiplier solving the dual problem that is non-increasing over  $t$  (which exists by Lemma [3\)](#page-11-0). I first check condition (i):

#### **Claim** (a).  $\phi^{opt}_{\hat{A}}$  $\langle \dot{t}$ agrees with  $\phi^o$  $\int_{0}^{\infty} \hat{\tau} \, ds$

*Proof for Claim (a).* I check by forward induction in t for  $t < \hat{t}$ . For  $t = 1, t < \hat{t}$  implies that  $\phi_1^d$  violates  $IC_1$  given the initial state distribution  $\mu_1$ . In this case, we must have  $\lambda_1^* > 0$  and thus  $IC_1$  is binding under  $\phi^{opt}$ . (Suppose not. Then  $\lambda_t^* = 0$  for all  $t \ge 1$  since it is non-increasing in t, and thus the Lagrangian problem  $\max \mathcal{L}(\phi; \lambda^*)$  would coincide with the dictator's problem. By Lemma [2,](#page-10-2) this implies that  $\phi^{opt}$  must also solve the dictator's problem. However, by construction  $\phi^d$  provides the highest time-1 expected consumer surplus given  $\mu_1$  among all optimal policies for the dictator's problem. Thus when  $\phi_1^d$  violates  $IC_1$ , so does  $\phi_1^{opt}$  $_{1}^{opt}$ , which is a contradiction.) Also notice the optimality of  $\phi^{opt}$  implies  $p > \bar{p} \Rightarrow \phi_1^{opt}$  $_1^{opt}(p) = 1$  ( $\mu_1$ -a.e.). Thus  $\phi_1^{opt}$  $_{1}^{\text{opt}}$  satisfies the properties of  $\phi_{1}^{\text{o}}$  in step 2 of the algorithm. By Lemma [B.5,](#page-39-2) we then must have  $\phi_1^{opt}$  $_{1}^{\text{opt}}$  and  $\phi_1^o$  agree  $\mu_1$ -a.e.

Now, assume  $\phi^{opt}$  and  $\phi^o$  agree a.s. for all periods before t and  $t < \hat{t}$ . Then, they induce the same distribution for  $p_t$ , which is just  $\mu_t^o$  constructed in the algorithm. I want to show  $\phi^{opt}$  must also satisfy  $p > \bar{p} \Rightarrow \phi_t^{opt}$  $t^{opt}_t(p) = 1 \mu_t^o$ -a.s. and  $\int_p \phi_t^{opt}$  $t_t^{opt}(p)u(p)\mu_t^o(dp)=0.$  The former is directly implied by the optimality of  $\phi^{opt}$ . For the latter, notice  $t < \hat{t}$  implies that  $\phi^d$  violates  $IC_t$  given  $p_t \sim \mu_t^o$ . We then must have  $\lambda_t^* > 0$  and thus  $IC_t$  is binding under  $\phi^{opt}$ . (Suppose not. Then  $\lambda^*_{t'} = 0$  for all  $t' \geq t$ , and thus the continuation Lagrangian problem starting with time t coincides with the corresponding dictator's continuation problem. The optimality of  $\phi^{opt}$  then implies that  $\phi^{opt}_{\geq t}$  must be optimal for this dictator's continuation problem given  $p_t \sim \mu_t^o$ . However, among all such policies,  $\phi_{\geq t}^d$  delivers the highest time-t expected consumer surplus. Thus when  $\phi_t^d$  violates  $IC_t$ , so does  $\phi_t^{opt}$  $t_t^{opt}$ , which is a contradiction.) Thus  $\int_p \phi_t^{opt}$  $t_t^{opt}(p)u(p)\mu_t^o(dp) = 0.$  Again by Lemma [B.5,](#page-39-2) we must have  $\phi_t^{opt}$  and  $\phi_t^o$  agree  $\mu_t^o$ -a.e. The proof is then completed by induction.  $\Box$ 

Now, since  $\phi^{opt}_{\hat{f}}$  $\lt t$ and  $\phi^o$  $\int_{\alpha}^{\infty}$  agree almost surely, they lead to the same distribution for  $p_{\hat{t}}$ , which is just  $\mu_{\hat{i}}^{\circ}$  $\hat{t}$ . Hence  $\phi_{\geq \hat{t}}^{opt}$  $\frac{\partial p_t}{\partial \hat{t}}$  must satisfy all IC constraints after time  $\hat{t}$  given  $p_{\hat{t}} \sim \mu_{\hat{t}}^d$  $\hat{\hat{t}}$ . The following claim checks the rest of condition (ii) in the proposition.

**Claim** (b).  $\phi^{opt}_{\leq \hat{t}}$  $\frac{\partial p_t}{\partial \hat{t}}$  is optimal for the dictator's continuation problem since time  $\hat{t}$  given  $p_{\widehat{t}} \sim \mu_{\widehat{t}}^o$  $\hat{\hat{t}}$ .

*Proof for Claim (b).* By the definition of  $\hat{t}$ ,  $\phi_t^d$ d satisfies  $IC_{\hat{t}}$  given  $p_{\hat{t}} \sim \mu_{\hat{t}}^{\omega}$  $\hat{\tau}$  and thus  $\phi^d_{\geq}$  $\geq t$ satisfies all later IC constraints by Lemma [B.7](#page-40-1) given  $p_{\hat{t}} \sim \mu_{\hat{t}}^o$  $\frac{1}{\hat{t}}$ . This implies that if we deviate from  $\phi^{opt}_{\leq \hat{i}}$  $\geq \widehat{t}$ to  $\phi^d$  $\frac{d}{dt}$  ince period  $\hat{t}$ , no IC constraint will be violated. Also notice such deviation can only improve the total surplus since  $\phi^d$ .  $\frac{d}{\geq \hat{t}}$  is optimal for the dictator's

continuation problem, which is more relaxed than the original continuation problem. For such deviation to be unprofitable, we then need  $\phi_{\mathcal{A}}^{opt}$  $\geq \hat{t}$ to achieve the same value as  $\phi^d$ .  $\sum_{i=1}^{d}$  for that continuation problem and thus  $\phi^{opt}_{\hat{\mathcal{F}}}$  $\geq t$ is also optimal for it.  $\Box$ 

Now, I turn to the "if" part. Given the existence of optimal policy (guaranteed by Proposition [1\)](#page-10-3), it suffices to see that all policies satisfying the two conditions (i) and (ii) are feasible and yield the same total payoff for the designer. For the total payoff, this is obviously true since: for periods before  $\hat{t}$ , all such policies agree a.s. and lead to  $p_{\hat{t}} \sim \mu_{\hat{t}}^o$ ; bt for periods  $t \geq \hat{t}$ , all such policies achieve the same total payoff as that under  $\phi_{\geq}^d$  $\sum_{i=1}^{d}$  given  $p_{\widehat{t}} \sim \mu_{\widehat{t}}^o$  $\hat{t}$ . For feasibility, it suffices to check  $\phi^o$ .  $\frac{\partial}{\partial \xi}$  is feasible. This is true since  $\phi^0$  $\frac{1}{n}$  satisfies all IC constraints for  $t < \hat{t}$  as equalities by construction. This completes the proof for the "if" part.

Finally, notice by construction  $\phi^o$  does satisfy conditions (i) and (ii). In particular, Lemma [B.7](#page-40-1) implies that  $\phi^d$ .  $\frac{d}{\geq \hat{t}}$  satisfies all IC constraints after time  $\hat{t}$  given  $p_{\hat{t}} \sim \mu_{\hat{t}}^{\omega}$  $\frac{\partial}{\partial t}$ , which implies the feasibility of  $\phi^o$  following time  $\hat{t}$ . Thus  $\phi^o$  is optimal.

 $Q.E.D.$ 

#### B.6 Proof for Proposition [4](#page-15-2)

*Proof.* Let  $\phi^*$  denote an optimal policy, let  $(\eta_t^*)_{t=1}^T$  denote its thresholds, and let  $\mu_t^*$ denote the distribution of  $p_t$  under it. Notice that under the full support condition in Assumption [2,](#page-14-4) Proposition [2](#page-12-3) implies that we must have  $\eta_t^* \leq \bar{p}$  for all t. (Recall that  $\bar{p}$  is the myopic threshold, i.e.,  $u(\bar{p}) = 0$ .) Moreover, under the atomless condition in Assumption [2,](#page-14-4) randomization at the thresholds does not matter.

For part (a), I first show the following observation:

**Claim**. For any  $\eta \in (0, \bar{p}]$ , we have  $\int_{p \geq \eta} \left[ \int_{p' \geq \eta} u(p') G(dp'|p) \right] \mu_t^*(dp) > \int_{p \geq \eta} u(p) \mu_t^*(dp)$ . *Proof for the claim.* Given any  $\eta \in (0, \bar{p}]$ , recall that property (P4) in Lemma [1](#page-9-5) implies  $G([0, \eta]|\eta) > 0$ . By the weak continuity of  $G(\cdot|p)$  on p (i.e., property (P[1\)](#page-9-5) in Lemma 1), this further implies that there exists  $\delta > 0$  s.t.  $G([0, \eta)|p) > 0 \,\forall p \in [\eta, \eta + \delta]$ .<sup>[52](#page-42-0)</sup> Together with the full support assumption on  $\mu_t^*$ , we then have  $\int_{p\geq \eta} G((0,\eta)|p)\mu_t^*(dp) > 0$ . Since  $\eta \leq \bar{p}$ ,  $u(p) < 0$  for all  $p < \eta$ . Thus  $\int_{p \geq \eta} \int_{p' < \eta} u(p') G(dp'|p) \mu_t^*(dp) < 0$ . This then implies  $\int_{p\geq \eta} \int_{p'\geq \eta} u(p') G(dp'|p) \mu_t^*(dp) > \int_{p\geq \eta} \int_{p'} u(p') G(dp'|p) \mu_t^*(dp) = \int_{p\geq \eta} u(p) \mu_t^*(dp)$ where the equality holds by property  $(P2)$  in Lemma [1.](#page-9-5)  $\Box$ 

<span id="page-42-0"></span> $52$ See Theorem 3.2.11 in [Durrett](#page-52-12) [\(2019\)](#page-52-12) (equivalence between conditions (i) and (ii)).

Now, I argue that the following holds given any  $t \le \hat{t} - 2$ :

$$
\int_{p \geq \eta_t^*} u(p) \mu_{t+1}^*(dp)
$$
\n
$$
= \int_p \int_{p' \geq \eta_t^*} u(p') \left[ \mathbb{1}_{\{p \geq \eta_t^*\}} G(dp'|p) + \mathbb{1}_{\{p < \eta_t^*\}} D(dp'|p) \right] \mu_t^*(dp)
$$
\n
$$
= \int_{p \geq \eta_t^*} \left[ \int_{p' \geq \eta_t^*} u(p') G(dp'|p) \right] \mu_t^*(dp) + \int_{p \geq \eta_t^*} \mathbb{1}_{\{p < \eta_t^*\}} u(p) \mu_t^*(dp)
$$
\n
$$
= \int_{p \geq \eta_t^*} \left[ \int_{p' \geq \eta_t^*} u(p') G(dp'|p) \right] \mu_t^*(dp)
$$
\n
$$
> \int_{p \geq \eta_t^*} u(p) \mu_t^*(dp) = 0
$$

The first equality holds by the transition rule of  $p_t$ ; the second equality is trivial identity; the third equality holds because the second term in line 3 is obviously zero; the last expression equals to zero because the IC constraint is binding for any  $t < \hat{t}$  by Proposi-tion [3.](#page-14-3) To see the inequality holds, notice that  $t \leq \hat{t} - 2$  necessarily implies  $\eta_t^* > 0$ , since otherwise  $\phi_t^d$  would be feasible at time t and the algorithm in Definition [2](#page-13-2) would have stopped in step 1 at time  $t$ . The desired inequality is then directly implied by the claim proved above.

For part (b), notice under the atomless assumption in Assumption [2,](#page-14-4) Lemma  $B.4(c)$  $B.4(c)$ (see Appendix [B.4\)](#page-30-3) implies that  $H_0(p_t, t)$  is non-zero almost surely under any policy. Lemma [B.2\(](#page-31-2)b) then implies that any optimal policy for the dictator must almost surely agree with  $\phi^d$ . By Proposition [3,](#page-14-3) this further implies that  $\phi^*_{\geq \hat{t}}$  must almost surely agree with  $\phi^d$  $\frac{d}{\geq \hat{t}}$  (given  $p_t \sim \mu_{\hat{t}}^*$  $\hat{t}$ ). Under the full support assumption in Assumption [2,](#page-14-4) this then requires that  $\phi^*$  $\sum_{i=1}^{n}$  and  $\phi_{\geq}^{d}$  $\frac{d}{dt}$  share the same thresholds. It thus suffices to prove the desired property for  $\phi^d$ .

Let  $(\eta_t^d)_{t=1}^T$  denote the sequence of thresholds of  $\phi^d$ . Recall that  $\phi_t^d(p) = \mathbb{1}_{\{H_0(p,t)>0\}}$ (as is defined in Appendix [B.5.1\)](#page-38-2). By the continuity of  $H_0(\cdot, t)$  (Lemma [B.1\)](#page-30-0), we then have  $H_0(\eta_t^d, t) = 0$  for any  $t^{53}$  $t^{53}$  $t^{53}$  By Lemma [B.4\(](#page-35-1)d), this further implies  $H_0(\eta_t^d, t + 1) < 0$ . Since  $H_0(\cdot, t+1)$  is increasing (Lemma [B.4\(](#page-35-1)b)), we thus must have  $\eta_{t+1}^d > \eta_t^d$  for any t. This completes the proof for part (b).

Finally,  $\eta_t^* \leq \bar{p}$  for all t is directly implied by Proposition [2](#page-12-3) under the full support assumption in Assumption [2.](#page-14-4)

$$
Q.E.D.
$$

<span id="page-43-0"></span><sup>&</sup>lt;sup>53</sup>Notice it is easy to see that  $H_0(0, t) < 0$  and  $H_0(1, t) > 0$ .

#### <span id="page-44-0"></span>B.7 Proof for Proposition [5](#page-19-2)

Proof. The designer's problem is formally written as:

$$
\max_{\phi \in \Phi} \left\{ \sum_{t=1}^{T} \mathbb{E}_{\phi}[a_t u(p_t)] \right\}
$$
  
s.t. 
$$
\mathbb{E}_{\phi}[a_t u(p_t)] \ge 0 \quad \forall t = 1, ..., T
$$

$$
p_{t+1}[p_t, a_t \sim a_t [\alpha G^I(\cdot | p_t) + (1 - \alpha)D(\cdot | p_t)] + (1 - a_t)D(\cdot | p_t)
$$

$$
p_1 \sim \mu_1
$$

Pick  $\alpha_a$  and  $\alpha_b$  with  $\alpha_a < \alpha_b$ . Corresponding to these two information generation rates respectively, let  $G^a$  and  $G^b$  be the transition kernels of  $p_t$  following one's consumption, as is defined in equation [\(9\)](#page-18-5); let  $V_0^a$  and  $V_0^b$  be the value functions for the dictator's problem (i.e., with  $\lambda = 0$ ), as is defined in Section [B.4;](#page-30-3) let  $H_0^a$  and  $H_0^b$  be the associated H-functions (with  $\lambda = 0$ ) as in equation [\(B.8\)](#page-30-2); let  $\hat{t}^a$  and  $\hat{t}^b$  denote the critical time points defined in Definition [2;](#page-13-2) let  $\phi^a$  and  $\phi^b$  be the optimal threshold policies and denote their sequences of thresholds as  $(\eta_t^a)_{t=1}^T$  and  $(\eta_t^b)_{t=1}^T$ . I show  $\eta_t^a \geq \eta_t^b$  for all t with a sequence of claims below.

First, since higher  $\alpha$  is beneficial, we have the following non-surprising result for  $V_0^a$ and  $V_0^b$ :

**Claim** (a).  $V_0^b(p,t) \geq V_0^a(p,t)$  for any pair of  $(p,t)$ .

*Proof for Claim (a).* I show by backward induction on t. For  $t = T + 1$ , the result holds trivially. Assuming it holds for all  $t' > t$ , I now consider time t.

By the Bellman equation, we have:

$$
V_0^b(p,t) - V_0^a(p,t) = \max \left\{ u(p) + \int_{p'} V_0^b(p', t+1) G^b(dp'|p), V_0^b(p, t+1) \right\}
$$

$$
- \max \left\{ u(p) + \int_{p'} V_0^a(p', t+1) G^a(dp'|p), V_0^a(p, t+1) \right\}
$$

By the induction hypothesis, we know  $V_0^b(p, t+1) \geq V_0^a(p, t+1)$ . It thus suffices to check

 $\int_{p'} V_0^b(p', t+1) G^b(dp'|p) \geq \int_{p'} V_0^a(p', t+1) G^a(dp'|p)$ . Notice the following relations hold:

$$
\int_{p'} V_0^b(p', t+1) G^b(dp'|p) - \int_{p'} V_0^a(p', t+1) G^a(dp'|p)
$$
  
\n
$$
= \alpha_b \int_{p'} V_0^b(p', t+1) G^I(dp'|p) + (1 - \alpha_b) V_0^b(p, t+1)
$$
  
\n
$$
- \alpha_a \int_{p'} V_0^a(p', t+1) G^I(dp'|p) - (1 - \alpha_a) V_0^a(p, t+1)
$$
  
\n
$$
\geq \alpha_a \int_{p'} V_0^b(p', t+1) G^I(dp'|p) + (1 - \alpha_a) V_0^b(p, t+1)
$$
  
\n
$$
- \alpha_a \int_{p'} V_0^a(p', t+1) G^I(dp'|p) - (1 - \alpha_a) V_0^a(p, t+1)
$$

where the equality is by the definition of  $G^a$  and  $G^b$ . To see the inequality holds, recall that Lemma [B.3\(](#page-32-0)b) implies  $\int_{p'} V_0^b(p', t+1) G^b(dp'|p) \geq V_0^b(p, t+1)$ , which further implies  $\int_{p'} V_0^b(p', t+1) G^I(dp'|p) \geq V_0^b(p, t+1)$  since  $G^b(\cdot|p)$  is a weighted average of  $G^I(\cdot|p)$ and  $D(\cdot|p)$ . The above inequality thus holds given  $\alpha_b > \alpha_a$ . Now, notice the induction hypothesis implies that the last expression above is indeed non-negative. We thus have the desired result. □

An important implication of the above claim is:

**Claim** (b).  $H_0^b(p,t) \leq 0 \Rightarrow H_0^a(p,t) \leq 0$  for all pairs of  $(p,t)$ .

*Proof for Claim (b)*. When  $H_0^b(p, t) \leq 0$ , by Lemma [B.4\(](#page-35-1)d) we know  $H_0^b(p, t') \leq 0$  for all  $t' > t$ . With  $\alpha = \alpha_b$ , it is thus optimal for the dictator to stop recommendation from time t on given  $p_t = p$ , which leads to the optimal continuation value being zero. Hence  $V_0^b(p,t) = 0$ . By Claim (a), this implies  $V_0^a(p,t) \leq 0$  and it is thus also optimal for the dictator to stop recommendation at  $(p, t)$  given  $\alpha = \alpha_a$ . This then implies  $H_0^a(p, t) \leq$  $0.$ 

With Claim (b), we can now prove the desired result for  $t \geq \hat{t}^b$ .

*Claim* (*c*).  $\eta_t^a \geq \eta_t^b$  for all  $t \geq \hat{t}^b$ .

*Proof for Claim (c)*. By Proposition [3,](#page-14-3) we know that  $\phi^b$ .  $\sum_{i=1}^{b} b_i$  is optimal for the dictator's continuation problem with  $\alpha = \alpha_b$  given the distribution of  $p_{\hat{t}b}$  under  $\phi^b$ . Under As-sumption [2,](#page-14-4) this together with Lemma [B.2\(](#page-31-2)b) implies that  $H_0^b(p, t) \leq 0$  for Lebesgue-a.e.  $p < \eta_t^b$  for all  $t \geq \hat{t}^b$ .

Now, suppose  $t \geq \hat{t}^b$  but  $\eta_t^a < \eta_t^b$ . Then the above conclusion together with Claim (b) implies that  $H_0^a(p,t) \leq 0$  for Lebesgue-a.e.  $p \in (\eta_t^a, \eta_t^b)$ . By Lemma [B.4,](#page-35-1) we know  $H_0^a(\cdot, t)$ is weakly increasing and has at most one zero point. Thus we must have  $H_0^a(p, t) < 0$  for all  $p \in (\eta_t^a, \eta_t^b)$ .

The above result further implies that given  $\alpha = \alpha_a$ , the most conservative optimal policy for the dictator (i.e.,  $\phi^d$  as is defined in Section [3.4\)](#page-13-1) has a threshold larger than  $\eta_t^a$  at time  $t$  and it is hence incentive compatible at time  $t$  provided that we have been following  $\phi^a$  before. By the algorithm in Definition [2](#page-13-2) and Proposition [3,](#page-14-3) we then must have  $t \geq \hat{t}^a$ . This in turn implies the optimality of  $\phi_{\geq t}^a$  for the dictator's continuation problem given  $\alpha = \alpha_a$  and hence under Assumption [2](#page-14-4) we must have  $H_0^a(p, t) < 0 \Rightarrow p < \eta_t^a$  (Lebesgue-a.e. ) by Lemma [B.2\(](#page-31-2)b). This contradicts with the result  $H_0^a(p, t) < 0$  for all  $p \in (\eta_t^a, \eta_t^b)$ above. Thus we must have  $\eta_t^a \geq \eta_t^b$  when  $t \geq \hat{t}^b$ . □

Now it suffices to show  $\eta_t^a \geq \eta_t^b$  for  $t < \hat{t}^b$ . For  $k = a, b$ , let  $F_t^k$  denote the cdf of  $p_t$  under  $\phi^k$  for any t given  $\alpha = \alpha_k$ . Under Assumption [2,](#page-14-4) notice that  $F_t^a$  and  $F_t^b$  are continuous and have full support over  $[0, 1]$  for any t. The following observation is the key part of the proof.

**Claim** (d). Let  $\tau$  be any fixed time. If  $\eta_t^a \geq \eta_t^b$  for all  $t < \tau$ , then  $F_\tau^b$  is a mean-preserving spread of  $F_{\tau}^a$  and  $F_{\tau}^a(p) \leq F_{\tau}^b(p)$  for any  $p \leq \min_{t \leq \tau} \{\eta_t^b\}.$ 

Proof for Claim (d). To prove the claim, I construct two belief processes (truncated at time  $\tau$ )  $(p_t^a)_{t=1}^{\tau}$  and  $(p_t^b)_{t=1}^{\tau}$  on the same probability space, where  $(p_t^a)_{t=1}^{\tau}$  follows the transition rule decided by  $\phi^a$  given  $\alpha_a$  and  $(p_t^b)_{t=1}^\tau$  follows the transition rule decided by  $\phi^b$  given  $\alpha_b$ .

Specifically, fix a probability space on which a Markov process  $(x_n)_{n=0}^{\infty}$  and a sequence of i.i.d. random variables  $(\xi_t)_{t=1}^{\tau}$  with  $\xi_t \sim$  Uniform[0, 1] are defined.  $(x_n)_{n=0}^{\infty}$  is independent from  $(\xi_t)_{t=1}^{\tau}$  and satisfies:

$$
x_0 \sim \mu_1
$$

$$
x_{n+1}|x_n \sim G^I(\cdot | x_n), \ \forall n
$$

Intuitively, one can interpret  $x_n$  as the value that the platform's belief will take after receiving the n'th informative signal from consumers;  $\xi_t$  will serve as a randomization device deciding whether an informative signal will be generated after consumption at time t. I define a filtration of  $\sigma$ -fields  $(\mathcal{F}_n)_{n=0}^{\infty}$  such that  $\mathcal{F}_n = \sigma((\xi_t)_{t=1}^{\tau}, x_0, ..., x_n)$  for all *n*. Then obviously  $(x_n)_{n=0}^{\infty}$  is a martingale w.r.t.  $(\mathcal{F}_n)_{n=0}^{\infty}$ .

For  $k = a, b$ , I now define process  $(p_t^k)_{t=1}^{\tau}$  together with an auxiliary process  $(n_t^k)_{t=1}^{\tau}$ by the following rule:

$$
n_1^k = 0; \quad p_1^k = x_0
$$

$$
n_{t+1}^k = n_t^k + \mathbb{1}_{\{p_t^k > \eta_t^k\}} \mathbb{1}_{\{\xi_t < \alpha_k\}}; \quad p_{t+1}^k = x_{n_{t+1}^k} \quad \forall t
$$

Intuitively, under the scenario with  $(\alpha_k, \phi^k)$ ,  $n_t^k$  tracks how many informative signals have been recorded at the beginning of time t. It is added by 1 after each period if and only if consumption has been made in that period (i.e.,  $p_t^k > \eta_t^k$ ) and an informative signal is generated (which is assumed to happen when  $\xi_t < \alpha^k$ ).<sup>[54](#page-47-0)</sup> Given that  $n_t^k$  informative signals have been received,  $p_t^k$  just equals to  $x_{n_t^k}$ , which reflects the posterior belief given those signals. It is easy to check that  $(p_t^k)_{t=1}^{\tau}$  indeed satisfy the initial distribution and the transition rule of the belief process under policy  $\phi^k$  given response rate  $\alpha_k$  and thus  $p_t^k \sim F_t^k$  for all  $t \leq \tau$ .

Now, I notice that  $n_{\tau}^{k}$  is a bounded stopping time w.r.t.  $(\mathcal{F}_{n})_{n=0}^{\infty}$ . The boundedness is obvious since  $n_{\tau}^k < \tau$ . To show it is a stopping time, notice by the construction of  $(n_t^k)_{t=1}^{\tau}$ , whether  $\{n_{\tau}^k \leq n\}$  happens is solely determined by  $(x_0, ..., x_n)$  together with  $(\xi_t)_{t=1}^{\tau}$ . (With  $(x_0, ..., x_n)$  and  $(\xi_t)_{t=1}^{\tau}$ , we can perfectly predict when the  $(n + 1)$ 'th informative signal will come.) Thus  $\{n_{\tau}^{k} \leq n\} \in \mathcal{F}_n$  and  $n_{\tau}^{k}$  is a stopping time w.r.t.  $(\mathcal{F}_n)_{n=0}^{\infty}$  by definition. Moreover, since  $\eta_t^a \geq \eta_t^b$  for all  $t < \tau$  and  $\alpha_b > \alpha_a$ , it is easy to see that  $n_\tau^b \geq n_\tau^a$ for sure.<sup>[55](#page-47-1)</sup> By the Doob's optional sampling theorem, we then have  $\mathbb{E}[x_{n_{\tau}^{b}}|\mathcal{F}_{n_{\tau}^{a}}] = x_{n_{\tau}^{a}}$ and thus  $\mathbb{E}[x_{n_{\tau}^{b}}|x_{n_{\tau}^{a}}] = x_{n_{\tau}^{a}}$ .<sup>[56](#page-47-2)</sup> This further implies  $\mathbb{E}[p_{\tau}^{b}|p_{\tau}^{a}] = p_{\tau}^{a}$  and therefore  $F_{\tau}^{b}$  is a mean-preserving spread of  $F_{\tau}^a$ .

Now, pick any  $p \leq \min_{t \leq \tau} {\{\eta_t^b\}}$ . Notice that because  $p \leq \eta_t^b \leq \eta_t^a$  for all  $t < \tau$ , by construction the processes of  $(p_t^a)_{t=1}^{\tau}$  and  $(p_t^b)_{t=1}^{\tau}$  will stop once they fall into  $[0, p]$ . Thus when  $p_{\tau}^{b} > p$ , we must have  $x_n > p$  for all  $n \leq n_{\tau}^{b}$ . Because  $n_{\tau}^{a} \leq n_{\tau}^{b}$ , we then must have  $p_{\tau}^{a} = x_{n_{\tau}^{a}} > p$ . This implies  $p_{\tau}^{b} > p \Rightarrow p_{\tau}^{a} > p$  for sure and therefore  $F_{\tau}^{a}(p) \leq F_{\tau}^{b}(p)$ .

□

We are now ready to complete the last piece of the proof:

*Claim* (e).  $\eta_t^a \geq \eta_t^b$  for all  $t < \hat{t}^b$ .

*Proof for Claim (e)*. It suffices to assume  $\hat{t}^b > 1$ , otherwise the result is vacuous. I prove the claim by induction in  $t = 1, ..., \hat{t}^b$ . For  $t = 1, t < \hat{t}^b$  implies that  $\int_{p \ge \eta_1^b} u(p) \mu_1(dp) = 0$ (by Proposition [3\)](#page-14-3). Feasibility of  $\phi^a$  also implies  $\int_{p \ge \eta_1^a} u(p) \mu_1(dp) \ge 0$ . Since  $\eta_1^a, \eta_1^b \le \bar{p}$ by the optimality of  $\phi^a$  and  $\phi^b$ , these obviously imply  $\eta_1^a \leq \eta_1^b$  under Assumption [2.](#page-14-4)

Now, assuming the result holds for all periods  $t < \tau$ , I show it for period  $\tau$ . First, notice  $\tau < \hat{t}^b$  implies that  $\eta_t^b$  has been decreasing over time up to time  $\tau$  by Proposition [4.](#page-15-2) Thus if  $\eta_{\tau}^a > \eta_t^b$  for some  $t < \tau$ , we must have  $\eta_{\tau}^a > \eta_{\tau}^b$  to hold. Therefore, it suffices to

<span id="page-47-0"></span><sup>&</sup>lt;sup>54</sup>Under Assumption [2,](#page-14-4) what happens when  $p_t^k = \eta_t^k$  does not matter since it has zero probability to occur.

<span id="page-47-1"></span><sup>&</sup>lt;sup>55</sup>By construction, whenever  $n_t^b = n_t^a$ , we have  $p_t^b = p_t^a$  and hence  $n_{t+1}^a = n_t^a + 1 \Rightarrow n_{t+1}^b = n_t^b + 1$ . Thus the sequence of  $(n_t^a)_t$  can never surpass  $(n_t^b)_t$ .

<span id="page-47-2"></span> $56$ See, for example, theorem 10.11 in [Klenke](#page-52-13) [\(2020\)](#page-52-13) (third edition).

consider the case where  $\eta_{\tau}^a \leq \min_{t \leq \tau} \eta_t^b$ . In this case, I argue that the following holds:

$$
\int_{\eta_{\tau}^{a}}^{1} u(p) dF_{\tau}^{b}(p) = u(1) - u(\eta_{\tau}^{a}) F_{\tau}^{b}(\eta_{\tau}^{a}) - (u_{H} - u_{L}) \int_{\eta_{\tau}^{a}}^{1} F_{\tau}^{b}(p) dp
$$
  
\n
$$
\geq u(1) - u(\eta_{\tau}^{a}) F_{\tau}^{a}(\eta_{\tau}^{a}) - (u_{H} - u_{L}) \int_{\eta_{\tau}^{a}}^{1} F_{\tau}^{a}(p) dp
$$
  
\n
$$
= \int_{\eta_{\tau}^{a}}^{1} u(p) dF_{\tau}^{a}(p) \geq 0
$$

where the two equalities are just by integration by parts,  $57$  and the last inequality is due to the incentive compatibility of  $\phi^a$ . To check the first inequality, notice under the induction hypothesis and the assumption that  $\eta_{\tau}^a \leq \min_{t \leq \tau} \eta_t^b$ , Claim (d) implies  $F_{\tau}^{b}(\eta_{\tau}^{a}) \geq F_{\tau}^{a}(\eta_{\tau}^{a})$  and  $F_{\tau}^{b}$  is a mean-preserving spread of  $F_{\tau}^{a}$ , the latter of which further implies  $\int_{\eta_{\tau}^a}^1 F_{\tau}^b(p) dp \leq \int_{\eta_{\tau}^a}^1 F_{\tau}^a(p) dp$ .<sup>[58](#page-48-1)</sup> Together with the fact that  $u(\eta_{\tau}^a) \leq 0$ ,<sup>[59](#page-48-2)</sup> these imply the desired inequality.

The above discussion has shown  $\int_{\eta_{\tau}}^1 u(p) dF_{\tau}^b(p) \geq 0$ . Since  $\tau < \hat{t}^b$ , by Proposition [3](#page-14-3) we also know  $\int_{\eta_{\tau}^b}^1 u(p) dF_{\tau}^b(p) = 0$ . Thus we must have  $\eta_{\tau}^b \leq \eta_{\tau}^a$  under Assumption [2.](#page-14-4) This completes the proof by induction.

□

 $Q.E.D.$ 

#### B.8 Proof for Proposition [6](#page-20-2)

Proof. Based on my discussion in the main text, the designer's problem is equivalent to:

$$
\max_{\phi \in \Phi} \left\{ \sum_{t=1}^{T} \mathbb{E}_{\phi}[a_t u(p_t)] \right\}
$$
\n  
\n
$$
\text{s.t. } \mathbb{E}_{\phi}[a_t u(p_t)] \geq 0 \quad \forall t = 1, ..., T
$$
\n  
\n
$$
p_{t+1}|p_t, a_t \sim a_t \left[ \rho G(\cdot | p_t) + (1 - \rho) D(\cdot | p_t) \right] + (1 - a_t) D(\cdot | p_t)
$$
\n  
\n
$$
p_1 \sim \mu_1
$$

With  $\rho$  replaced by  $\alpha$  and G replaced by  $G<sup>I</sup>$ , this is equivalent to the designer's problem studied in Appendix [B.7.](#page-44-0) Thus the effect of an increment in  $\rho$  here is equivalent to the

<span id="page-48-0"></span><sup>&</sup>lt;sup>57</sup>Notice  $u(p) = u_L + (u_H - u_L)p$  in the current application, and  $F^a_\tau$  and  $F^b_\tau$  are continuous under Assumption [2.](#page-14-4)

<span id="page-48-1"></span><sup>&</sup>lt;sup>58</sup>See, e.g., Theorem 3.A.1(a) in [Shaked & Shanthikumar](#page-53-14) [\(2007\)](#page-53-14). In their notation, we have  $F^b_\tau \geq_{cx} F^a_\tau$ . The theorem implies  $\int_{\eta_{\tau}^a}^{\infty} [1-F_{\tau}^b(p)]dp \geq \int_{\eta_{\tau}^a}^{\infty} [1-F_{\tau}^a(p)]dp$ , which implies my result since  $F_{\tau}^a(p) = F_{\tau}^b(p)$ 1 for  $p > 1$ .

<span id="page-48-2"></span><sup>&</sup>lt;sup>59</sup>Otherwise, there would be a non-empty interval of p in which  $p > \bar{p}$  but  $p < \eta_{\tau}^a$ . This would violate the optimality of  $\phi^a$  under Assumption [2.](#page-14-4)

effect of an increment in  $\alpha$  there. The result is hence directly implied by Proposition [5.](#page-19-2)  $Q.E.D.$ 

#### B.9 Proof for Proposition [A.1](#page-25-4)

In the following proof, I assume  ${Q_z}_{z \in Z}$  and the conditional distributions of  $s_i$   $(i \geq 1)$ conditional on  $\tilde{\theta}$  are all continuous distributions, so the dominating measure for their densities is chosen as the Lebesgue measure. In the general case, the proof remains the same with Lebesgue measure replaced by proper dominating measures on  $\mathbb{R}$  (e.g., counting measure for discrete distributions).

*Proof.* For any  $z \in Z$  and  $s \in S$ , I define  $\psi_{(z,s)}$  as a probability density over R such that

$$
\psi_{(z,s)}(\theta) = \frac{q_z(\theta)\ell(s|\theta)}{\int q_z(\theta)\ell(s|\theta)d\theta}
$$

That is,  $\psi_{(z,s)}$  is the density function of the posterior about  $\tilde{\theta}$  computed from Bayes rule given prior  $Q_z$  and consumption-generated signal realization s.

**Claim** (a). For any  $x, y \in Z$  and  $s_a, s_b \in S$ , we have  $Q_y \geq_{LR} Q_x$  and  $s_b \geq s_a$  together imply  $\psi_{(y,s_b)} \geq_{LR} \psi_{(x,s_a)}$ . *Proof for Claim (a).* Assume  $Q_y \geq_{LR} Q_x$  and  $s_b \geq s_a$ . By the definition of  $\psi$ , we have:

$$
\frac{\psi_{(y,s_b)}(\theta)}{\psi_{(x,s_a)}(\theta)} = \frac{q_y(\theta)}{q_x(\theta)} \cdot \frac{\ell(s_b|\theta)}{\ell(s_a|\theta)}
$$

Since  $Q_y \geq_{LR} Q_x$  and  $\ell(\cdot|\theta)$  increases in likelihood-ratio order in  $\theta$ , both fractions on the right-hand-side are increasing in  $\theta$ . Thus  $\psi_{(y,s_b)} \geq_{LR} \psi_{(x,s_a)}$ . . □

Now, I show the following observation:

**Claim** (b). Assume  $\lambda_t$  is non-increasing over t. Then, for any  $x, y \in Z$ , we have  $Q_y \geq_{LR}$  $Q_x \Rightarrow H_\lambda(y,t) \geq H_\lambda(x,t)$  for all t.

*Proof for Claim (b).* I show by backward induction in t. The result holds with  $t = T$ since  $H_{\lambda}(z,T) = (1 + \lambda_T) \int \theta dQ_z(\theta)$ . Now, assuming the result holds for all periods since time  $t + 1$ , I show it for period t. Recall that equation [\(B.14\)](#page-36-0) derived in Section [B.4](#page-30-3) implies

$$
H_{\lambda}(z,t) = (\lambda_{t} - \lambda_{t+1})u(z) + \min\{H_{\lambda}(z,t+1),0\}
$$
  
+ 
$$
\int_{z' \in Z} \max\{H_{\lambda}(z',t+1),0\} [\rho G(dz';z) + (1-\rho)D(dz';z)]
$$
  
= 
$$
(\lambda_{t} - \lambda_{t+1})u(z) + \rho \min\{H_{\lambda}(z,t+1),0\} + (1-\rho)H_{\lambda}(z,t+1)
$$
  
+ 
$$
\rho \int_{z' \in Z} \max\{H_{\lambda}(z',t+1),0\} G(dz';z)
$$

(Since we have random consumer arrivals with arrival rate  $\rho$ , the transition kernel G in equation [\(B.14\)](#page-36-0) is replaced with  $\rho G + (1 - \rho)D$ .)

Pick any  $x, y \in Z$  s.t.  $Q_y \geq_{LR} Q_x$ . We obviously have  $(\lambda_t - \lambda_{t+1})u(y) \geq (\lambda_t - \lambda_{t+1})u(x)$ given the assumption that  $\lambda_t$  is non-increasing in t. Moreover, the induction hypothesis implies min $\{H_{\lambda}(y,t+1), 0\} \ge \min\{H_{\lambda}(x,t+1), 0\}$  and  $(1-\rho)H_{\lambda}(y,t+1) \ge (1-\rho)H_{\lambda}(x,t+1)$ 1). It then suffices to show  $\int_{z' \in Z} \max\{H_\lambda(z', t+1), 0\} G(dz'; y) \ge \int_{z' \in Z} \max\{H_\lambda(z', t+1), 0\}$ 1),  $0$ } $G(dz';x)$  below.

Notice in the current setting, state z matters only through the belief it represents. With slight abuse of notation, I write  $H_{\lambda}(q_z, t + 1) = H_{\lambda}(z, t + 1)$ . Then, we have:

$$
\int_{z' \in Z} \max\{H_{\lambda}(z', t+1), 0\} G(dz'; y) = \int_{\theta} \Big[ \int_{s} \max\{H_{\lambda}(\psi_{(y,s)}, t+1), 0\} \ell(s|\theta) ds \Big] q_{y}(\theta) d\theta
$$
\n
$$
\geq \int_{\theta} \Big[ \int_{s} \max\{H_{\lambda}(\psi_{(x,s)}, t+1), 0\} \ell(s|\theta) ds \Big] q_{y}(\theta) d\theta
$$
\n
$$
\geq \int_{\theta} \Big[ \int_{s} \max\{H_{\lambda}(\psi_{(x,s)}, t+1), 0\} \ell(s|\theta) ds \Big] q_{x}(\theta) d\theta
$$
\n
$$
= \int_{z' \in Z} \max\{H_{\lambda}(z', t+1), 0\} G(dz'; x)
$$

where the two equalities hold by the definition of  $\psi_{(z,s)}$ . The first inequality holds due to the induction hypothesis and that Claim (a) above implies  $\psi_{(y,s)} \geq_{LR} \psi_{(x,s)}$ . To see the second inequality, notice Claim (a) implies that  $\psi_{(x,s)}$  increases in likelihood-ratio order in s. Together with the induction hypothesis, this implies that  $\max\{H_{\lambda}(\psi_{(x,s)},t+1),0\}$ increases in s, which further implies that  $\int_s \max\{H_\lambda(\psi_{(x,s)}, t+1), 0\} \ell(s|\theta) ds$  increases in θ since ℓ(·|θ) increases in likelihood-ratio order in θ. The inequality is hence implied by  $q_y \geq_{LR} q_x.$ 

Now, I slightly strengthen both the condition and the conclusion in Claim (b).

**Claim** (c). Assume  $\lambda_t$  is non-increasing over t. Then, for any  $x, y \in Z$ , we have  $Q_y \geq_{LR}$  $Q_x$  and  $\int_{\theta} \theta dQ_y(\theta) > \int_{\theta} \theta dQ_x(\theta)$  together imply  $H_{\lambda}(y, t) > H_{\lambda}(x, t)$  for all t. *Proof for Claim (c).* I show by backward induction in t. The result holds with  $t = T$ since  $H_{\lambda}(z,T) = (1 + \lambda_T) \int \theta dQ_z(\theta)$ . Now, assuming the result holds for all periods since time  $t + 1$ , I show it for period t. By the same argument as in the proof of Claim (b), we

have

$$
H_{\lambda}(z,t) = (\lambda_t - \lambda_{t+1})u(z) + \rho \min\{H_{\lambda}(z,t+1),0\} + (1-\rho)H_{\lambda}(z,t+1) + \rho \int_{z' \in Z} \max\{H_{\lambda}(z',t+1),0\}G(dz';z)
$$

and that for any x and y satisfying the conditions in the claim: (i)  $(\lambda_t - \lambda_{t+1})u(y) \ge$  $(\lambda_t - \lambda_{t+1})u(x)$ ; (ii)  $\min\{H_\lambda(y, t+1), 0\} \ge \min\{H_\lambda(x, t+1), 0\}$ ; (iii)  $\int_{z' \in Z} \max\{H_\lambda(z', t+1)\}$ 

1), 0} $G(dz'; y) \ge \int_{z' \in Z} \max\{H_\lambda(z', t+1), 0\} G(dz'; x)$ . Moreover, the induction hypothesis directly imply that  $(1 - \rho)H_{\lambda}(y, t + 1) > (1 - \rho)H_{\lambda}(x, t + 1)$  for  $\rho < 1$ . These together imply  $H_{\lambda}(y, t) > H_{\lambda}(x, t)$  as is desired.

Now, define functions  $V_{\lambda}$  and  $H_{\lambda}$  in the same way as in Appendix [B.4,](#page-30-3) but with  $p_t$ replaced with  $z_t$ . Then the dynamic programming result – Lemma  $B.2$  – still applies to the current setting, because its proof only relies on property (P1) in Lemma [1,](#page-9-5) which has its counterpart in Lemma [A.1.](#page-24-2) Given the result of Claim  $(c)$  and Lemma [A.2,](#page-24-3) Lemma [B.2\(](#page-31-2)b) implies that any solution to the Lagrangian optimization  $\max_{\phi} \mathcal{L}(\phi; \lambda^*)$  (with  $\lambda^*$  solves the dual problem) is almost surely equivalent to some  $\phi^*$  satisfying the property specified in Proposition [A.1.](#page-25-4) The proposition hence holds by the duality result in Lemma [2.](#page-10-2)

Q.E.D.

### References

- <span id="page-51-0"></span>Altman, E. (1999). Constrained markov decision processes: stochastic modeling. Routledge.
- <span id="page-51-2"></span>Bahar, G., Smorodinsky, R., & Tennenholtz, M. (2015). Economic recommendation systems. arXiv preprint arXiv:1507.07191 .
- <span id="page-51-3"></span>Bahar, G., Smorodinsky, R., & Tennenholtz, M. (2021). Recommendation systems and self motivated users. arXiv preprint arXiv:1807.01732.
- <span id="page-51-1"></span>Ball, I. (2019). Dynamic information provision: Rewarding the past and guiding the future. Available at SSRN 3103127.
- Banerjee, A. V. (1992). A simple model of herd behavior. The quarterly journal of economics,  $107(3)$ , 797–817.
- <span id="page-51-4"></span>Beutler, F. J., & Ross, K. W. (1985). Optimal policies for controlled markov chains with a constraint. Journal of mathematical analysis and applications,  $112(1)$ ,  $236-252$ .
- <span id="page-51-5"></span>Beutler, F. J., & Ross, K. W. (1986). Time-average optimal constrained semi-markov decision processes. Advances in Applied Probability, 18(2), 341–359.
- Bikhchandani, S., Hirshleifer, D., & Welch, I. (1992). A theory of fads, fashion, custom, and cultural change as informational cascades. Journal of political Economy,  $100(5)$ , 992–1026.
- Bolton, P., & Harris, C. (1999). Strategic experimentation. *Econometrica*,  $67(2)$ , 349– 374.
- <span id="page-52-1"></span>Che, Y.-K.,  $\&$  Hörner, J. (2018). Recommender systems as mechanisms for social learning. The Quarterly Journal of Economics, 133 (2), 871–925.
- <span id="page-52-8"></span>Chen, B., Frazier, P., & Kempe, D. (2018). Incentivizing exploration by heterogeneous users. In Conference on learning theory (pp. 798–818).
- <span id="page-52-12"></span>Durrett, R. (2019). *Probability: theory and examples* (Vol. 49). Cambridge university press.
- <span id="page-52-6"></span>Ely, J. C. (2017). Beeps. American Economic Review, 107(1), 31–53.
- <span id="page-52-7"></span>Ely, J. C., & Szydlowski, M. (2020). Moving the goalposts. Journal of Political Economy,  $128(2)$ , 468–506.
- <span id="page-52-11"></span>Feinberg, E. A., & Piunovskiy, A. B. (2000). Multiple objective nonatomic markov decision processes with total reward criteria. Journal of mathematical analysis and applications,  $247(1)$ , 45-66.
- <span id="page-52-2"></span>Glazer, J., Kremer, I., & Perry, M. (2021). The wisdom of the crowd when acquiring information is costly. Management Science,  $67(10)$ , 6443–6456.
- <span id="page-52-10"></span>Hörner, J., & Skrzypacz, A. (2017). Learning, experimentation and information design. Advances in Economics and Econometrics, 1, 63-98.
- <span id="page-52-9"></span>Immorlica, N., Mao, J., Slivkins, A., & Wu, Z. S. (2019). Bayesian exploration with heterogeneous agents. In *The world wide web conference* (pp. 751–761).
- <span id="page-52-5"></span>Kamenica, E., & Gentzkow, M. (2011). Bayesian persuasion. American Economic Review,  $101 (6)$ , 2590-2615.
- Keller, G., Rady, S., & Cripps, M. (2005). Strategic experimentation with exponential bandits. *Econometrica*,  $73(1)$ , 39–68.
- <span id="page-52-13"></span>Klenke, A. (2020). Probability theory: a comprehensive course. Springer Cham.
- <span id="page-52-3"></span>Komiyama, J., & Noda, S. (2021). Deviation-based learning. arXiv preprint arXiv:2109.09816 .
- <span id="page-52-0"></span>Kremer, I., Mansour, Y., & Perry, M. (2014). Implementing the "wisdom of the crowd". Journal of Political Economy, 122 (5), 988–1012.
- <span id="page-52-4"></span>Küçükgül, C., Özer, Ö., & Wang, S. (2022). Engineering social learning: Information design of time-locked sales campaigns for online platforms. Management Science,  $68(7)$ , 4899–4918.
- <span id="page-53-0"></span>Liang, Y., Sun, P., Tang, R., & Zhang, C. (2020). Efficient resource allocation contracts to reduce adverse events (Tech. Rep.). Working Paper.
- <span id="page-53-8"></span>Lorecchio, C. (2021). Pursuading crowds. unpublished manuscript.
- <span id="page-53-1"></span>Lorecchio, C., & Monte, D. (2021). Dynamic information design under constrained communication rules. American Economic Journal: Microeconomics.
- <span id="page-53-13"></span>Luenberger, D. G. (1997). Optimization by vector space methods. John Wiley & Sons.
- <span id="page-53-3"></span>Mansour, Y., Slivkins, A., & Syrgkanis, V. (2020). Bayesian incentive-compatible bandit exploration. Operations Research,  $68(4)$ , 1132–1161.
- <span id="page-53-9"></span>Mansour, Y., Slivkins, A., Syrgkanis, V., & Wu, Z. S. (2016). Bayesian exploration: Incentivizing exploration in bayesian games.  $arXiv$  preprint  $arXiv:1602.07570$ .
- Martin, K., Ryan, C. T., & Stern, M. (2016). The slater conundrum: duality and pricing in infinite-dimensional optimization. SIAM Journal on Optimization,  $26(1)$ , 111–138.
- <span id="page-53-12"></span>Müller, A., & Stoyan, D.  $(2002)$ . Comparison methods for stochastic models and risks (Vol. 389). Wiley.
- <span id="page-53-10"></span>Myerson, R. B. (1986). Multistage games with communication. Econometrica: Journal of the Econometric Society, 323–358.
- <span id="page-53-7"></span>Orlov, D., Skrzypacz, A., & Zryumov, P. (2020). Persuading the principal to wait. Journal of Political Economy, 128(7), 2542-2578.
- <span id="page-53-2"></span>Papanastasiou, Y., Bimpikis, K., & Savva, N. (2018). Crowdsourcing exploration. Management Science, 64 (4), 1727–1746.
- <span id="page-53-4"></span>Rayo, L., & Segal, I. (2010). Optimal information disclosure. Journal of Political Econ $omy, 118(5), 949-987.$
- <span id="page-53-5"></span>Renault, J., Solan, E., & Vieille, N. (2017). Optimal dynamic information provision. Games and Economic Behavior, 104, 329–349.
- <span id="page-53-14"></span>Shaked, M., & Shanthikumar, J. G. (2007). Stochastic orders. Springer.
- <span id="page-53-11"></span>Smith, L., & Tian, J. (2018). Informational inertia. unpublished mimeo.
- <span id="page-53-6"></span>Smolin, A. (2021). Dynamic evaluation design. American Economic Journal: Microeconomics,  $13(4)$ ,  $300-331$ .